

## Original Research Article

# Evaluation of pigeonpea genotypes against sterility mosaic disease

### ABSTRACT

Pigeonpea (*Cajanuscajan* L.) belonging to family leguminosae is an important leguminous pulse crop of semi-arid tropic and subtropic regions (Asia and Africa). It is also known as red gram, arhar, tur dal. The crop is affected by several phytopathogens, of which, pigeonpea sterility mosaic disease/virus, and transmissible by eriophyid mites (*Aceria-cajani*) has been a major bottleneck. Yield losses up to 95 per cent or even 100 per cent in severe Sterility Mosaic Disease incidence were reported. Therefore, present investigation on screening of pigeon pea genotypes against the sterility mosaic disease was carried out during Kharif, 2018 at the Department of Plant Pathology, College of Agriculture, Latur.

Formatted: Font: Italic

Formatted: Font: Italic

In present study, about 27 entries of pigeonpea were screened under natural epiphytotic against pigeonpea sterility mosaic disease. Of these GRG-152, ICP-2376, BRG-5, BRG-4, ICPL-15048, BSMR853, BSMR-736, BRG-1 and BRG-3 were resistant; MPV-106, RVSA-16-1, IPA-16-8 and BRG-2 were moderately resistant, whereas, PUSA-2017-01, TDRG-58, ICP-8863, PUSA-2018, PUSA-2018-1, PUSA-2018-2, PUSA-2018-3, PUSA-2018-5, AKTE-12-04, KRG-244, PADT-16, RKPV-912, JKM-189 and TJT-501 were susceptible to Sterility Mosaic disease.

**Key words:** Sterility mosaic of disease, Pigeonpea, Screening, Genotypes.

### Introduction:

Pigeonpea (*Cajanuscajan* L.) is an important pulse crop of semi-arid tropic and subtropic regions viz., Asia and Africa. Pigeonpea is grown in India over an area of 5.39 M ha with an annual production of 4.60 M tones and productivity of 854 kg/ha, whereas, in Maharashtra these were 1037.0 ha, 661.0 tonnes and 637 kg/ha, respectively (Anonymous, 2017).

Pigeonpea crop being affected by around 50 diseases/pathogens, in mild to severe form. Of these, Sterility Mosaic (SM), *Fusarium* wilt and *Phytophthora* blight are economically important. SMD causes substantial yield losses up to 95 per cent (Ganapathy et al. 2011).

Comment [CM1]: Add information about the initial identification of the virus in the region. If there is a molecular identification, add Genebank accession numbers

Comment [CM2]: Add more information about viral particles nature

Formatted: List Paragraph, Space Before: 0 pt

Comment [CM3]: Add information about virus-vector relationship

Sterility Mosaic Disease (SMD) caused by Pigeonpea Sterility Mosaic Virus (PPSMV) is a widespread and economically important disease. It is transmitted by the eriophyid mite (*Aceria-cajani*). Sterility Mosaic Disease (SMD), was first described in 1931 from Pusa, Bihar State of India (Mitra, 1931), is a major disease limiting the pigeonpea production in the Indian sub-continent. SMD causes substantial yield losses up to 95 per cent (Ganapathy et al. 2011).

During recent years, research programs have focused on the development of high yielding genotypes with combined resistance to PPSMV and *Fusariumwilt*, as both these diseases are endemic in the subcontinent. Therefore, the present study was carried out with aimed to screen the pigeonpea genotypes against sterility mosaic disease to identify the resistant sources.

**Material and Methods:**

A field experiment was conducted in the research farm at the Department of Plant Pathology, College of Agriculture, Latur during *Kharif* 2018-19. All twenty seven pigeonpea germplasm, cultivars and varieties obtained from different sources were evaluated for their reactions against pigeonpea sterility mosaic disease (SMD). These genotypes were planted in each row with row to row spacing of 60 cm and plant to plant spacing of 15 cm. SMD check (Maruthi) was included after every 10 test rows. Based on disease reactions, 27 test entries of pigeonpea germplasm were categorized as given below, by applying the rating scale of AICRP on Pigeonpea.

**Table 1. Pigeonpea SMD rating scale**

Sr. No.	Disease incidence	Reactions
1	0-10%	Resistant
2	11-30%	Moderately Resistant
3	>30%	Susceptible

Per cent disease incidence (PDI) was calculated by using following formula:

$$PDI = \frac{\text{Number of diseased plants}}{\text{Total number of plant observed}} \times 100$$

**Results and Discussion:**

Twenty seven pigeonpea elite entries were screened against sterility mosaic disease (SMD) under field conditions, at the research farm Department of Plant Pathology, College of Agriculture, Latur. Average per cent incidence of SMD was calculated and accordingly the test entries were categorized.

The result (Table 2) revealed zero disease incidence in ICP-2376, BRG-4, BRG-5, BSMR-853, BRG-1 and BRG-3, followed by ICPL-15048 (2.22%) and BSMR-736 (2.94%). Whereas, highest disease incidence (100%) was reported in ICP-8863, PUSA-2018 and AKTE-12-04, followed by PUSA-2018-2 (97.56%) and PUSA-2018-1 (86.84%).

Based on per cent SMD incidence, the test pigeonpea entries were categorized (Table 3), which revealed the entries viz., GRG-152, ICP-2376, BRG-4, BRG-5, ICPL-15048, BSMR-853, BSMR-736, BRG-1 and BRG-3 as resistant with the

**Comment [CM4]:** Add more information about it. Is a control cv? Resistant or susceptible?

**Comment [CM5]:** Been a field experiment, several other virus can be present, add information if there are no other virus infecting pigeonpea in the region.

**Comment [CM6]:** Add what AICRP means

**Comment [CM7]:** Add how PPSMV was identify. If was based on symptoms or an ELISA, PCR test?

**Comment [CM8]:** Discussion can be more informative

SMD incidence in the range of 0-10 per cent. The entries viz., MPV-106, RVSA-16-1, IPA-16-18 and BRG-2 were moderately resistant (11-30% incidence) and the entries viz., PUSA-2017-01, TDRG-58, ICP-8863, PUSA-2018, PUSA-2018-1, PUSA-2018-2, PUSA-2018-3, PUSA-2018-5, AKTE-12-04, KRG-244, PADT-16, RKPV-912, JKM-189, TJT-501 were susceptible (> 30% incidence).



Healthy plant

Infected sterile plant



Comparison between healthy plant and infected plant

**Comment [CM9]:** Resolution does not permit symptoms identification. I suggest to change this photo

**Comment [CM10]:** Add which ones are the healthy plants in the picture

Various stages of sterility mosaic symptoms development



Bushy and pale green appearance Increase in secondary and tertiary branches

**Comment [CM11]:** Interesting photo. Similar to a phytoplasma infection

**Comment [CM12]:** It more like a vein yellowing than a mosaic symptom



Mosaic pattern of leaves



Mottling of leaves

**Fig 1.** Symptoms produced in experimental field

**Comment [CM13]:** Check number of figures

**Table 2. Screening of pigeonpea genotypes against *Pigeonpea Sterility Mosaic Virus* at the Department of Plant Pathology, College of Agriculture, Latur. 2018-19. PPSMV**

Formatted: Font: Italic, Font color: Auto

Sr. No.	Entries	Total plants	SMD plants	% Disease Incidence	Disease Reactions
1.	GRG-152	44	4	9.09	R
2.	MPV-106	43	12	27.90	MR
3.	PUSA- 2017-01	48	29	60.41	S
4.	RVSA-16-1	38	11	28.94	MR
5.	TDRG-58	39	17	43.58	S
6.	ICP-8863 (S.Check)	44	44	100	S
7.	PUSA-2018	37	37	100	S
8.	PUSA-2018-1	38	33	86.84	S
9.	PUSA-2018-2	41	40	97.56	S
10.	PUSA-2018-3	35	25	71.42	S
11.	PUSA-2018-5	31	19	61.29	S
12.	ICP-2376	42	0	0	R
13.	AKTE-12-04	32	32	100	S
14.	IPA-16-18	30	5	16.66	MR
15.	KRG-244	29	19	65.51	S
16.	PADT-16	30	19	63.33	S
17.	RKPV-912	32	14	43.75	S
18.	BRG-2	38	11	28.94	MR
19.	BRG-4	33	0	0	R
20.	BRG-5	37	0	0	R
21.	ICPL-15048	45	1	2.22	R
22.	JKM-189	39	12	30.76	S
23.	TJT-501	33	17	51.51	S
24.	BSMR-853	41	0	0	R
25.	BSMR-736	34	1	2.94	R
26.	BRG-1	36	0	0	R
27.	BRG-3	43	0	0	R

SMD: Sterility mosaic diseased R:Resistant MR:Moderately Resistant S: Susceptible

Comment [CM14]: This information its already presented table 3

**Table 3. Categorization of pigeonpea test entries based on reactions to SMD disease, during Kharif 2018-19.**

Comment [CM15]: This information its already presented in the first two tables. I suggest to delete.

Sr. No.	Disease Reactions	List of entries
1.	Resistant (0-10%) Entries= 09	GRG-152, ICP-2376, BRG-4, BRG-5, ICPL-15048, BSMR-853, BSMR-736, BRG-1, BRG-3
2.	Moderately resistant (11-30%) Entries= 04	MPV-106, RVSA-16-1, IPA-16-18, BRG-2
3.	Susceptible (>30%) Entries= 14	PUSA-2017-01, TDRG-58, ICP-8863, PUSA-2018, PUSA-2018-1, PUSA-2018-2, PUSA-2018-3, PUSA-2018-5, AKTE-12-04, KRG-244, PADT-16, RKPV-912, JKM-189, TJT-501

These results on the reactions of pigeonpea entries against SMD are reported on similar line to the reports of several earlier workers such as Manjunatha *et al.* (2013), Barhate *et al.* (2000), Bhaskar (2016), Sudharani *et al.* (2017) and

Roy and Kumar (2018).Manjunatha *et al.* (2013) reported BRG-3 as resistant to SMD with high seed yield, BRG-2 as moderately resistant and ICP-8863 as susceptible.Barhate *et al.* (2000) reported 100 per cent incidence (100%) of sterility mosaic in cv.ICP-8863. Bhaskar (2016) reported BSMR-736 and BSMR-853 as resistant to sterility mosaic, BRG-2 as moderately resistant and ICP-8863 as susceptible to sterility mosaic disease. Sudharani *et al.* (2017) reported cv.GRG-152 as moderately resistant to SMD, Roy and Kumar (2018) reported PADT-16, ICP-8863, TDRG-58 and JKM-189 as susceptible to sterility mosaic disease.

### Conclusion:

These results on the reactions of pigeonpea entries against SMD are nine genotypes are resistant, four are moderately resistant and fourteen were susceptible to the pigeonpea sterility mosaic disease. Need further investigation for resistant breeding.

### References:

- Anonimus (2017). *Agricultural Statistics at a glance*, Agricultural statistics division, Ministry of Agriculture, Government of India, New Delhi: 252.
- Barhate, B. G., Bendre, N. J., Kute, N. S. and Gaikwad, R. T. (2000). Sources of resistance to *Fusarium* wilt and sterility mosaic disease of pigeonpea. *Legume Res.*, **23**(2):136-138.
- Bhaskar Vijay (2016). Screening of pigeonpea genotypes against wilt and sterility mosaic disease in Telangana State, India. *Indian J. Agric. Res.*, **50**(2): 172-176.
- Ganapathy, K. N., Gnanesh, B. N., Byregowda, M., Venkatesha, S. C., Sunil, S., Gomashe and Channamallikarjuna, V. (2011). AFLP analysis in pigeonpea (*Cajanus cajan* (L.) Millsp.) revealed close relationship of cultivated genotypes with some of its wild relatives. *Genet. Res. Crop Evol.*, **58**: 837-847.
- Manjunatha, L., Ramappa, H. K., Gowda, M. B., Rangaswamy, K. T. and Nagaraju (2013). Prevalence of sterility mosaic disease in Karnataka and its resistance source. *Mysore J. Agric. Sci.*, **47**(2): 314-319.
- Mitra, M. (1931). Report of the Imperial Mycologist, *Scientific Rep. Agric. Res. Inst., Pusa* **19**: 58-71.
- Nene, Y. L., Reddy, M. V., Beniwal, S. P. S., Mahmood, M., Zote, K. K., Singh, R. N. and Sivaprakasam, K. (1989). Multilocal testing of pigeonpea for broad-based resistance to sterility mosaic in India. *Indian Phytopath.*, **42**(3): 444-448.
- Roy, A. N. and Kumar, B. (2018). Evaluation of pigeonpea genotypes against sterility mosaic disease under artificial inoculation conditions. *Pl. Dis. Res.*, **33**(2): 208-212.

**Comment [CM16]:** Since there are no serological or molecular identification of the virus, incidence were calculated based of the presence of symptoms. However, asymptomatic infection are common in apparently resistant cultivars. So breeding of this materials are not recommended. This information it's important for other scientists working on this subject

Sudharani, Y. S., Amaresh, S. M. and Naik, M. K. (2017). Evaluation of pigeonpea (*Cajanuscajan* L. Millspaugh) genotypes against sterility mosaic disease. *Internat. J. Pure App. Biosci.*,5(4): 1841-1844.

UNDER PEER REVIEW