

Evaluation of pigeonpea genotypes against sterility mosaic disease

ABSTRACT

Pigeonpea (*Cajanus cajan* L.) belonging to family Leguminosae is an important leguminous pulse crop of semi-arid tropic and subtropic regions (Asia and Africa). It is also known as red gram, arhar, tur dal. The crop is affected by several phytopathogens, of which, pigeonpea sterility mosaic disease/virus, and transmissible by eriophyid mites (*Aceria cajani*) has been a major bottleneck. Yield losses up to 95 per cent or even 100 per cent in severe Sterility Mosaic Disease incidence were reported. Therefore, present investigation on screening of pigeon pea genotypes against the sterility mosaic disease was carried out during Kharif, 2018 at the Department of Plant Pathology, College of Agriculture, Latur.

In present study, about 27 entries of pigeonpea were screened under natural epiphytotic against pigeonpea sterility mosaic disease. Of these GRG-152, ICP-2376, BRG-5, BRG-4, ICPL-15048, BSMR853, BSMR-736, BRG-1 and BRG-3 were resistant; MPV-106, RVSA-16-1, IPA-16-8 and BRG-2 were moderately resistant, whereas, PUSA-2017-01, TDRG-58, ICP-8863, PUSA-2018, PUSA-2018-1, PUSA-2018-2, PUSA-2018-3, PUSA-2018-5, AKTE-12-04, KRG-244, PADT-16, RKPV-912, JKM-189 and TJT-501 were susceptible to Sterility Mosaic disease.

Key words: Sterility mosaic of disease, Pigeonpea, Screening, Genotypes.

Introduction:

“Pigeonpea (*Cajanus cajan* L.) is an important pulse crop of semi-arid tropic and subtropic regions viz., Asia and Africa. Pigeonpea is grown in India over an area of 5.39 M ha with an annual production of 4.60 M tones and productivity of 854 kg/ha, whereas, in Maharashtra these were 1037.0 ha, 661.0 tones and 637 kg/ha, respectively” (Anonymous, 2017).

“Pigeonpea crop being affected by around 50 diseases/pathogens, in mild to severe form. Of these, Sterility Mosaic Disease (SMD) caused by Pigeonpea Sterility Mosaic Virus (PPSMV) is a widespread and economically important disease. It is transmitted by the eriophyid mite (*Aceria cajani*). Sterility Mosaic Disease (SMD), was first described in 1931 from Pusa, Bihar State of India, is a major disease limiting the pigeonpea production in the Indian sub-continent”. (Mitra, 1931) “SMD causes substantial yield losses up to 95 per cent” (Ganapathy et al. 2011). “The vector, eriophyid mite, is host specific with a narrow host range confined to pigeonpea and few of its wild relatives” (Kumar et al., 2003). “It is the sole vector responsible for the transmission of SMD in pigeonpea” (Kulkarni et al., 2002 and Patil and Kumar, 2015)

PSMD is the most destructive disease of pigeonpea (Kannaiyan et al., 1984) causing yield losses up to 95 percent (Ganapathy et al., 2011; Reddy and Nene, 1981). The early stage (<45-days old plants) of infection results into 95 to 100

percent yield losses as reported by Reddy et al. (1990); Kulkarni et al. (2003). The disease results in 100 percent yield loss when symptoms appear at the pre-flowering and podding stage. Whereas, at maturity stage loss could be of 67 percent and at pre-harvest stage up to 30 percent.

“Manifestation of PSMD chiefly depends on the availability of mite populations” (Singh et al., 1999). During recent years, research programs have focused on the development of high yielding genotypes with combined resistance to PPSMV and *Fusarium*wilt, as both these diseases are endemic in the subcontinent. Therefore, the present study was carried out with aimed to screen the pigeonpea genotypes against sterility mosaic disease to identify the resistant sources.

Material and Methods:

A field experiment was conducted in the research farm at the Department of Plant Pathology, College of Agriculture, Latur during *Kharif* 2018-19. All twenty-seven pigeonpea germplasms, cultivars and varieties obtained from different sources were evaluated for their reactions against pigeonpea sterility mosaic disease (SMD). These genotypes were planted in each row with row to row spacing of 60 cm and plant to plant spacing of 15 cm SMD susceptible check (variety Maruthi) was included after every 10 test rows. The symptoms showing sterility mosaic were recorded and disease incidence were recorded. Based on disease reactions to sterility mosaic ,pigeonpea germplasms were categorized as given below, by applying the rating scale of All India Coordinated Research Project on Pigeonpea.

Table 1. Pigeonpea Sterility Mosaic Disease rating scale

Sr. No.	Disease incidence	Reactions
1	0-10%	Resistant
2	11-30%	Moderately Resistant
3	>30%	Susceptible

Based on symptoms of sterility mosaic disease in field disease incidence was recorded. Per cent disease incidence (PDI) was calculated by using following formula:

$$\text{PDI} = \frac{\text{Number of diseased plants}}{\text{Total number of plants observed}} \times 100$$

Results and Discussion:

Twenty-seven pigeonpea elite entries were screened against sterility mosaic disease (SMD) under field conditions, at the research farm Department of Plant Pathology, College of Agriculture, Latur. Average per cent incidence of SMD was calculated and accordingly the test entries were categorized.

The symptoms of sterility mosaic of pigeonpea observed during screening were bushy and pale green appearance of plants followed by reduction in size, increase in number of secondary branches and mosaic mottling of leaves and finally partial or complete cessation of reproductive structures. In some plants affected plants remain green with more vegetative growth and have no flower or seed pods (Fig.1 and 2a and b). The result (Table 1) revealed zero disease incidence in ICP-2376, BRG-4, BRG-5, BSMR-853, BRG-1 and BRG-3, followed by ICPL-15048 (2.22%) and BSMR-736 (2.94%). Whereas, highest disease incidence (100%) was reported in ICP-8863, PUSA-2018 and AKTE-12-04, followed by PUSA-2018-2 (97.56%) and PUSA-2018-1 (86.84%).

Based on per cent SMD incidence, the test pigeonpea entries were categorized, which revealed the entries viz., GRG-152, ICP-2376, BRG-4, BRG-5, ICPL-15048, BSMR-853, BSMR-736, BRG-1 and BRG-3 as resistant with the SMD incidence in the range of 0-10 per cent. The entries viz., MPV-106, RVSA-16-1, IPA-16-18 and BRG-2 were moderately resistant (11-30% incidence) and the entries viz., PUSA-2017-01, TDRG-58, ICP-8863, PUSA-2018, PUSA-2018-1, PUSA-2018-2, PUSA-2018-3, PUSA-2018-5, AKTE-12-04, KRG-244, PADT-16, RKPV-912, JKM-189, TJT-501 were susceptible (> 30% incidence).



Infected sterile plant

Healthy Plant

Fig 1. Comparison between healthy plant and infected plant



Fig 2a. Various stages of sterility mosaic symptoms development



and pale green appearance



Increase in secondary and tertiary branches

Bushy



Mosaic pattern of leaves



Mottling of leaves

Fig 2b.Symptoms produced in experimental field

Table 2.Screening of pigeonpea genotypes against *Pigeonpea Sterility Mosaic Virus*disease.

Sr. No.	Entries	Total plants	SMD plants	% Disease Incidence	Disease Reactions
1.	GRG-152	44	4	9.09	R
2.	MPV-106	43	12	27.90	MR
3.	PUSA- 2017-01	48	29	60.41	S
4.	RVSA-16-1	38	11	28.94	MR
5.	TDRG-58	39	17	43.58	S
6.	ICP-8863 (S.Check)	44	44	100	S
7.	PUSA-2018	37	37	100	S
8.	PUSA-2018-1	38	33	86.84	S
9.	PUSA-2018-2	41	40	97.56	S
10.	PUSA-2018-3	35	25	71.42	S
11.	PUSA-2018-5	31	19	61.29	S
12.	ICP-2376	42	0	0	R
13.	AKTE-12-04	32	32	100	S
14.	IPA-16-18	30	5	16.66	MR
15.	KRG-244	29	19	65.51	S
16.	PADT-16	30	19	63.33	S
17.	RKPV-912	32	14	43.75	S
18.	BRG-2	38	11	28.94	MR
19.	BRG-4	33	0	0	R

20.	BRG-5	37	0	0	R
21.	ICPL-15048	45	1	2.22	R
22.	JKM-189	39	12	30.76	S
23.	TJT-501	33	17	51.51	S
24.	BSMR-853	41	0	0	R
25.	BSMR-736	34	1	2.94	R
26.	BRG-1	36	0	0	R
27.	BRG-3	43	0	0	R

SMD: Sterility mosaic diseased R:Resistant MR:Moderately Resistant S: Susceptible

These results on the reactions of pigeonpea entries against SMD are reported on similar line to the reports of several earlier workers such as Manjunatha *et al.* (2013), Barhate *et al.* (2000), Bhaskar (2016), Sudharani *et al.* (2017) and Roy and Kumar (2018), Tharageshwari *et al.* (2019) Dhanushasree *et al.* (2019). Manjunatha *et al.* (2013) reported BRG-3 as resistant to SMD with high seed yield, BRG-2 as moderately resistant and ICP-8863 as susceptible. Barhate *et al.* (2000) reported 100 per cent incidence (100%) of sterility mosaic in cv. ICP-8863. Bhaskar (2016) reported BSMR-736 and BSMR-853 as resistant to sterility mosaic, BRG-2 as moderately resistant and ICP-8863 as susceptible to sterility mosaic disease. Sudharani *et al.* (2017) reported cv. GRG-152 as moderately resistant to SMD, Roy and Kumar (2018) reported PADT-16, ICP-8863, TDRG-58 and JKM-189 as susceptible to sterility mosaic disease. According to Tharageshwari *et al.* (2019), out of the ninety-four genotypes studied, only four genotypes, DPP 2-89, DPP 3-182, IC 22557, and ICP 3666 showed highly resistant reaction to SMD infection, whereas fifty-four genotypes showed highly sensitive reaction. Genotypes viz., CRG 16-07, BWR 153, ICP 7919, IC 339057, IC74016, IPAE 15-05, AL 2250, CRG 16-01, Pusa Arhar 21-14, Pusa Arhar 21-27, BWR 253, ICP 9808 and ICP 7234 were found to be SMD resistant ones and can be utilized as donors for resistant breeding program to reduce yield loss as compared to susceptible types.

Conventional plant breeding methods have been effective in bringing about improvement in crops but efforts are still being made to develop more efficient breeding methods to overcome specific problems.

Conclusion:

These results on the reactions of pigeonpea entries against SMD are nine genotypes are resistant, four are moderately resistant and fourteen were susceptible to the pigeonpea sterility mosaic disease.

References:

Anonymus (2017). *Agricultural Statistics at a glance*, Agricultural statistics division, Ministry of Agriculture, Government of India, New Delhi: 252.

Barhate, B. G., Bendre, N. J., Kute, N. S. and Gaikwad, R. T. (2000). Sources of resistance to *Fusarium* wilt and sterility mosaic disease of pigeonpea. *Legume Res.*, 23(2):136-138.

- Bhaskar Vijay (2016). Screening of pigeonpea genotypes against wilt and sterility mosaic disease in Telangana State, India. *Indian J. Agric. Res.*,50(2): 172-176.
- Dhanushasree M., A. Thanga Hemavathy, R.P. Gnanamalar and L. Karthiba (2022). Evaluation of Pigeonpea Genotypes against Sterility Mosaic Virus (PPSMV) Disease. *Biological Forum – An International Journal*, 14(2): 1263-1268.
- Ganapathy, K. N., Gnanesh, B. N., Byregowda, M., Venkatesha, S. C., Sunil, S., Gomashe and Channamallikarjuna, V. (2011). AFLP analysis in pigeonpea (*Cajanuscajan* (L.) Millsp.) revealed close relationship of cultivated genotypes with some of its wild relatives. *Genet. Res. Crop Evol.*,58: 837-847.
- Kannaiyan J, Nene YL, Reddy MV, Ryan G, Raju T N.(1984) Prevalence of pigeonpea disease and associated crop losses in Asia Africa and America. *Trop Pest Manag.* 30:62–71.
- Kulkarni NK, Reddy AS, Kumar PL, Vijayanarasimha J, Rangaswamy KT, Muniyappa V, Reddy LJ, Saxena KB, Jones AT, Reddy DV. (2003). Broad based resistance to pigeonpea sterility mosaic disease in accessions of *Cajanus scarbaeoides*(L.) Benth. *Ind J Plant Pro.* 31:6–11.
- Kumar, P. L., Jones, A. T., and Reddy, D. V. R. (2003). A novel mite-transmitted virus with a divided RNA genome closely associated with pigeonpea sterility mosaic disease. *Phytopathology* 93, 71–81.
- Patil, B. P., and Kumar, P. L. (2015). *Pigeonpea sterility mosaic virus*: a legume infecting Emaravirus from South Asia. *Mol. Plant Pathol.* 16, 775–786.
- Manjunatha, L., Ramappa, H. K., Gowda, M. B., Rangaswamy, K. T. and Nagaraju (2013).Prevalence of sterility mosaic disease in Karnataka and its resistance source.*Mysore J. Agric. Sci.*,47(2): 314-319.
- Mitra, M. (1931). Report of the Imperial Mycologist, *Scientific Rep.Agric.Res.Inst.,Pusa*19: 58-71.
- Nene, Y. L., Reddy, M. V., Beniwal, S. P. S., Mahmood, M., Zote, K. K., Singh, R. N. and Sivaprakasam, K. (1989). Multilocational testing of pigeonpea for broad-based resistance to sterility mosaic in India.*Indian Phytopath.*,42(3): 444-448.
- Reddy MV, Nene YL.1981 Estimation of yield loss in pigeonpea due to sterility mosaic *Proc. International workshop on Pigeonpeas* 215–19 December 1980 ICRI SAT Center; Patancheru, AP, India: 305–312.
- Reddy MV, Sharma SB, Nene YL. (1990). Pigeonpea: Disease management. In: Nene YL, Hall SD, Sheila VK, editors. *The Pigeonpea*. CAB International, Wallingford; UK. pp. 303–347.
- Roy, A. N. and Kumar, B. (2018). Evaluation of pigeonpea genotypes against sterility mosaic disease under artificial inoculation conditions.*Pl. Dis. Res.*, 33(2): 208-212.
- Sudharani, Y. S., Amaresh, S. M. and Naik, M. K. (2017). Evaluation of pigeonpea (*Cajanuscajan* L. Millspaugh) genotypes against sterility mosaic disease. *Internat. J. Pure App. Biosci.*,5(4): 1841-1844.

Singh AK, Rathi YPS, Agrawal KC. (1999). Sterility mosaic of pigeonpea: a challenge of the 20th century. *Indian J Virol.* 15:85-92.

Tharageshwari, L. M., Hemavathy, A. T., Jayamani, P. and Karthiba, L. (2019). Evaluation of pigeonpea (*Cajanus cajan*) genotypes against pigeonpea sterility mosaic disease. *Electronic Journal of Plant Breeding*, 10(2): 727-731.

UNDER PEER REVIEW