

Sustainable development attributes of zero budget natural farming (ZBNF) to agricultural practices

Abstract

The agricultural sector is fundamental to human civilization since it provides food and other basic needs to billions of people. Given that 9.7 billion people are expected to live on Earth by 2050, food production needs to increase by 70%. The population is predicted to become more urbanized by 2050, with 75% of people likely to live in metropolitan regions. Improved agricultural techniques and the promotion of high-yielding cultivars are essential to addressing the issues of sustainability and food security. The Green Revolution changed agriculture in India, resulting in a food surplus. However, it has resulted in difficulties such as lower crop output, resource depletion, and environmental deterioration. In response, new techniques such as zero-budget natural farming (ZBNF) have evolved. Zero Budget Natural Farming (ZBNF) is a sustainable farming approach aimed at improving soil health, increasing crop output, and lowering production costs. The foundations of ZBNF are examined in this review study along with how they affect agricultural output, environmental sustainability, and socioeconomic advantages. ZBNF places a strong emphasis on using organic inputs and natural resources such as *Jeevamrita*, *bijamrita*, *acchadana* and *whapasato* to increase microbial diversity, improve soil fertility, encourage nutrient cycling, and save water. ZBNF can lower production costs, boost crop yield, and enhance soil health. It also confers better emergence parameters, relative leaf water content, root parameters, and overall yield and productivity on plant growth. Additionally, ZBNF supports the UN Sustainable Development Goals by encouraging sustainable agriculture, lowering greenhouse gas emissions, and improving rural lives.

Keywords – Zero budget natural farming (ZBNF), sustainable agriculture, soil health, crop productivity, microbial diversity

Introduction

Agricultural sector is the major foundation of human civilization which provides food and essentials to billions of people on earth. The World's population is projected to hike about 9.7 billion people by 2050, and therefore, the food production levels need to be increased by 70 per cent to ensure the agricultural productivity and to enhance sustainability (Shubham *et al.*, 2023). Furthermore, 75% of the World's population is expected to reside in urban areas by 2050 due to the World's growing urbanization (Anonymous, 2018). In order to meet the everlasting demands for food safety and security, ensuring food security becomes a major concern for the world. Adoption of high yielding cultivars and improved farming practices with sound technologies can be an alternative for overcoming such issues (Shubham *et al.*, 2022). Green revolution technology has proven itself a double-edged sword for both the Indian agricultural sectors and worldwide. However, Indian agricultural sector has progressed from a food-scarce to a food-surplus yet although it has faced various challenges like reduction in crop factor productivity, depletion of natural resources, lack of water and nutrients and negative implications on climate change and soil health (Bharucha *et al.*, 2020).

In the last decade, various crop management approaches like biofertilizers/biopesticides, vermicompost and farm yard manure (FYM) have been used

to reduce the detrimental effects of chemical fertilizers and pesticides on soil (Ray *et al.*, 2020). Despite such scientific attempts, unfortunately there is still no established strategy for replacing the chemical-based fertilizers with long-term negative outcomes. Moreover, organic farming reduced the greenhouse gases (GHG's) emissions by about 65 per cent and mitigated the climate change. However, challenges such as low production and complex certification procedure still remain major hurdles (Timsina, 2018). Zero budget natural farming (ZBNF), a historic and traditional farming approach has been reintroduced as a self-sufficient alternative to assist and to address these concurrent concerns. ZBNF was promoted by agriculturist Mr. Subhash Palekar in the mid-1990's as an alternative to green revolution's methods based on chemical fertilizers and pesticides and intensive irrigation. ZBNF has emerged as a sustainable and cost-effective agricultural practice that prioritizes soil health, biodiversity and farmers well-being. In the face of increasing global food demand and environmental challenges, the adoption of ZBNF offers a promising solution to enhance the agricultural productivity along with preserving the natural resources. The approach of ZBNF relies on the exemption of external sources like synthetic fertilizers and pesticides and therefore cutting the cost to almost zero. Additionally, crops are grown without the use of chemicals, utilizing natural resources like cow dung and urine among others (Bharucha *et al.*, 2020; Timsina, 2018; Khadse *et al.*, 2021). Activity of earthworms and microbial populations are found to be increased many times with the ZBNF formulations like *jeevamrita*, *bijamarita* and *panchgavya*. Incorporation of these formulations in soil boosts the soil nutrients availability, fortifies the resistance mechanism and ultimately raises the crop yield (Ray *et al.*, 2020; Maduka *et al.*, 2019; Patelet *et al.*, 2021).

Recent research on ZBNF by Niti Ayog in the Indian states such as Andhra Pradesh, Maharashtra and Karnataka showed that comparing the irrigated and rainfed conditions, the overall cost of cultivation was found to be dropped by 23.7 %, however the yield, gross and net revenue was improved from 14.2 to 50 % (Kumar *et al.*, 2020). Therefore, after such a reliable result, number of states including Himachal Pradesh, Uttar Pradesh, Haryana, Gujarat, Tamil Nadu and Kerala are shifting towards this farming approach (Khadse and Rosset, 2021). ZBNF is likely to handle sustainability, resource conservation, reduce global warming and other global concerns (Munster, 2018). Therefore, it presents itself as an effective solution to the serious and major problems of modern agriculture era, and in addition, providing a practical and affordable strategy which aims to increase crop productivity without harming the environment. Therefore, the present review mainly explores the pillars of ZBNF and its impacts on agricultural production, environmental and socio-economic benefits.

Zero budget natural farming (ZBNF)

ZBNF approach was introduced by Mr. Subhash Palekar. His keen interest in field experiments showed that how constant chemical usage is affecting the soil fertility and turning the area barren (Jannoura *et al.*, 2014). As the name implies, ZBNF is a farming practice in which there is no expenditure oriented for the cultivation and harvesting of plants. Therefore, it means farmers will save money on fertilizers and insecticides in order to

keep their crops healthy. ZBNF approach reduces the amount of money farmers have to invest. It also helps to prevent the soil from degradation and desertification. It aims to encourage traditional indigenous techniques which lessen the demand for foreign inputs. It focuses largely on on-farm biomass recycling with a particular emphasis on biomass mulching, on-farm formulations of cow dung-urine, frequent soil aeration and the avoidance of synthetic chemicals. In essence, ZBNF is entirely dependent on the exclusive use of numerous natural resources, just as ancient agriculture was. As ZBNF improves soil health through biodiversity, microbial activities, nutrient recycling and beneficial biological interactions which leads to sustainable crop production, it is becoming more popular at a time when production costs are rising sharply, production rates are stagnating and environmental footprints associated with chemical pesticides and fertilizers are increasing (Moss and Bittman, 2018).

This approach has the potential to boost the rural economies and lowering the farmer's credit risks associated with chemical farming, which uses a lot of resources. Additionally, it will enable farming households to save more money for necessities like financial stability, healthcare and education. The Andhra Pradesh government gave Rythu Sadhikara Samstha (RySS), a state-owned nonprofit organization, an order in 2015 to expand ZBNF practices to encompass all six million farmers and eight million hectares of agricultural land in the state by 2024. The initiative seeks to support ecological, chemical-free and climate-resilient agriculture in the face of agrarian distress and to give small and marginal farmers lucrative agricultural incomes. The initiative was introduced in 2015–16 and in 2016–17 and began to be implemented in the field. In the 13 districts of Andhra Pradesh, approximately 500,000 farmers had registered for the program, totalling over 2,04,000 acres as of July 2019. Large-scale execution of this initiative might contribute significantly to India's progress toward nearly 25 per cent of the 169 SDG objectives (Tripathi *et al.*, 2018). ZBNF is based on its "four wheels": *Whapasa* (moisture), *Bijamrita* (treating seeds), *Jeevamrita* (inoculating soil) and *Acchadana* (mulching). Two of them are microbial combinations that can be made in less than 48 hours, that is *Jeevamrita* and *Bijamrita* (Khadse and Rosset, 2019). In order to release the soil's natural nutrients through *Jeevamrita*, the notion of "Annapurna" was presented. This idea states that each farm family does not need to possess a cow because one cow's dung and urine are enough to cultivate the thirty acres of land (Anonymous, 2018).

Pillars of ZBNF

- 1. *Jeevamrita*:** A fermented microbial culture which acts as a catalyst and promotes the soil microbial activity, increase the number of native earthworms in soil and also enrich soil by supplying nutrients. For the production of *jeevamrita*, one need to fill the barrel with 200 litres of water, add 10 kg of fresh cow dung from the area, 5 to 10 litres of old cow urine, 2 kilogram of jaggery, a sort of brown sugar from the area, 2 kg of pulse flour and a handful of dirt from the farm's bund. Give the mixture a good stir, then let it to ferment in the shade for 48 hours then, *jeevamrita* is ready for use. One acre of land may be sufficiently covered with 200 litres of *jeevamrita*. Over the course of the 48-hour fermentation process, the anaerobic and aerobic bacteria present in cow dung and urine proliferate as they consume organic ingredients such as pulse flour and jaggery. Native species of bacteria and organisms are inoculated by a little amount of undisturbed soil. Additionally, *jeevamrita* aids in the prevention of bacterial and fungal plant diseases. It should be applied to the crops twice a month using irrigation water or a 10 per cent foliar spray. The preparation is kept in storage

for a maximum of 15 days. *Jeevamrita* is administered to the individual plant in horticultural crops. Most of the sample farmers in Maharashtra are using the drip irrigation method to apply *jeevamrita*. Using *jeevamrita* enhanced agricultural yield and encouraged the growth of beneficial soil microorganisms (Shaikh and Gachande, 2015).

2. ***Bijamrita***: *Bijamrita* is a locally produced microbial seed treatment suitable for seeds, seedlings and other planting materials. It is made using components similar to those found in *jeevamrita*. It is effective in preventing fungal growth on early roots and protecting seedlings from diseases spread by seeds or the soil (Khadse *et al.*, 2017). It helps to stimulate plant development in addition to acting as a plant protector. The same elements used to make *jeevamrita* are used in its preparation: lime, water, soil, and locally produced Indian cow dung and urine. In particular, five litres of cow urine and 5 kg of cow dung are put into a container that holds 20 litres of water. After that, a fistful of local soil and 50 grams of lime are thoroughly combined with it. Before the seed is placed in the ground, it has to be soaked in *bijamrita*, covered, and planted (Anonymous, 2016).
3. ***Acchadana***: A layer of *acchadana* (mulching) is applied to the soil create optimal ecosystem necessary for the optimal growth, reproduction and activity of *jeevamrita*'s helpful bacteria. Three methods are used to supply this environment: soil mulching, live mulching and mulching with straw. Mulching is a useful technique for raising crop quality and production because it keeps the soil wet, regulates temperature, suppress weeds and reduces evaporation (Chakraborty *et al.*, 2008).

Types of mulching

- a) **Soil Mulching**: This kind of mulching is intended to prevent tilling from destroying topsoil during farming by enhancing aeration and water retention in the soil. It is suggested to avoid deep ploughing.
 - b) **Straw Mulching**: Straw material is the term used to describe dried biomass waste from previous crops. Microbial cultures and the soil biota stimulate the organic materials as they break down and ultimately turn into humus.
 - c) **Live Mulching**: It is essential to construct various cropping patterns for monocotyledons and dicotyledons planted in the same region in order to supply the necessary nutrients to the soil and crops. Plants in the dicot group, which includes pulses, fix nitrogen. Monocots like wheat and rice provide additional elements including phosphate, sulfur and potash.
4. ***Whapasa***: *Whapasa* is the condition where there are both air molecules and water molecules present in the soil which helps in reducing irrigation requirement. Farmers adopting ZBNF who irrigated only at noon and in alternating furrows reported a considerable reduction in irrigation requirements. It lowered dependency on irrigation while improving aeration and soil hydration. Palekar disputes the notion that plant roots need a lot of water. According to him, plant roots require water vapor condensation to improve soil moisture. ZBNE methods which includes mulching can cut water usage by up to 90%, making them suitable for rain-fed farming (Bharucha *et al.*, 2020).

Effect of zero budget natural farming on soil fertility and crop production

ZBNF encourages the use of organic and natural inputs such as agricultural leftovers, cow dung, and urine, as well as other locally accessible resources. By adding necessary minerals to the soil, such as potassium, phosphorus, and nitrogen (NPK), these inputs improve soil fertility. The addition of mulch and other organic inputs raises the amount of organic matter in the soil through organic matter enrichment. As a result, there is an increase in nutrient availability due to improved soil structure, water-holding capacity, nutrient retention, and beneficial microbial activity (Ghosh *et al.*, 2017). Use of *jeevamritha* (fermented microbial culture), cultivates beneficial microorganisms in the soil. This improves microbial variety, which is important for the cycling of nutrients, the prevention of illness, and the general health of the soil. Better soil aeration, nutrient availability, and enhanced soil structure are all facilitated by increased microbial activity in the soil, which results in healthier and more productive soils. Proteobacteria and bacterial phyla including *Bacillus*, *Pseudomonas*, *Rhizobium*, and *Panibacillus* with the application of *jeevamritha* on soil (Saharan *et al.*, 2023). One essential element of ZBNF is inter-cropping, which encourages soil nutrient cycling. By ensuring that vital nutrients are accessible to the crops, this approach helps prevent nutrient imbalances and depletion. The research conducted by Saharan *et al.* (2023) demonstrated a significant increase in available potassium, accessible phosphorus, and organic carbon in the soil increases of up to 46%, 439%, and 142%, respectively. Moreover, there were notable improvements in micronutrients including manganese, iron, copper, and zinc, which increased to 98%, 23%, 62%, and 55%, respectively. ZBNF stimulates nutrient recycling mechanisms in the soil, lowering the need for outside inputs and improving long-term soil fertility by upholding a balanced ecosystem and encouraging biodiversity (Kumar *et al.*, 2019).

One of the four pillars in ZBNF is *acchadana* (mulching), which is frequently done in ZBNF, improves water penetration, lowers evaporation, and controls soil temperature to assist or maintain soil moisture. This guarantees a steady supply of water for crops, particularly in areas where water is scarce. Effective water management techniques, such as rainwater collection and drip irrigation, are promoted in ZBNF to assist save water resources and satisfy the unique moisture requirements of crops without wasting them. Natural farming is a highly effective approach that has been shown to increase water retention capacity. It needs minimal water usage and is known to lessen reliance on water and power. As a result, groundwater reserves will be preserved, water tables will improve, and farmers will have less financial and labor stress. *Whapasa* practices help to improve soil fertility and water use efficiency (Kumar *et al.*, 2020).

ZBNF approach has showed its positive effects on the crop production as it have increased production per unit area in many crops. ZBNF which combines minimum tillage, mulching, and intercropping, shows higher emergence count and emergence velocity. This suggests that ZBNF methods might encourage early plant establishment and improved seed germination (Sreenivasa *et al.*, 2010). Compared to plants grown under conventional tillage practices, plants subjected to ZBNF treatments—including ZBNF combined with minimum tillage (MT), mulching (M), and intercropping (IC)—displayed noticeably higher relative leaf

water content throughout all growth stages. The research conducted by *McMillen*, (2013) suggests that ZBNF techniques may be able to maintain more appropriate water levels in plant tissues, which may result in improved plant performance and health. In research, plants cultivated using Zero Budget Natural Farming (ZBNF) treatments—that is, ZBNF in conjunction with minimum tillage, mulching, and intercropping produced the highest root mass and length densities when compared to plants grown using traditional tillage techniques. This result emphasizes how ZBNF procedures might encourage strong root growth. Plants require a well-developed root system in order to efficiently absorb nutrients and water, which lays the groundwork for healthy development and increased resistance to environmental challenges (*Yang et al.*, 2018). Zero Budget Natural Farming (ZBNF) has become a very promising agricultural strategy that greatly increases yields in comparison to conventional tillage methods, particularly when combined with minimum tillage, mulching, and intercropping. These techniques work in combination to produce a favourable environment for plant development and increased yield. ZBNF gradually contributes to improved soil health by removing artificial inputs and depending only on natural resources. This is a critical improvement since healthier soils hold on more water and nutrients, which promotes healthy root development and overall plant growth (*Parihar et al.*, 2018; *Shao et al.*, 2016).

Impact of zero budget natural farming on sustainable development goals

In addition to potentially helping millions of farmers and their families worldwide, the Zero Budget Natural Farming (ZBNF) effort might make a major contribution to the UN Sustainable Development Goals (SDGs) (Table 1). Chandrababu Naidu, the Andhra Pradesh chief minister, and Erik Solheim, the executive director of the United Nations Environment Programme, announced the results. The research maps the possible social, economic, and environmental impacts of the ZBNF program to the specific goals of the SDGs. The ZBNF farmers in Andhra Pradesh have stopped using artificial pesticides and fertilizers in favor of natural farming practices. Instead, according to the release, they employ inexpensive, naturally produced mixtures, inoculums, and decoctions created from cow dung, cow urine, jaggery, lilac, green chilies, and a number of other similar natural components, all of which are acquired locally.

Table 1. Major practices of ZBNF in line with the UN SDGs.

Sustainable Development Goals	Impact on SDG
SDG 1 (End Poverty)	Considerably reduced input costs and increased yields. Their net earnings are higher as a result.

SDG 2 (End hunger, food security, improved nutrition, and promote sustainable agriculture)	By giving all small-scale food producers fair and safe access to land, we can double agricultural output and profits. Use reliable techniques and make sure food production systems are long-lasting. Funding for rural infrastructure, extension services, and agricultural research should all be increased.
SDG 3 (Health lives and promotes well-being for all)	Encourage mental wellness and well-being while reducing the early death rate from non-communicable diseases by one-third. Cut down on the amount of illnesses and fatalities caused due to contaminated soil, water, and air as well as hazardous components and other aspects.
SDG 4 (Ensure inclusive and equitable quality education)	Increase the percentage of adults and youth who possess the abilities needed for gainful employment, entrepreneurship, and respectable jobs. Make sure that each student has the abilities and information required to promote sustainable growth.
SDG 5 (Gender equality and empower all women and girls)	Reforms to guarantee women's equal access to natural resources, land sovereignty, and economic resources. Increase the way that enabling technology is used to support women's empowerment.
SDG 6 (Ensure availability and sustainable management of water and sanitation for all)	To enhance water quality, decrease untreated wastewater, minimize pollution, reduce hazardous chemical emissions, and boost recycling and safe reuse. Protect and restore aquatic ecosystems.
SDG 7 (Affordable, reliable, sustainable and modern energy for all)	By eliminating the use of chemical pesticides and fertilizers, ZBNF will drastically reduce the amount of energy required along their value chain.
SDG 8 (Encourage inclusive, long-term economic growth and the creation of employment)	Maximizing the effective use of resources in both production and consumption, and keeping environmental damage apart from growth. Ensure that everyone, especially the youth, is fully and productively employed and doing high-quality work.
SDG 9 (Reliable infrastructure, encourage equitable and environmentally friendly development)	Businesses should adapt resource-saving strategies and environmentally friendly practices, and infrastructure should be

	upgraded, in order to make them more sustainable.
SDG 10 (Reduce inequality within and among countries)	Aiming to achieve and maintain income growth rates for the poorest 40% of the population that are higher than the national average
SDG 11 (Cities and human settlement inclusive, safe, resilient, and sustainable)	Limit the number of people affected and the financial losses brought on by catastrophes, especially those involving water, with an emphasis on the weak and disadvantaged. Compared to non-ZBNF plots, ZBNF farmlands could be more resilient to flooding, severe winds, and droughts.
SDG 12 (Ensure sustainable consumption and production patterns)	Ensure sustainable and effective management of natural resources. Achieve environmentally sound waste management for all wastes, including chemicals, and restrict their release into the air, water, and soil in order to minimize any unfavorable consequences. Use prevention, reduction, recycling, and reuse to lessen the amount of waste produced. Promote the application of moral public procurement procedures. Make sure everyone is aware of the environment and knows how to live sustainably.
SDG 13 (Take urgent action to combat climate change and its impacts)	Strengthen the ability and resilience of the world to withstand natural catastrophes and risks associated with climate change.
SDG 14 (Seas, oceans, and marine resources should be preserved and used responsibly for sustainable development.)	Prevent and minimize marine pollution, especially that which is brought on by activities that are done on land, such as the contamination of nutrients and debris. Take steps that will mitigate the impacts of ocean acidification.
SDG 15 (Combat desertification, preserve, restore, and encourage the sustainable use of terrestrial ecosystems, manage forests responsibly, stop and reverse land degradation, and stop the loss of biodiversity.)	Ensure that wetlands, mountains, drylands, forests, and other terrestrial and inland freshwater ecosystems are preserved, restored, and utilized sustainably. Prevent desertification and restore land and soil that have been harmed by floods, droughts, and other natural calamities. Minimize the destruction of natural area habitats, stop the decline in biodiversity, and prevent the extinction of fragile species.

SDG 16 (Encourage open and peaceful societies for long-term growth, guarantee everyone's access to justice, and create inclusive, responsible, and successful institutions at all levels.)	Ensures that decision-making is responsive, inclusive, participatory, and representational at all levels.
SDG 17 (Increase the Global Partnership for Sustainable Development's implementation strategies and resuscitate it.)	Establishing multistakeholder partnerships that mobilize and share knowledge, expertise, technology, and other resources helps improve international cooperation for sustainable development.

Source: (Tripathi *et al.*, 2018)

Conclusion

In conclusion, this review highlights Zero Budget Natural Farming (ZBNF) as a transformative agricultural system with significant implications for sustainability and resilience. ZBNF promotes soil health and an ecosystem that flourishes in agricultural settings by prioritizing the use of natural resources such as *jeevamrita*, *bijamrita*, *acchadana*, and *whapasa*, as well as organic inputs. The emphasis on soil aeration, on-farm biomass recycling, and the avoidance of synthetic pesticides demonstrates ZBNF's commitment to regenerative measures that promote soil fertility and productivity over time. Additionally, ZBNF's cost-effectiveness allows farmers to save money on pesticides and fertilizers which helps both individual farmers and the greater goal of sustainable agriculture. Due to its capacity to reduce production costs, boost crop yield, and enhance overall productivity, ZBNF seems to be able to address food security and environmental sustainability. ZBNF represents an integrated strategy to agricultural development that prioritizes the welfare of the land and people by advocating for resource conservation, greenhouse gas emissions reduction, and rural life enhancement. Further, ZBNF is compatible to the UN Sustainable Development Goals; demonstrates its ability to encourage progress in a broad range of areas, including poverty reduction, climate action, and sustainable agriculture. By embracing and applying the ZBNF principles into agricultural policies and practices, a more resilient, equitable, and environmentally sensitive future can be created for future generations. However, further scientific research and assessment are urgently required before recommending this procedure. Several researchers propose that the effectiveness of ZBNF may be area, agro-climatic zone, or soil-specific. Multi-location trials are necessary to accurately assess the effectiveness of ZBNF in various soil types and agro-climatic zones.

Competing interests: Authors have declared that no competing interests exist.

References

1. Bharucha, Z.P., S. Mitjans and J. Pretty, (2020). Towards redesign at scale through zero budget natural farming in Andhra Pradesh, India. *Int. J. Agric. Sustain*, 18: 1-20
2. Chakraborty D, Nagarajan S, Aggarwal P, Gupta VK, Tomar RK, Garg RN, Sahoo RN, Sarkar A, Chopra U K, Sarma K S and Kalra N, (2008). Effect of mulching on soil and

- plant water status, and the growth and yield of wheat (*Triticum aestivum* L.) in a semi-arid environment. *Agric. Water Manag.* 95(12):1323- 34.
3. FAO , (2016). *Via Campesina L. Zero Budget Natural Farming in India*. Family Farming Knowledge Platform of FAO. Accessed 30 June 2023.
 4. Ghosh, S.; Sarkar, S.; Sau, S.; Karmakar and Brahmachari, K, (2017) .Influence of guava(*Psidium guajava* L.) based intercropping systems on soil health and productivity in alluvial soil of West Bengal, India. *International Journal of Current Microbiology and Applied Sciences.*6(11):241-251.
 5. IIMR. *Natural farming in millets- A Revolution in Indian Agriculture*, (2018). Accessed 26 June 2023 Available:https://www.manage.gov.in/nf/re_sextn/publications/PUB-IIMR-HYD.pdf
 6. Jannoura R, Joergensen GR, Bruns C, (2014). Organic fertilizer effects on growth, crop yield, and soil microbial biomass indices in sole and intercropped peas and oats under organic farming conditions. *Eur. J Agron.* 52(B):259- 270.
 7. Khadse A and Rosset P M, (2019). Zero budget natural farming in India—from inception to institutionalization. *Agroecol. Sustain. Food Syst.*43(7-8):848-871.
 8. Khadse A, Rosset P M, Morales H and Ferguson B G, (2017) Taking agroecology to scale: the Zero Budget Natural Farming peasant movement in Karnataka, India. *J. Peasant Stud.*45(1):192-219.
 9. Khadse, A.; Rosset, P, (2021) Zero Budget Natural Farming in India—From inception to institutionalization. *Desenvolv. Meio Ambient.*58, 579–603.
 10. Khadse, A.; Rosset, P, (2021). Zero Budget Natural Farming in India—From inception to institutionalization. *Desenvolv. Meio Ambient.* 2 58, 579–603.
 11. Kumar R., Kumar S., Yashavanth B. S., Meena P. C., Indoria A. K., Kundu S., & Manjunath M, (2020). Adoption of Natural Farming and its Effect on Crop Yield and Farmers' Livelihood in India. ICAR-National Academy of Agricultural Research Management, Hyderabad, India.
 12. Kumar, R., S. Kumar, B. S. Yashavanth and P. C. Meena, (2019). Natural Farming practices in India: Its adoption and impact on crop yield and farmers' income. *Indian. J. Agric. Econ*, 74(3): 420-432.
 13. Kumar, R.; Kumar, S.; Yashavanth, B.S.; Meena, P.C.; Indoria, A.; Kundu, S.; Manjunath, M, (2020). Adoption of Natural Farming and Its Effect on Crop Yield and Farmers' Livelihood in India; ICAR: New Delhi, India.
 14. Maduka, C.M.; Udensi, C, (2019). Comparative analysis of the effect of some organic manure on soil microorganisms. *Bionatura* 4, 922–925.
 15. McMillen M,(2013). The effect of mulch type and thickness on the soil surface evaporation rate, pp: 69. California Polytechnic State University, San Luis Obispo, California, USA.
 16. Moss D, Bittman M, (2018). Bringing farming back to nature. *New York Times* ,26.
 17. Munster, D, (2018). Performing alternative agriculture: Critique and recuperation in Zero Budget Natural Farming, South India. *J. Polit. Ecol.* 25, 748–764.
 18. Parihar CM, Yadav MR, Jat SL, Singh AK, Kumar B, Pooniya V, Pradhan S, Verma RK, Jat M.L, Jat RK and Saharawat YS, (2018). Long term conservation agriculture and intensified cropping systems: Effect on growth, yield, water and energy-use efficiency of maize in north-western India. *Pedosphere* 28 (6): 952-963.
 19. Patel J. S., Kumar G., Bajpai R., Teli B., Rashid M., Sarma B. K, (2021). “Chapter 18 - PGPR formulations and application in the management of pulse crop health,” in *Biofertilizers*, eds. A. Rakshit, V. S. Meena, M. Parihar, H. B. Singh, and A. K. Singh (Woodhead Publishing), 239–251. 10.1016/B978-0-12-821667-5.00012-9

20. Ray, P.; Lakshmanan, V.; Labbé, J.L.; Craven, K.D, (2020). Microbe to Microbiome: A Paradigm Shift in the Application of Microorganisms for Sustainable Agriculture. *Front Microbiol.* 11, 622926
21. Saharan, B.S.; Tyagi, S.; Kumar, R.; Vijay; Om, H.; Mandal, B.S.; Duhan, J.S, (2023) Application of Jeevamrit Improves Soil Properties in Zero Budget Natural Farming Fields. *Agriculture*.13, 196. [https:// doi.org/10.3390/ agriculture13010196](https://doi.org/10.3390/agriculture13010196)
22. Shaikh NF, Gachande BD, (2015). Effect of organic bio-booster and inorganic inputs on rhizosphere mycoflora population and species diversity of wheat. *Int Jour of Sci and Res*.4:295-302.
23. Shao YH, Xie YX, Wang CY, Yue JQ, Yao YQ, Li XD, Liu WX, Zhu YJ and Guo TC, (2016). Effects of different soil conservation tillage approaches on soil nutrients, water use and wheat-maize yield in rain-fed dry-land regions of North China. *European Journal of Agronomy* 81: 37-45.
24. Shubham, Sharma U and Kaushal R, (2023). Effect of soil applied natural and synthetic nitrification inhibitors on nitrogen transformations and nitrification inhibition in NW Himalayan region of Himachal Pradesh. *Indian Journal of Soil Conservation*51(2): 95-101.
25. Shubham, Sharma U, Kaushal R and Sharma YP, (2022). Effect of Forest Fires on Soil Carbon Dynamics in Different Land Uses under NW Himalayas. *Indian Journal of Ecology* 49(6): 2322-2329. DOI: <https://doi.org/10.55362/IJE/2022/3828>.
26. Sreenivasa MN, Nagaraj MN and Bhat SN,(2010). Beejamruth: A source for beneficial bacteria. *Karnataka Journal of Agricultural Sciences* 17 (3): 72-77.
27. Timsina J, (2018). Can Organic Sources of Nutrients Increase Crop Yields to Meet Global Food Demand? *Agronomy* 8, 214.
28. Tripathi, S., Nagbhushan, S., and Shahidi, T, (2018). Zero Budget Natural Farming for the Sustainable Development Goals: Andhra Pradesh, India. New Delhi: Council on Energy, Environment and Water.
29. United Nations for Food and Agriculture Organization (FAO), (2018).How to Feed the Worldin2050,(2018).Availableonline: https://www.fao.org/fileadmin/templates/wsfs/docs/expert_paper/How_to_Feed_the_World_in_2050.pdf (accessed on 20 December 2021).
30. Yang X, Zhen L, Yang Q, Wang Z, Cui S and Shen Y,(2018). Modelling the effects of conservation tillage on crop water productivity, soil water dynamics and evapotranspiration of a maize-winter wheatsoybean rotation system on the Loess Plateau of China using APSIM. *Agricultural Systems* 166:111-123