

EFFECT OF HUMIC ACID ON GROWTH, YIELD AND SOIL PROPERTIES IN RICE: A REVIEW

ABSTRACT:

Humic acids, natural organic compounds derived from decaying organic matter, have gained significant attention in agriculture due to their potential to promote plant growth, yield, and soil fertility. Humic acid additions boost plant metabolism, promote root development, and increase nutrient absorption, leading to enhanced growth and development. Furthermore, humic acids have a significant role in improving the structure, moisture retention, and availability of nutrients in soil, thereby creating a favorable environment for plant growth. Studies have illustrated that humic acid application can lead to increased yields in a wide range of crops. Humic acid application efficiently increases hormonal activities and various enzyme activities which helps in tricarboxylic acid cycle, glycolysis, and respiration process. The mechanisms underlying this enhancement involve the modulation of physiological processes, such as photosynthesis, nutrient assimilation, and stress tolerance. In addition to promoting plant growth and yield, humic acid amendments have been shown to positively influence on soil properties. These include increased microbial activity, enhanced soil aggregation, and improved nutrient cycling. Moreover, humic acids can mitigate soil degradation processes, such as salinization, thereby promoting sustainable agricultural practices. Through the utilization of humic acids, farmers may maximize crop yields while reducing their ecological footprint, therefore promoting food security and sustainable agriculture.

Key words: Sustainable agriculture; nutrient cycling; stress tolerance; tricarboxylic acid cycle; glycolysis; respiration; salinization.

1. INTRODUCTION:

Rice (*Oryza sativa* L.) represents one of the most important cereals in the world. China is the world's leading producer of rice, with India coming in second and being the world's largest exporter. It occupies one-third of the total cultivable land and grown as a major staple food crop in India [1]. It is one of the richest sources of carbohydrates, vitamins, proteins, and minerals; the endosperm of grains counts for about 90% of carbohydrates of its total produced dry weight [2, 3]. Milling products of rice yield contains 70 percent of

33 the major product as rice (endosperm) and 30 percent of the by-product as 20 percent husk, 8 percent
34 bran, and 2 percent germ [4]. In recent times, it has been investigated that rice bran is one of the important
35 components which has various potential biological properties (i.e. anti-inflammatory action and antioxidant
36 functions) which reduce the occurrence of cancer and cholesterol levels and impede coronary artery
37 diseases and rice bran is also rich in phytic acid (59.4 - 60.9 gm per kg) [5,6,7]. The application of synthetic
38 fertilizers alone under continuous practices of intensive cropping does not positively sustain the crop
39 productivity status, but the addition of organic substances with applied chemical fertilizers effectively
40 improves the physical properties of soil and maintains greater soil fertility status with better yield production
41 [8]. Among the organic substances, humic acid is one of the greater organic molecules that have various
42 important roles on many agronomic parameters and soil properties [9]. Humic acid is mainly produced from
43 the degraded waste materials of plants and animals and plays some vital roles in agriculture systems such
44 as increment in soil biochemical and physical properties through greater texture, structure, microbial
45 activities, and water holding capacity [10,11]; enhance micronutrient nutrient availability in soil as a
46 chelating agent and increase nutrients uptake to the plants [12]; decreases the intake of heavy metal which
47 are toxic to the plants [13]. The application of humic acid also improved crop growth status by enhancing
48 growth promoting hormones like cytokinin and auxin, which help in the metabolic process of nutrients, stress
49 resistance, and photosynthesis activity [14,15]. Additionally, humic acid has positive effects on plant cell
50 membranes which helps in greater transportation of minerals, higher protein synthesis, greater enzyme
51 activities, lesser activity of toxic elements, and higher microbial population [16]. This review explores the
52 effects of humic acid application on many aspects of agricultural productivity, soil properties, and plant
53 physiology.

54

55 2. RELATIONSHIP BETWEEN STRUCTURES AND FUNCTIONALITY OF HUMIC ACIDS:

56 The functionality of humic acid is directly associated with the structural composition, which is dependent on
57 the type of sources [17]. Humic acid structure contains several functional groups, the most prevalent among
58 them are the phenolic (OH) and carboxylic (COOH) groups [18]. These two functional groups act as major
59 primary groups that help in the improvement of soil chemical and physical properties and promote greater
60 growth of the plants (**Fig 1**) [18,19]. The disintegration of these two functional groups forms nonpolar and
61 polar ends (hydrophobic and hydrophilic parts, respectively), which are involved in the beneficial processes
62 of humic acid functions [20]. The hydrophobic end is associated with repelling activities, whereas the
63 hydrophilic end is mostly engaged in chelating activities [21]. Once the disintegration of COOH and OH
64 occurs, the cationic metals are retained in the soil by aggregation with the polar end (anionic parts) through
65 electrostatic bonding [22]. The water-loving hydrophilic part forms micelle, which helps in the increment of
66 water holding capacity of the soil. Similarly, the hydrophobic parts reduce the rate of water infiltration and
67 improve the aggregation stability of clay particles [21]. Recently, a research study conducted by de Castro
68 *et al.*, (2021) [17], reported that the aliphatic and aromatic groups of humic acid effectively contributed to
69 the increment of soluble sugars and nitrogen uptake, leading to enhancement in rice yield production.

70 García *et al.*, (2016) [23] also reported that these functional groups of humic acid effectively stimulate the
71 growth of the root system of rice seedlings.

72

73

74

75

76

77

78

79

80

81


82

83

84

85

86



83

84

85

Fig 1: Functions of humic acid on plant and soil attributing characters

87

3. MECHANISM OF HUMIC ACID ON PLANT GROWTH:

88

89

90

91

92

93

94

95

96


97

98

99

Humic acid has an important role in plant growth by stimulating carbon and nitrogen metabolism. Humic acid triggered enzyme activities such as glutamine synthetase, nitrate reductase, and glutamate dehydrogenase, which are linked to the nitrogen assimilation processes [24]. Furthermore, humic substances stimulate nitric oxide (NO) at the emergence point of lateral roots. Nitric oxide is a bioactive substance that has a variety of roles in the physiological activities of plants, including the formation of their roots [25]. The incorporation of humic substances primarily increases NO accumulation, which helps in the morphological changes in the root system such as the increment of secondary roots, increment in thickness of the roots, and higher fresh weight of roots [26]. Canellas *et al.*, (2014) [27] reported that humic acid increases the density and length of the root hairs as well as cell proliferation of root tissues. The humic substances also improve hormonal effects on plant's root and shoot system [28]. Humic substances can stimulate ion transport and H⁺-ATPase activities in the plasma membranes of the root system [29]. Both these impacts are directly involved in nutrient acquisition and uptake of nutrients in the plant system. Nardi

100 *et al.*, [30] reported that humic substances increase enzyme activities those are involved in tricarboxylic
 101 acid cycle and glycolysis process. The enzyme activities related to the glycolysis process are pyruvate
 102 kinase, phosphofructokinase, glucokinase, phosphoglucose isomerase, and the enzyme activities related
 103 to the respiration process are malate dehydrogenase, NADP+-isocitrate dehydrogenase, and citrate
 104 synthase. Humic substance application in the root zone can trigger physiological responses for shoot
 105 development by the activation of auxin, like phospholipase A2 as a stomatal opening regulator [31]. **Fig 2**
 106 represents the activities of humic on plant growth by its various mechanisms.



117 Fig 2: Mechanisms of humic acid on plant growth

118 4. IMPACTS OF HUMIC ACID ON THE GROWTH PARAMETERS:

119 Humic acid is one of the most effective biostimulants among the humic substances, which have many
 120 positive influences on rice growth and development. It efficiently promotes the photosynthesis rate, higher
 121 plant growth, greater amount of dry matter accumulation, and production and accumulation of more amount
 122 of organic matter [32, 33]. The addition of humic acid with various synthetic fertilizers increases plant growth
 123 with greater amount of nutrient uptake. Humic acid can improve soil's biochemical properties by enhancing
 124 enzyme activities, microbial population and their activities, water holding capacity (WHC) and cation
 125 exchange capacity (CEC) in the soil that ultimately improve plant's nutrient uptake and efficient plant growth
 126 [34]. This impact of humic acid ultimately reflects on plant height, leaf area index (LAI), and biomass
 127 production through greater photosynthesis [35]. Various scientific studies mentioned that humic acid also
 128 has an effect on chlorophyll content and root-shoot development of plants. It improves the growth of shoots
 129 and roots by increasing growth promoting hormones i.e. production of cytokinin and auxin, and enzyme
 130 activities in the plants [36, 37]. The increased uptake of plant nutrients (micro and macronutrients) through
 131 humic acid enhances the chlorophyll concentration in the leaf, which positively improves the shoot growth

132 of the plants [38]. With the slow release of soil nutrients due to the humic acid, plants can uptake more
 133 amount of nutrients, which leads to the higher number of tillers in the plants [39]. **Table 1** shows that the
 134 addition of humic acid both in soil or foliar spray significantly increases growth attributing characteristics
 135 such as plant height, number of tillers, dry weight, etc. **However, Saha *et al.*, (2013) [40] reported that higher**
 136 **dose of humic acid@ 6 L ha⁻¹ decreases plant growth attributes compared to humic acid@ 3 L ha⁻¹. It may**
 137 **be due to an imbalance in nutrients and toxic effects on plant metabolism, which damages plant**
 138 **physiological processes and nutrient uptake to the plants.**

139 Table 1: Effect of humic acid on various growth attributes of rice

SI. No.	Treatments	Plant height (cm)	Number of tillers	Dry weight (g)	Reference
1	90 kg ha ⁻¹ nitrogen fertilizer + no humic acid	71.77	24.50	-	[39]
	90 kg ha ⁻¹ nitrogen fertilizer + 5 kg ha ⁻¹ humic acid	76.77	33.37	-	
2	No humic acid	86.11	10.33	-	[40]
	3 L ha ⁻¹ humic acid	86.33	11.56	-	
	6 L ha⁻¹ humic acid	86.22	11.00		
3	RDF + 2.5 kg ha ⁻¹ humic acid-based biostimulant	75.98	14.50	33.23	[41]
	RDF + 5 kg ha ⁻¹ humic acid based biostimulant	77.39	14.81	34.49	
4	Without foliar spray	85.95	501.5	-	[42]
	Foliar spray of humic acid	91.45	552.5	-	
5	No humic acid	69.24	11.05	32.09	[43]
	6 kg ha ⁻¹ of humic acid	70.12	12.94	39.07	
6	Control	94.8 ± 2.19	-	3.44 ± 0.317	[44]
	Humic acid	103 ± 2.53	-	4.62 ± 0.373	
7	Control	114.33 ± 0.60	-	6.424 ± 0.299	[45]
	Humic acid	116.36 ± 0.30	-	7.784 ± 0.622	

140

141 5. IMPACTS OF HUMIC ACID ON THE YIELD AND QUALITY ATTRIBUTES:

142 The yield attributing characteristics of the plants are effectively influenced by humic acid application both
 143 as soil application and foliar spray. Humic acid enhances physical and chemical properties and microbial
 144 activities in the soil as well as increases hormonal activities, which leads to greater uptake of plant nutrients
 145 resulting in more production of photosynthates and greater source to sink translocation of assimilates, which
 146 finally produces higher number of effective tillers, panicle length, number of grains panicle⁻¹, grain and straw
 147 yield, and the test weight of grains [46]. Sivakumar *et al.*, (2007) [47] reported that humic acid application
 148 significantly increases N, P, and K uptake in plants, which helps in the greater accumulation of dry matter

149 production and higher grain yield production. Additionally, it also functions as a naturally occurring
 150 biostimulant, improving plant resistance to abiotic stresses including drought and salinity, and minimizing
 151 production losses. On the other hand, humic acid increases protein content in the grains by improving
 152 nitrogen uptake in the sink [48, 49]. The increment in carbohydrate content also happens by humic acid,
 153 which helps in the improved rate of photosynthesis, finally leading to greater translocation of assimilates
 154 from the source to sink [50]. The application of humic acid produces better root system and increases cell
 155 membrane permeability, leading to greater uptake of plant nutrients, and finally, concentrations of nitrogen,
 156 potassium and phosphorus increase in the plants [51, 52]. **Table 2** shows the effects of humic acid on grain
 157 and straw yield of rice, which indicates the impact of humic acid on yield attributing characteristics such as
 158 number of effective tillers, panicle length, number of grains panicle⁻¹, test weight, etc. **However, according**
 159 **to Sivakumar *et al.*, (2007) [47], soil application of humic acid up to 20 kg ha⁻¹ significantly increases crop**
 160 **yield, but beyond this, it decreases yield production due to its toxic effects on plant metabolic processes.**

161 Table 2: Effect of humic acid on yield parameters of rice


SI. No.	Treatments	Grain yield (t ha ⁻¹)	Straw yield (t ha ⁻¹)	Reference
1	No humic acid	3.08	4.62	[47]
	20 kg ha ⁻¹ humic acid	3.63	5.45	
	30 kg ha ⁻¹ humic acid	3.48	5.22	
2	No humic acid	3.09	8.22	[40]
	3 L ha ⁻¹ humic acid	3.10	8.33	
	6 L ha ⁻¹ humic acid	3.20	8.45	
3	100% RDF	4.74	-	[53]
	100% RDF + Humic acid	5.12	-	
4	Humic acid based biostimulant @ 2.5 kg ha ⁻¹	4.10	5.24	[41]
	Humic acid based biostimulant @ 5 kg ha ⁻¹	4.17	5.39	
5	Without foliar spray	3.05	2.7	[42]
	Foliar spray of humic acid	3.82	3.4	
6	15 kg nitrogen fed ⁻¹	2.98	2.89	[54]
	15 kg nitrogen fed ⁻¹ + humic acid	3.94	3.07	

162

163 6. IMPACTS OF HUMIC ACID ON THE SOIL PROPERTIES:

164 Humic acid acts as a chelating agent by attracting positive ions and making the greater availability of
 165 nutrients to the plants by their slow release ability [52]. The chelating properties help in the prevention of
 166 fixation, oxidation, leaching, and precipitation of micronutrients in the soil [55]. The soil pH is also influenced
 167 by humic acid but it depends on the number of phenolic and carboxylic functional groups present in their
 168 structure [56]. Further, humic acid can stabilize ammonium in the soil, which leads to better nitrogen
 169 availability [57]. The nitrogen molecules are also present in humic substances which become accessible to

170 the plants after their application in soil [21]. Apart from nitrogen, phosphorus content in the soil also
 171 improves by humic acid application due to their synergistic effects on phosphatase activity, leading to
 172 greater solubilization of phosphorus in soil [58]. Zhu *et al.*, (2018) [59] mentioned that humic acid increases
 173 the desorption of phosphate ions and reduces sorption, which makes the availability of phosphorus in the
 174 soil solution. Humic acid also improves the plasma membrane permeability and increases nutrient
 175 absorption in plants [60]. While the higher rate of humic acid application improves soil's physical properties,
 176 its binding capability reduces micronutrient availability for plant uptake. The phenolic and carboxylic
 177 functional groups help in the chelating process, which makes metal complexes of humic acid in soil, leading
 178 to greater availability of micronutrients in the soil but their availability for plant uptake decreases [61].
 179 However, the application of humic acid can effectively increase micronutrient availability in the soil but
 180 cannot be assumed to have a synergistic linear relationship with plant uptake due to the different
 181 compositions of physical and chemical properties of humic acid. **Fig 3** shows the impacts of humic acid on
 182 soil properties.



183
 184
 185
 186
 187
 188
 189
 190 Fig 3: Effect of humic acid on soil properties and availability of nutrients for plant uptake

191
 192 **7. CONCLUSION:**

193 This review paper has revealed that humic acid has greater potential impacts on various agronomic
 194 parameters and soil properties. It increases the intake of nutrients, strengthens the structure of the soil,
 195 reduces stress, encourages biological activity, and may even benefit farmers commercially. Although there
 196 may be an initial expense involved, it is overcome in the long term by its benefits to soil health and
 197 productivity. However, the current study suggests that introducing humic acid in cropping practices may
 198 help to create productive and sustainable agricultural systems.

199 **ACKNOWLEDGMENTS:**

200 The authors would like to acknowledge the School of Agriculture, Lovely Professional University for their
 201 consistent support to write this review paper.

202 **COMPETING INTERESTS:**

203 The authors have declared that there are no competing interests.

204 **AUTHOR'S CONTRIBUTIONS:**

205 Bidyabhusan Bera and Kangujam Bokado have selected the topic for review and prepared the framework
206 of the review paper. Bidyabhusan Bera and Shainika Arambam compiled all the research findings and
207 completed the review writing. The final editing and proof reading was carried out by by Kangujam Bokado
208 and Barkha.

REFERENCE:

1. Yengkokpam P, Kumari P, Singh VK, Ningthi KC, Dhiman S, Chaudhary M, Sharma R. Application and Management of Nitrogenous Fertilizer in Rice Field: A Review. *Asian Journal of Soil Science and Plant Nutrition*. 2024;10(2):64-71.
2. Rathna Priya TS, Eliazar Nelson AR, Ravichandran K, Antony U. Nutritional and functional properties of coloured rice varieties of South India: a review. *Journal of Ethnic Foods*. 2019;6(1):1-1. Doi: <https://doi.org/10.1186/s42779-019-0017-3>.
3. Das P, Adak S, Lahiri Majumder A. Genetic manipulation for improved nutritional quality in rice. *Frontiers in Genetics*. 2020;11:531308. Doi:<https://doi.org/10.3389/fgene.2020.00776>.
4. Van Hoed V, Depaemelaere G, Ayala JV, Santiwattana P, Verhé R, De Greyt W. Influence of chemical refining on the major and minor components of rice bran oil. *Journal of the American Oil Chemists' Society*. 2006;83:315-321.
5. Liu SX, Mamidipally PK. Quality comparison of rice bran oil extracted with d-limonene and hexane. *Cereal Chemistry*. 2005;82(2):209-215.
6. Canan C, Cruz FT, Delarozza F, Casagrande R, Sarmiento CP, Shimokomaki M, Ida EI. Studies on the extraction and purification of phytic acid from rice bran. *Journal of food composition and analysis*. 2011;24(7):1057-1063.
7. Kumar A, Sahu C, Panda PA, Biswal M, Sah RP, Lal MK, Baig MJ, Swain P, Behera L, Chattopadhyay K, Sharma S. Phytic acid content may affect starch digestibility and glycemic index value of rice (*Oryza sativa* L.). *Journal of the Science of Food and Agriculture*. 2020;100(4):1598-1607.
8. Elankavi S, Nambi J, Ramesh S, Jawahar S, Lavanya K. Influence of different doses of fertilizers and foliar spray of nutrients on yield and yield attributes of rice. *Annals of the Romanian Society for Cell Biology*. 2020:1127-1134.
9. Ampong K, Thilakaranthna MS, Gorim LY. Understanding the role of humic acids on crop performance and soil health. *Frontiers in Agronomy*. 2022;4:848621.
10. Nardi S, Schiavon M, Francioso O. Chemical structure and biological activity of humic substances define their role as plant growth promoters. *Molecules*. 2021;26(8):2256. Doi: [10.3390/molecules26082256](https://doi.org/10.3390/molecules26082256).
11. Shah ZH, Rehman HM, Akhtar T, Alsamadany H, Hamooh BT, Mujtaba T, Daur I, Al Zahrani Y, Alzahrani HA, Ali S, Yang SH. Humic substances: Determining potential molecular regulatory processes in plants. *Frontiers in plant science*. 2018;9:263. Doi: [10.3389/fpls.2018.00263](https://doi.org/10.3389/fpls.2018.00263).
12. Yang F, Tang C, Antonietti M. Natural and artificial humic substances to manage minerals, ions, water, and soil microorganisms. *Chemical Society Reviews*. 2021;50(10):6221-6239. Doi: [10.1039/D0CS01363C](https://doi.org/10.1039/D0CS01363C).

13. Wu S, Li R, Peng S, Liu Q, Zhu X. Effect of humic acid on transformation of soil heavy metals. In IOP Conference Series: Materials Science and Engineering. 2017;207(1): 012089. Doi: [10.1088/1757-899X/207/1/012089](https://doi.org/10.1088/1757-899X/207/1/012089).
14. Canellas LP, Canellas NO, da S. Irineu LE, Olivares FL, Piccolo A. Plant chemical priming by humic acids. Chemical and Biological Technologies in Agriculture. 2020;7:1-7. Doi: [10.1186/s40538-020-00178-4](https://doi.org/10.1186/s40538-020-00178-4).
15. Jindo K, Canellas LP, Albacete A, Figueiredo dos Santos L, Frinhani Rocha RL, Carvalho Baia D, Oliveira Aguiar Canellas N, Goron TL, Olivares FL. Interaction between humic substances and plant hormones for phosphorous acquisition. Agronomy. 2020 May 1;10(5):640.
16. Khaled H, Fawy HA. Effect of different levels of Humic acids on the nutrient content, plant growth, and soil properties under conditions of salinity. Soil and Water Research. 2011;6(1):21-29.
17. de Castro TA, Berbara RL, Tavares OC, da Graca Mello DF, Pereira EG, de Souza CD, Espinosa LM, García AC. Humic acids induce a eustress state via photosynthesis and nitrogen metabolism leading to a root growth improvement in rice plants. Plant Physiology and Biochemistry. 2021;162:171-184. Doi: [10.1016/j.plaphy.2021.02.043](https://doi.org/10.1016/j.plaphy.2021.02.043).
18. Nardi S, Schiavon M, Francioso O. Chemical structure and biological activity of humic substances define their role as plant growth promoters. Molecules. 2021;26(8):2256. Doi: [10.3390/molecules26082256](https://doi.org/10.3390/molecules26082256).
19. de Melo BA, Motta FL, Santana MH. Humic acids: Structural properties and multiple functionalities for novel technological developments. Materials Science and Engineering: C. 2016;62:967-974. Doi: [10.1016/j.msec.2015.12.001](https://doi.org/10.1016/j.msec.2015.12.001).
20. Mirza MA, Agarwal SP, Rahman MA, Rauf A, Ahmad N, Alam A, Iqbal Z. Role of humic acid on oral drug delivery of an antiepileptic drug. Drug development and industrial pharmacy. 2011;37(3):310-9. Doi: [10.3109/03639045.2010.512011](https://doi.org/10.3109/03639045.2010.512011).
21. Billingham KL. "Humic products-potential or presumption for agriculture? Can humic products improve my soil?" in 27th Annual Conference (Orange, NSW: Grassland Society of NSW Inc.). 2012.
22. Among K, Thilakaranthna MS, Gorim LY. Understanding the role of humic acids on crop performance and soil health. Frontiers in Agronomy. 2022;4:848621.
23. García AC, de Souza LG, Pereira MG, Castro RN, García-Mina JM, Zonta E, Lisboa FJ, Berbara RL. Structure-property-function relationship in humic substances to explain the biological activity in plants. Scientific reports. 2016;6(1):20798. Doi: [10.1038/srep20798](https://doi.org/10.1038/srep20798).
24. Canellas LP, Balmori DM, Médici LO, Aguiar NO, Campostrini E, Rosa RC, Façanha AR, Olivares FL. A combination of humic substances and *Herbaspirillum seropedicae* inoculation enhances the growth of maize (*Zea mays* L.) Plant Soil. 2013;366:119–132. Doi: [10.1007/s11104-012-1382-5](https://doi.org/10.1007/s11104-012-1382-5).
25. Lamattina L, García-Mata C, Graziano M, Pagnussat G. Nitric oxide: the versatility of an extensive signal molecule. Annual review of plant biology. 2003;54(1):109-36.

26. Mora V, Baigorri R, Bacaicoa E, Zamarreno AM, García-Mina JM. The humic acid-induced changes in the root concentration of nitric oxide, IAA and ethylene do not explain the changes in root architecture caused by humic acid in cucumber. *Environmental and Experimental Botany*. 2012;76:24-32.
27. Canellas LP, Olivares FL. Physiological responses to humic substances as plant growth promoter. *Chemical and Biological Technologies in Agriculture*. 2014;1:1-1.
28. Muscolo A, Sidari M, Nardi S. Humic substance: relationship between structure and activity. Deeper information suggests univocal findings. *Journal of Geochemical Exploration*. 2013;129:57-63.
29. Pinton R, Cesco S, Varanini Z. Role of humic substances in the rhizosphere. *Biophysico-chemical processes involving natural nonliving organic matter in environmental systems*. 2009;2:341-66.
30. Nardi S, Muscolo A, Vaccaro S, Baiano S, Spaccini R, Piccolo A. Relationship between molecular characteristics of soil humic fractions and glycolytic pathway and krebs cycle in maize seedlings. *Soil Biology and Biochemistry*. 2007;39(12):3138-46.
31. Russell L, Stokes AR, Macdonald H, Muscolo A, Nardi S. Stomatal responses to humic substances and auxin are sensitive to inhibitors of phospholipase A2. *Plant and Soil*. 2006;283:175-85.
32. Dan C, Liang-Gang Z, Jun X, Qian Z, Yan Z. Effects of biofertilizer on organically cultured cucumber growth and soil biological characteristics. *Yingyong Shengtai Xuebao*. 2010;21(10).
33. Zhang XC, Shangguan ZP. Effect of nitrogen fertilization on photosynthetic pigment and fluorescence characteristics in leaves of winter wheat cultivars on dryland. *J. Nuclear Agric. Sci*. 2007;21:299-304.
34. Khattak RA, Dost M. Seed cotton yield and nutrient concentrations as influenced by lignitic coal derived humic acid in salt-affected soils. *Sarhad Journal of Agriculture*. 2010;26(1):43-49.
35. Tahir MM, Khurshid M, Khan MZ, Abbasi MK, Kazmi MH. Lignite-derived humic acid effect on growth of wheat plants in different soils. *Pedosphere*. 2011;21(1):124-31.
36. Rose MT, Patti AF, Little KR, Brown AL, Jackson WR, Cavagnaro TR. A meta-analysis and review of plant-growth response to humic substances: practical implications for agriculture. *Advances in agronomy*. 2014;124:37-89. Doi: [10.1016/B978-0-12-800138-7.00002-4](https://doi.org/10.1016/B978-0-12-800138-7.00002-4)
37. Olaetxea M, Mora V, Baigorri R, Zamarreño AM, García-Mina JM. The singular molecular conformation of humic acids in solution influences their ability to enhance root hydraulic conductivity and plant growth. *Molecules*. 2020;26(1):3. Doi: [10.3390/molecules26010003](https://doi.org/10.3390/molecules26010003)
38. Sible CN, Seebauer JR, Below FE. Plant biostimulants: A categorical review, their implications for row crop production, and relation to soil health indicators. *Agronomy*. 2021;11(7):1297. Doi: [10.3390/agronomy11071297](https://doi.org/10.3390/agronomy11071297)
39. Sunarya Y, Tedjaningsih T. Increasing nitrogen fertilizer efficiency on wetland rice by using humic acid. *Journal of Tropical Soils*. 2016;20(3):143-148. Doi: [10.5400/jts.2015.20.3.143](https://doi.org/10.5400/jts.2015.20.3.143)

40. Saha R, Saieed MA, Chowdhury MA. Growth and yield of rice (*Oryza sativa*) as influenced by humic acid and poultry manure. *Universal Journal of Plant Science*. 2013;1(3):78-84.
Doi: [10.13189/ujps.2013.010304](https://doi.org/10.13189/ujps.2013.010304)
41. Karennavar SS, Chavan VG, More VG, Dhopavkar RV, Desai SS, Bodake PS. Effect of humic acid based bio-stimulant on growth, yield and yield attributing characters of kharif rice (*Oryza sativa* L.). *Pharma Innovation*. 2022;11(11):319-322.
42. Osman EA, El-Masry AA, Khatab KA. Effect of nitrogen fertilizer sources and foliar spray of humic and/or fulvic acids on yield and quality of rice plants. *Advances in Applied Science Research*. 2013;4(4):174-83.
43. Nugroho WA, Suparman S, Wakhid N, Bhermana A. Rice growth and yield on acid sulfate soils following humic acid and NPK fertilizer application. In *AIP Conference Proceedings*. 2024;2957(1).
Doi: <https://doi.org/10.1063/5.0184029>
44. Abu-Ria M, Shukry W, Abo-Hamed S, Albaqami M, Almuqadam L, Ibraheem F. Humic acid modulates ionic homeostasis, osmolytes content, and antioxidant defense to improve salt tolerance in rice. *Plants*. 2023;12(9):1834. Doi: <https://doi.org/10.3390/plants12091834>
45. Shukry WM, Abu-Ria ME, Abo-Hamed SA, Anis GB, Ibraheem F. The efficiency of humic acid for improving salinity tolerance in salt sensitive rice (*Oryza sativa*): growth responses and physiological mechanisms. *Gesunde Pflanzen*. 2023;75(6):2639-53.
46. Elankavi S, Nambi J, Ramesh S, Jawahar S, Lavanya K. Influence of different doses of fertilizers and foliar spray of nutrients on yield and yield attributes of rice. *Annals of the Romanian Society for Cell Biology*. 2020:1127-1134.
47. Sivakumar K, Devarajan L, Dhanasekaran K, Venkatakrisnan D, Surendran U. Effect of humic acid on the yield and nutrient uptake of rice. *ORYZA-An International Journal on Rice*. 2007;44(3):277-279.
48. Abd El-Gawad MA. Effect of humic acid application as well as bio and mineral phosphorus fertilization on growth, green pods, dry seed yield and quality of peas (*Pisum sativum* L.) under the newly reclaimed soil conditions. *Egypt. J. Appl. Sci*. 2013;28(7):338-360.
49. El-Galad MA, Sayed DA, El-Shal RM. Effect of humic acid and compost applied alone or in combination with sulphur on soil fertility and faba bean productivity under saline soil conditions. *Journal of Soil Sciences and Agricultural Engineering*. 2013;4(10):1139-1157.
50. Azcona I, Pascual I, Aguirreolea J, Fuentes M, García-Mina JM, Sánchez-Díaz M. Growth and development of pepper are affected by humic substances derived from composted sludge. *Journal of Plant Nutrition and Soil Science*. 2011;174(6):916-924.
51. Mahmoud MM, Hassanein AH, Mansour SF, Khalefa AM. Effect of soil and foliar application of humic acid on growth and productivity of soybean plants grown on a calcareous soil under different levels of mineral fertilizers. *Journal of Soil Sciences and Agricultural Engineering*. 2011;2(8):881-890.

52. Eshwar M, Srilatha M, Rekha KB, Sharma SH. Effect of humic substances (humic, fulvic acid) and chemical fertilizers on nutrient uptake, dry matter production of aerobic rice (*Oryza sativa* L.). *Journal of Pharmacognosy and Phytochemistry*. 2017;6(5):1063-1066.
53. Vanitha K, Mohandass S. Effect of humic acid on plant growth characters and grain yield of drip fertigated aerobic rice (*Oryza sativa* L.). *The bioscan*. 2014;9(1):45-50.
54. El-Gohary AA, Osman EA, Khatib KA. Effect of nitrogen fertilization, humic acid and compost extract on yield and quality of rice plants. *Journal of Soil Sciences and Agricultural Engineering*. 2010;1(1):77-91.
55. Kadam SR, Amrutsagar VM, Deshpande AN. Influence of organic nitrogen sources with fulvic acid spray on yield and nutrient uptake of soyabean on inceptisol. *Journal of Soils and Crops*. 2010; 20(1):58-63.
56. Rupiasih NN, Vidyasagar P. A review: Compositions, structures, properties and applications of humic substances. *J. Adv. Sci. Technol*. 2005;8:16-25.
57. Zhang SQ, Liang YU, Wei LI, Lin ZA, LI YT, HU SW, ZHAO BQ. Effects of urea enhanced with different weathered coal-derived humic acid components on maize yield and fate of fertilizer nitrogen. *Journal of integrative agriculture*. 2019;18(3):656-666.
Doi: [10.1016/S2095-3119\(18\)61950-1](https://doi.org/10.1016/S2095-3119(18)61950-1)
58. Sharma SB, Sayyed RZ, Trivedi MH, Gobi TA. Phosphate solubilizing microbes: sustainable approach for managing phosphorus deficiency in agricultural soils. *SpringerPlus*. 2013;2:1-4.
59. Zhu J, Li M, Whelan M. Phosphorus activators contribute to legacy phosphorus availability in agricultural soils: A review. *Science of the Total Environment*. 2018;612:522-537.
Doi: [10.1016/j.scitotenv.2017.08.095](https://doi.org/10.1016/j.scitotenv.2017.08.095)
60. Nardi S, Pizzeghello D, Muscolo A, Vianello A. Physiological effects of humic substances on higher plants. *Soil Biology and Biochemistry*. 2002;34(11):1527-1536.
Doi: [10.1016/S0038-0717\(02\)00174-8](https://doi.org/10.1016/S0038-0717(02)00174-8)
61. Yang F, Tang C, Antonietti M. Natural and artificial humic substances to manage minerals, ions, water, and soil microorganisms. *Chemical Society Reviews*. 2021;50(10):6221-6239.
Doi: [10.1039/D0CS01363C](https://doi.org/10.1039/D0CS01363C)