

# Current knowledge of enhancing power of Salicylic acid for biotic and abiotic stress tolerance in field crops

**Comment [i1]:** Current knowledge of salicylic acid ability to increase field crops resistance to biotic and abiotic stress

### ABSTRACT

Salicylic acid is involved in the control of plant development in addition to being a well-known signal molecule that mediates plant immunity. On the flip side of side, its effect on plant development has not yet been thoroughly defined, despite its well-established involvement in plant immunity. The corpus of evidence demonstrating salicylic acid's vital roles in regulating cell division and expansion—two processes that eventually establish the structure of a plant. The present understanding of the action and molecular processes by which salicylic acid controls plant development via a variety of channels is summed up in this review. Here, it is emphasized how salicylic acid influences cell division and expansion to regulate growth control. The relationships between salicylic acid and other hormones as well as their significance in determining plant development were also covered. Future crop improvement will greatly benefit from a deeper understanding of the process underpinning salicylic acid-mediated growth.

*Keywords: Salicylic acid; plant immunity; cell division; hormones.*

**Comment [i2]:** Salicylic acid is a well-known signal molecule that mediates plant resistance and is also involved in control for plant development. Conversely, despite its well-established role in plant resistance, its impact on plant development is still poorly understood. The body of research indicating the essential functions of salicylic acid in controlling cell division and expansion, two processes that ultimately determine a plant's structure. This study summarizes the current knowledge of the mechanisms and molecular mechanisms via which salicylic acid regulates plant development through a range of pathways. Here, the role of salicylic acid in controlling growth regulation through effects on cell division and expansion is highlighted.

### 1. INTRODUCTION

Ortho-hydroxybenzoic acid, or salicylic acid (SA)—derived from the Latin word *Salix*, which means willow tree—is another name for this phenolic derivative that is widely found in the plant kingdom and is recognized for its ability to regulate several physiological and biochemical processes, including plant signaling or defense mechanism, thermogenesis, and response to different abiotic and biotic stress [1,2].

Salicylic acid may be extracted from plants in both free and conjugated form, and it is a member of a broad class of plant phenolics from a chemical perspective. The conjugated form specifically starts with the aromatic ring being hydroxylated, methylated, and/or glucosylated [3,4]. Johan Büchner first extracted salicin, one of the naturally occurring salicylic acid derivatives, from the willow tree's

**Comment [i3]:** The methods and molecular processes by which salicylic acid controls plant development through a variety of pathways are compiled in this study. Here, it is illustrated how salicylic acid regulates growth by having an impact on cell division and expansion.

**Comment [i4]:** Delete (.)

(*Salix* sp.) bark in 1828 [5,6]. The concentration of this natural compound in plants varies significantly with the seasons with values of 3 mg/g of fresh biomass in *S. lapponum* plants [7]. The highest content of salicylic acid is found in spring and summer and the lowest content in autumn and winter. Subsequently, it was found that nearly all willow trees, including *Salix daphnoides*, *Salix purpurea*, *Salix alba*, and *Salix fragilis* were particularly rich in it [7]. The Italian chemist Raffaele Piria obtained salicylic acid in the bloom and buds of the European plant *Spiraea ulmaria*, later renamed *Filipendula ulmaria* (L.) Maxim. Piria was the first scientist to find this natural substance in species other than *Salix* sp. in late 1838. The identification of this molecule as non-specific to the *Salix* genus has allowed for further research into its production, biochemical properties, and physiological roles in plants [8].

Comment [i5]: by

Comment [i6]: Phytohormones

Two metabolic pathways are known to create salicylic acid via the shikimate pathway in terms of its production. The first route—also referred to as the phenylalanine route—occurs in the cytoplasm of the cell. Trans-cinnamic acid (t-CA), which is oxidised to benzoic acid (BA) is produced by the enzyme phenylalanine ammonia lyase (PAL) from phenylalanine (Phe). Salicylic acid is subsequently formed via the hydroxylation of the aromatic ring of benzoic acid (BA), which is catalyzed by the enzyme benzoic-acid-2-hydroxylase (BA2H). Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) must be present for BA2H to convert benzoic acid (BA) into salicylic acid [9- 11]. The initial evidence for the first pathway came from Ellis and Amrhein, who noted that salicylic acid was produced when *Gaultheria procumbens* plants were fed <sup>14</sup>C-cinnamic acid or <sup>14</sup>C-benzoic acid [12]. Nevertheless, new findings suggest that salicylic acid is more likely to originate directly from benzoyl glucose, a conjugated form of benzoic acid (BA) [11,13].

Comment [i7]: In terms of production, two metabolic pathways are known to produce salicylic acid via the shikimate pathway

Comment [i8]: plants are fed with

Comment [i9]: most likely derived

The second step, known as the isochorismate (IC) pathway, takes place within the chloroplast [14-16]. Isochorismate pyruvate lyase (IPL) and Isochorismate synthase (ICS) are the two enzymes that catalyze the conversion of chorismate in plants into isochorismate and ultimately salicylic acid. It is well recognized from a physiological perspective that salicylic acid is essential for controlling plant development and growth, defense against different abiotic and biotic stress, and immunological responses [4,17- 21]. From that point on, there was an exponential rise in the number of articles focusing on salicylic acid as a plant growth regulator, signaling molecule, and plant elicitor that protects plants from different abiotic and biotic stress [20- 27].

Comment [i10]: growth regulation

Comment [i11]: stresses

By concentrating on these elements, the current study offers a thorough compilation of information on the roles that salicylic acid plays in plant growth and development (fig 1). The objective is to create a clear picture of salicylic acid and help guide future research on this subject.

Comment [i12]: The current study provides an extensive compilation of data on the roles that salicylic acid plays in plant growth and development by focusing on these factors (fig 1). The goal is to provide a clear image of salicylic acid and aid in directing further studies

## 2. BOOSTING POWER OF SA ON GROWTH AND DEVELOPMENT OF PLANT

Salicylic acid might play disputed functions in the growth and development of plants, depending on its concentration, the growing environment of the plant, and its developmental stage [28]. High concentrations of salicylic acid often have a detrimental effect on the growth and development of plants (depending on the type of plant; nonetheless, >1 mM salicylic acid is considered a high concentration). However, using the right amounts of salicylic acid had positive benefits on it. Salicylic

acid stimulated growth in various plant species under both normal and varied abiotic stress conditions[29].

Exogenous salicylic acid application diversely impacts plant growth, such as germination of seed, budding, blooming, fruit setting, and ripening. Salicylic acid-induced blooming in finger millet plants[30]. Seed germination of maize and barley was inhibited when infused with more than 3 mM of salicylic acid [31]. However, ingesting maize seeds in 0.3 mM to 0.9 mM salicylic acid resulted in increased shoot length, germination rate, and germination percentage[32]. The strongest germination-stimulating impact was notably shown by 0.43 mM salicylic acid; however, at higher doses, its effect was diminished. So, various salicylic acid concentrations in various plants may either promote or inhibit plant growth.


**Comment [i13]:**

Salicylic acid may exhibit controversial roles in the growth and development of plants, contingent on its concentration, the plant's growing environment, and its stage of development( 28). Elevated levels of salicylic acid can often impede the growth and development of plants (which is contingent upon the plant type; however, concentrations exceeding 1 mM are deemed high). Nevertheless, utilizing appropriate doses of salicylic acid has advantageous effects. Salicylic acid has shown to promote growth in various plant species under both normal and diverse abiotic stress conditions (29).

**Comment [i14]:** seed germination

**Comment [i15]:** Salicylic acid-induced blooming in finger millet plants

**Comment [i16]:** can either promote or inhibit plant growth in various plant species



**Fig. 1. Role of Salicylic acid on growth and development of field crops**

### 3. BOOSTING POWER OF SA ON BIOTIC STRESS TOLERANCE

Salicylic acid is a plant hormone associated with defense that is essential for resistance against several microbial diseases, including fungi, bacteria, viruses, and oomycetes[33]. It is widely known that endogenous salicylic acid levels in plants are positively correlated with resistance mechanisms to both biotrophic and hemibiotrophic diseases [34]. Furthermore, the application of exogenous salicylic acid results in the induction of both local and systemic acquired resistance in a variety of plant species against a range of pathogens, such as *Alternaria alternata*, *Fusarium oxysporum*, *Colletotrichum gloeosporoides*, *Magnaporthe grisea*, *Xanthomonas* spp., various viruses, and so forth [35- 37](Table 1). Notably, the growth of the powdery mildew disease in cucumber plants was almost entirely inhibited by the exogenous application of salicylic acid. Due to its intricacy, salicylic acid's functions in plant defense against necrotrophic diseases are yet unclear. There have been a few reports of exogenous salicylic acid treatment-induced higher sensitivity among various plant-necrotrophic pathogen interactions. Salicylic acid treatment in broad beans reduced red light-induced resistance to the necrotrophic fungus *Botrytis cinerea*, but it did not increase black light-induced vulnerability[38]. Application of tomato SA-induced increased susceptibility in a dose-dependent manner against *B. cinerea*. It is also controversially suggested that salicylic acid increases the resistance of Arabidopsis and tomato plants to *B. cinerea* [39,40].

**Comment [i17]:** defense-related hormone essential for resistance to

**Comment [i18]:** Additionally, the use of exogenous salicylic acid induces local and systemic acquired resistance to several pathogens

**Comment [i19]:** established way

**Table 1. Enhancement of disease resistance mechanism by foliar spray of SA in different plants**

Host	Pathogen	Salicylic acid concentration	Effect	Reference
<i>Oryza sativa</i> (Rice)	<i>Xanthomonas oryzae</i>	1 mM	Reduction of leaf blight lesion	[41]
		1 mM	Reduction of severity of disease (30%)	[42]
	<i>Magnaporthe grisea</i>	8 mM	Reduction of severity of disease (70%)	[43]
		<i>Ooebaluspugnax</i>	16 mM	Reduce the number of bugs (35%)
<i>Cicer arietinum</i> (Chickpea)	<i>Fusarium oxysporum</i>	14.5 mM (stem)	Reduction of severity of disease (20%)	[45]
		0.58 mM (soil)	Reduction of severity of disease (20 %)	
<i>Vigna mungo</i> (Black gram)	<i>Mungbean yellow mosaic Indian virus (MYMIV)</i>	0.1 mM	Reduction of severity of disease (71%)	[36]

### 4. BOOSTING POWER OF SA ON ABIOTIC STRESS TOLERANCE

Plant productivity is threatened by climate change and continuous cropping due to several abiotic stressors, including salt, ozone, UV light, temperature, drought, and heavy metals [46]. It's interesting to note that salicylic acid regulates tolerance to a variety of abiotic stimuli in addition to resistance to biotic challenges[47] (Table 2). The following are the mechanisms of salicylic acid-induced abiotic stress tolerance: (1) osmolytes accumulation, which may support the maintenance of osmotic homeostasis; (2) regulation of the absorption of minerals; (3) increased activity of scavenging reactive oxygen species; (4) increased production of secondary metabolites, comprising nitrogen (alkaloids,

**Comment [i20]:** crop production

**Comment [i21]:** salinity

**Comment [i22]:** It is interesting to note that in addition to resistance to biotic stresses, salicylic acid regulates tolerance to various abiotic stimuli.

**Comment [i23]:** accumulation of osmolytes that can

**Comment [i24]:** regulating minerals absorption

**Comment [i25]:** including

non-protein amino acids, and cyanogenic glucosides,) and sulphur-containing compounds (allinin, glutathione, thionins, phytoalexins, defensins, and glucosinolates) and (5) control of additional hormone pathways [47,48].

A group of pathogenesis-related (*PR*) genes, including *PR1*, *PR2*, and *PR5*, are expressed when exogenous salicylic acid treatment is applied [49]. Transgenic overexpression of several *PR* genes improved tolerance to various abiotic stressors as well as resistance to various infections [50- 52]. Increased resistance to heavy metals was shown by transgenic tobacco that overexpressed pepper *PR-1* [51]. In *Arabidopsis* plants, overexpression of pepper *PR-1* increased resistance to salt and drought stress [50]. Further research is necessary to understand the underlying molecular processes by which these *PR* proteins improve resistance to abiotic stress.

Comment [i26]: upon

Comment [i27]: Additional studies are needed

Comment [i28]: enhance

**Table 2. Improvement of abiotic stress resistance by exogenous application of SA in different plants**

Host	Abiotic stress	Salicylic acid concentration	Effect	Reference
<i>Triticum aestivum</i> (Wheat)	Freezing	0.01, 0.1, and 1 mM	Cell mortality and the loss of PS II quantum yield brought on by freezing stress were dramatically reduced by 0.01 and 0.1 mM salicylic acid.	[53]
<i>Zea mays</i> (Maize)	Cadmium (Cd)	0.5 mM	Root DW and shoot FW are raised by around 121% and 262%, respectively.	[54]
<i>Hordeum vulgare</i> (Barley)	Cadmium (Cd)	0.5 mM	Root DW and shoot FW are raised by around 127% and 133%, respectively.	[55]
	Osmotic stress	30, 60, and 120 nM	Approximately 50% less osmotic stress-induced membrane damage occurred.	[56]

Comment [i29]: Exogenous application of SA in various plants increases their resistance to abiotic stresses.

## 5. SALICYLIC ACID AND PLANT MICROBES

The plant science community has recently shown increased interest in studies examining the relationship between plant health and microbiome [57, 58]. The impact of salicylic acid on the microbiome of the model plant *Arabidopsis thaliana* was examined using either exogenous salicylic acid application or mutants with changed endogenous salicylic acid levels [59]. The results showed that the application of salicylic acid significantly increased the amount of certain bacterial isolates from the Synthetic Community (SynCom) experiment and decreased the amount of *Mitsuaria* sp. 370 ( $\beta$ -Proteobacteria). Furthermore, in *cpr5* mutants that constitutively manufacture salicylic acid, the population density of twelve Proteobacteria and nine Actinobacteria groups was decreased and raised, respectively. This implies that salicylic acid may significantly change the microbiome of the soil or rhizosphere. The stimulation of the systemic immune response has been the primary focus of salicylic

Comment [i30]: the microbiome

Comment [i31]: Results

Comment [i32]: densities of 12 groups of

Comment [i33]: delete

acid's effects on plants so far after soil drench application; however, not much is known regarding the compound's effects on endophytic microbiomes or plant roots. Therefore, further research on the impact of salicylic acid on a variety of soil or endophytic microbiomes will provide insight into how salicylic acid affects many aspects of plant physiology, such as immunity, growth, development, and so on.

**Comment [i34]:** so far been the main effect of salicylic acid

**Comment [i35]:** about the effects of compounds

**Comment [i36]:** on various soil

**Comment [i37]:** different

**Comment [i38]:** delete

## 6. SALICYLIC ACID WITH OTHER PLANT GROWTH REGULATORS (PGRs)

Salicylic acid controls many plant responses through interactions with other plant growth regulators or plant hormones in both favourable and unfavourable environments. Under both ideal and stressful conditions, the relationship between salicylic acid and other hormones, including cytokinin, auxin, gibberellins, abscisic acid, brassinosteroids, and ethylene has been investigated. In stressful situations, the interaction between salicylic acid and hormones may have an antagonistic or synergistic effect. Tamás et al. (2015) [60] recently examined how salicylic acid controlled the reduction of Cd-induced auxin-mediated ROS (reactive oxygen species) generation in barley roots, hence mitigating Cd stress. The authors hypothesize that salicylic acid plays a part in the IAA (indole-3-acetic acid) signaling system since salicylic acid treatment reduces the stress responses that IAA generates in plants. Agtuca et al. (2014) [61] documented that salicylic acid and IAA had opposing roles in maize roots. IAA applied exogenously promoted lateral development by inhibiting primary root growth, while salicylic acid increased the total biomass of roots [61].

**Comment [i39]:** by interacting

**Comment [i40]:** under both favorable and unfavorable conditions

**Comment [i41]:** root biomass

Plants may experience oxidative stress and increased ethylene production when exposed to several environmental conditions, such as heavy metals (HM) [47]. Peaked expression of ethylene-related biosynthetic genes or expression of ethylene-responsive genes is the cause of the enhanced ethylene synthesis. The exogenous spray of salicylic acid helped wheat under Cd stress by raising GSH levels, which led to metal detoxification and scavenged ROS (reactive oxygen species) produced by HM (heavy metal)-triggered ethylene synthesis. Under Cd stress, salicylic acid supplementation led to elevated ABA (Abscisic acid) levels in wheat seedlings, which were linked to the biosynthesis of ABA [62]. Additionally, during HM stress, endogenous ABA regulated SA-mediated changes in the concentration of dehydrin proteins, indicating the protective function of salicylic acid in wheat plants [62].

**Comment [i42]:** Peak

**Comment [i43]:** metals

**Comment [i44]:** Addition of salicylic acid under Cd stress increased abscisic acid (ABA)

**Comment [i45]:** dehydrin protein concentration,

Crosstalk between salicylic acid and jasmonates is essential for controlling plant development in the presence of abiotic stressors [63]. The signaling pathways for jasmonic acid and salicylic acid often function antagonistically. The antagonistic action between salicylic acid and jasmonic acid cell signaling is mediated by the Mitogen-Activated Protein Kinase (MAPK) signaling pathway [64]. Nonantagonistic interactions between salicylic acid and jasmonic acid have also been recorded, although further research is necessary to determine the precise mechanism [63]. Cu stress caused the production of salicylic acid in maize plants, which in turn caused jasmonic acid priming and jasmonic acid-induced volatile organic molecules.

**Comment [i46]:** jasmonate is required to regulate

**Comment [i47]:** effect

**Comment [i48]:** salicylic acid production

## 7. CONCLUSION

Salicylic acid and its derivatives show promise as eco-friendly plant protectors due to their positive impacts on plant and human health. Determining the optimal concentrations, from micromolar to low millimolar levels, is crucial to provide disease resistance without hindering plant growth. Higher 2 mM concentrations may serve as effective growth regulators to slow development for disease control. Exploring natural salicylic acid derivatives, like amorfrutins, with improved efficiency could lead to better plant protection techniques. Further research is needed to understand the practical implementation of salicylic acid across crop species and develop sustainable, cost-effective crop management systems utilizing these versatile compounds.

## REFERENCES

1. Chen Z, Zheng Z, Huang J, Lai Z, Fan B. Biosynthesis of salicylic acid in plants. *Plant signaling & behavior*. 2009;4(6):493-496. DOI: <https://dx.doi.org/10.4161/psb.4.6.8392>
2. Wani AB, Chadar H, Wani AH, Singh S, Upadhyay N. Salicylic acid to decrease plant stress. *Environmental Chemistry Letters*. 2017;15(1):101-123. DOI: <https://dx.doi.org/10.1007/s10311-016-0584-0>
3. Lovelock DA, Šola I, Marschollek S, Donald CE, Rusak G, van Pée KH, Ludwig-Müller J, Cahill DM. Analysis of salicylic acid-dependent pathways in *Arabidopsis thaliana* following infection with *Plasmodiophora brassicae* and the influence of salicylic acid on disease. *Molecular plant pathology*. 2016;17(8):1237-1251. DOI: <https://dx.doi.org/10.1111/mpp.12361>
4. Maruri-López I, Aviles-Baltazar NY, Buchala A, Serrano M. Intra and extracellular journey of the phytohormone salicylic acid. *Frontiers in Plant Science*. 2019;10:455069. DOI: <https://dx.doi.org/10.3389/fpls.2019.00423>
5. Raskin I. Role of salicylic acid in plants. *Annual review of plant biology*. 1992;43(1):439-463.
6. Muthulakshmi S, Lingakumar K. Role of salicylic acid (SA) in plants—A review. *Int. J. Appl. Res.* 2017;3(3):33-37.
7. Petrek J, Havel L, Petrlova J, Adam V, Potesil D, Babula P, Kizek R. Analysis of salicylic acid in willow barks and branches by an electrochemical method. *Russian Journal of Plant Physiology*. 2007;54:553-558. DOI: <https://dx.doi.org/10.1134/S1021443707040188>
8. Sharma A, Sidhu GP, Araniti F, Bali AS, Shahzad B, Tripathi DK, Brestic M, Skalicky M, Landi M. The role of salicylic acid in plants exposed to heavy metals. *Molecules*. 2020;25(3):540. DOI: <https://dx.doi.org/10.3390/molecules25030540>
9. Shine MB, Yang JW, El-Habbak M, Nagyabhyru P, Fu DQ, Navarre D, Ghabrial S, Kachroo P, Kachroo A. Cooperative functioning between phenylalanine ammonia lyase and isochorismate synthase activities contributes to salicylic acid biosynthesis in soybean. *New Phytologist*. 2016;212(3):627-636. DOI: <https://dx.doi.org/10.1111/nph.14078>
10. Zhang Y, Fu X, Hao X, Zhang L, Wang L, Qian H, Zhao J. Molecular cloning and promoter analysis of the specific salicylic acid biosynthetic pathway gene phenylalanine ammonia-lyase(AaPAL1) from *Artemisia annua*. *Biotechnology and Applied Biochemistry*. 2016;63(4):514-524. DOI: <https://dx.doi.org/10.1002/bab.1403>

**Comment [i49]:** Salicylic acid and its derivatives are promising as environmentally friendly plant protection products because of their positive effects on plant and human health. Determining the optimal concentration from micromolar to low millimolar levels is important to provide disease resistance without interfering with plant growth. Higher concentrations of 2mM can act as effective growth regulators to slow development and control disease. Research into natural salicylic acid derivatives, such as amorfrutin, with improved efficacy may lead to the development of more effective plant protection methods. Further research is needed to understand the practical applications of salicylic acid in various crop species and to develop sustainable and cost-effective crop management systems using these versatile compounds.

11. Chong J, Pierrel MA, Atanassova R, Werck-Reichhart D, Fritig B, Saindrenan P. Free and conjugated benzoic acid in tobacco plants and cell cultures. Induced accumulation upon elicitation of defense responses and role as salicylic acid precursors. *Plant Physiology*. 2001;125(1):318-28. DOI: <https://dx.doi.org/10.1104/pp.125.1.318>
12. Hayat S, Ali B, Ahmad A. Salicylic acid: biosynthesis, metabolism and physiological role in plants. *Salicylic acid: A plant hormone*. 2007:1-4.
13. Yalpani N, León J, Lawton MA, Raskin I. Pathway of salicylic acid biosynthesis in healthy and virus-inoculated tobacco. *Plant physiology*. 1993;103(2):315-321. DOI: <https://dx.doi.org/10.1104/pp.103.2.315>
14. Métraux JP. Recent breakthroughs in the study of salicylic acid biosynthesis. *Trends in plant science*. 2002;7(8):332-334. DOI: [https://dx.doi.org/10.1016/S1360-1385\(02\)02313-0](https://dx.doi.org/10.1016/S1360-1385(02)02313-0)
15. Garcion C, Lohmann A, Lamodièrre E, Catinot J, Buchala A, Doermann P, Métraux JP. Characterization and biological function of the *ISOCHORISMATE SYNTHASE2* gene of *Arabidopsis*. *Plant physiology*. 2008;147(3):1279-1287. DOI: <https://dx.doi.org/10.1104/pp.108.119420>
16. Rekhter D, Lüdke D, Ding Y, Feussner K, Zienkiewicz K, Lipka V, Wiermer M, Zhang Y, Feussner I. Isochorismate-derived biosynthesis of the plant stress hormone salicylic acid. *Science*. 2019;365(6452):498-502. DOI: <https://dx.doi.org/10.1126/science.aaw1720>
17. Wei Y, Liu G, Chang Y, He C, Shi H. Heat shock transcription factor 3 regulates plant immune response through modulation of salicylic acid accumulation and signalling in cassava. *Molecular plant pathology*. 2018;19(10):2209-2220. DOI: <https://dx.doi.org/10.1111/mpp.12691>
18. Hartmann M, Zeier J. N-hydroxypipicolinic acid and salicylic acid: a metabolic duo for systemic acquired resistance. *Current opinion in plant biology*. 2019;50:44-57. DOI: <https://dx.doi.org/10.1016/j.pbi.2019.02.006>
19. El-Shazoly RM, Metwally AA, Hamada AM. Salicylic acid or thiamin increases tolerance to boron toxicity stress in wheat. *Journal of plant Nutrition*. 2019;42(7):702-722. DOI: <https://dx.doi.org/10.1080/01904167.2018.1549670>
20. Luo J, Xia W, Cao P, Xiao ZA, Zhang Y, Liu M, Zhan C, Wang N. Integrated transcriptome analysis reveals plant hormones jasmonic acid and salicylic acid coordinate growth and defense responses upon fungal infection in poplar. *Biomolecules*. 2019;9(1):12. DOI: <https://dx.doi.org/10.3390/biom9010012>
21. Pasternak T, Groot EP, Kazantsev FV, Teale W, Omelyanchuk N, Kovrizhnykh V, Palme K, Mironova VV. Salicylic acid affects root meristem patterning via auxin distribution in a concentration-dependent manner. *Plant Physiology*. 2019;180(3):1725-1739. DOI: <https://dx.doi.org/10.1104/pp.19.00130>
22. Dempsey DM, Klessig DF. How does the multifaceted plant hormone salicylic acid combat disease in plants and are similar mechanisms utilized in humans?. *BMC biology*. 2017;15:1-11. DOI: <https://dx.doi.org/10.1186/s12915-017-0364-8>

23. Klessig DF, Choi HW, Dempsey DM. Systemic acquired resistance and salicylic acid: past, present, and future. *Molecular plant-microbe interactions*. 2018;31(9):871-888. DOI: <https://dx.doi.org/10.1094/MPMI-03-18-0067-CR>
24. Subban K, Subramani R, Srinivasan VP, Johnpaul M, Chelliah J. Salicylic acid as an effective elicitor for improved taxol production in endophytic fungus *Pestalotiopsis microspora*. *PloS one*. 2019;14(2):e0212736. DOI: <https://dx.doi.org/10.1371/journal.pone.0212736>
25. Tripathi D, Raikhy G, Kumar D. Chemical elicitors of systemic acquired resistance—Salicylic acid and its functional analogs. *Current Plant Biology*. 2019;17:48-59. DOI: <https://dx.doi.org/10.1016/j.cpb.2019.03.002>
26. Nadeem M, Ahmad W, Zahir A, Hano C, Abbasi BH. Salicylic acid-enhanced biosynthesis of pharmacologically important lignans and neo lignans in cell suspension culture of *Linum usitatissimum* L. *Engineering in Life Sciences*. 2019;19(3):168-74. DOI: <https://dx.doi.org/10.1002/elsc.201800095>
27. Li N, Han X, Feng D, Yuan D, Huang LJ. Signaling crosstalk between salicylic acid and ethylene/jasmonate in plant defense: do we understand what they are whispering?. *International journal of molecular sciences*. 2019;20(3):671. DOI: <https://dx.doi.org/10.3390/ijms20030671>
28. Rivas-San Vicente M, Plasencia J. Salicylic acid beyond defence: its role in plant growth and development. *Journal of experimental botany*. 2011;62(10):3321-3338. DOI: <https://dx.doi.org/10.1093/jxb/err031>
29. Manzoor K, Ilyas N, Batool N, Ahmad B, Arshad M. Effect of Salicylic Acid on the Growth and Physiological Characteristics of Maize under Stress Conditions. *Journal of the Chemical Society of Pakistan*. 2015;37(3).
30. Appu M, Muthukrishnan S. Foliar application of salicylic acid stimulates flowering and induce defense related proteins in finger millet plants. *Universal Journal of Plant Science*. 2014;2(1):14-18. DOI: <https://dx.doi.org/10.13189/ujps.2014.020102>
31. Xie Z, Zhang ZL, Hanzlik S, Cook E, Shen QJ. Salicylic acid inhibits gibberellin-induced alpha-amylase expression and seed germination via a pathway involving an abscisic-acid-inducible WRKY gene. *Plant molecular biology*. 2007;64:293-303. DOI: <https://dx.doi.org/10.1007/s11103-007-9152-0>
32. Sallam AM, Ibrahim HI. Effect of grain priming with salicylic acid on germination speed, seedling characters, anti-oxidant enzyme activity and forage yield of teosinte. *American-Eurasian Journal of Agricultural & Environmental Sciences*. 2015;15(5):744-753.
33. Vlot AC, Dempsey DM, Klessig DF. Salicylic acid, a multifaceted hormone to combat disease. *Annual review of phytopathology*. 2009;47:177-206. DOI: <https://dx.doi.org/10.1146/annurev.phyto.050908.135202>
34. Glazebrook J. Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. *Annu. Rev. Phytopathol.* 2005;43:205-227. DOI: <https://dx.doi.org/10.1146/annurev.phyto.43.040204.135923>

35. Jendoubi W, Harbaoui K, Hamada W. Salicylic acid-induced resistance against *Fusarium oxysporum* f. sp. *pradicislycopercisi* in hydroponic grown tomato plants. *Journal of New Sciences*. 2015;21.
36. Kundu S, Chakraborty D, Pal A. Proteomic analysis of salicylic acid induced resistance to Mungbean Yellow Mosaic India Virus in *Vigna mungo*. *Journal of proteomics*. 2011;74(3):337-349. DOI: <https://dx.doi.org/10.1016/j.jprot.2010.11.012>
37. Le Thanh T, Thumanu K, Wongkaew S, Boonkerd N, Teamroong N, Phansak P, Buensanteai N. Salicylic acid-induced accumulation of biochemical components associated with resistance against *Xanthomonas oryzae* pv. *oryzae* in rice. *Journal of Plant Interactions*. 2017;12(1):108-120. DOI: <https://dx.doi.org/10.1080/17429145.2017.1291859>
38. Khanam NN, Ueno M, Kihara J, Honda Y, Arase S. Suppression of red light-induced resistance in broad beans to *Botrytis cinerea* by salicylic acid. *Physiological and molecular plant pathology*. 2005;66(1-2):20-29. DOI: <https://dx.doi.org/10.1016/j.pmpp.2005.03.006>
39. Ferrari S, Plotnikova JM, De Lorenzo G, Ausubel FM. *Arabidopsis* local resistance to *Botrytis cinerea* involves salicylic acid and camalexin and requires *EDS4* and *PAD2*, but not *SID2*, *EDS5* or *PAD4*. *The Plant Journal*. 2003;35(2):193-205. DOI: <https://dx.doi.org/10.1046/j.1365-313X.2003.01794.x>
40. Li L, Zou Y. Induction of disease resistance by salicylic acid and calcium ion against *Botrytis cinerea* in tomato (*Lycopersicon esculentum*). *Emirates Journal of Food and Agriculture*. 2017:78-82. DOI: <https://dx.doi.org/10.9755/ejfa.2016-10-1515>
41. Mohan Babu R, Sajeena A, VijayaSamundeeswari A, Sreedhar A, Vidhyasekaran P, Seetharaman K, Reddy MS. Induction of systemic resistance to *Xanthomonas oryzae* pv. *oryzae* by salicylic acid in *Oryza sativa* (L.). *Journal of Plant Diseases and Protection*. 2003;110:419-431. DOI: <https://dx.doi.org/10.1007/bf03356119>
42. Le Thanh T, Thumanu K, Wongkaew S, Boonkerd N, Teamroong N, Phansak P, Buensanteai N. Salicylic acid-induced accumulation of biochemical components associated with resistance against *Xanthomonas oryzae* pv. *oryzae* in rice. *Journal of Plant Interactions*. 2017;12(1):108-120. DOI: <https://dx.doi.org/10.1080/17429145.2017.1291859>
43. Daw BD, Zhang LH, Wang ZZ. Salicylic acid enhances antifungal resistance to *Magnaporthe grisea* in rice plants. *Australasian Plant Pathology*. 2008;37:637-644. DOI: <https://dx.doi.org/10.1071/AP08054>
44. Stella de Freitas TF, Stout MJ, Sant'Ana J. Effects of exogenous methyl jasmonate and salicylic acid on rice resistance to *Oebaluspugnax*. *Pest management science*. 2019;75(3):744-752. DOI: <https://dx.doi.org/10.1002/ps.5174>
45. Saikia R, Singh T, Kumar R, Srivastava J, Srivastava AK, Singh K, Arora DK. Role of salicylic acid in systemic resistance induced by *Pseudomonas fluorescens* against *Fusarium oxysporum* f. sp. *ciceri* in chickpea. *Microbiological Research*. 2003;158(3):203-213. DOI: <https://dx.doi.org/10.1078/0944-5013-00202>
46. Connor D. Climate change and global crop productivity. *Crop Science*. 2002;42(3):978-979. DOI: <https://dx.doi.org/10.2135/cropsci2002.9780>

47. Khan MI, Fatma M, Per TS, Anjum NA, Khan NA. Salicylic acid-induced abiotic stress tolerance and underlying mechanisms in plants. *Frontiers in plant science*. 2015;6:135066. DOI: <https://dx.doi.org/10.3389/fpls.2015.00462>
48. Horváth E, Csiszár J, Gallé Á, Poór P, Szepesi Á, Tari I. Hardening with salicylic acid induces concentration-dependent changes in abscisic acid biosynthesis of tomato under salt stress. *Journal of Plant Physiology*. 2015;183:54-63. DOI: <https://dx.doi.org/10.1016/j.jplph.2015.05.010>
49. Ali S, Ganai BA, Kamili AN, Bhat AA, Mir ZA, Bhat JA, Tyagi A, Islam ST, Mushtaq M, Yadav P, Rawat S. Pathogenesis-related proteins and peptides as promising tools for engineering plants with multiple stress tolerance. *Microbiological research*. 2018;212:29-37. DOI: <https://dx.doi.org/10.1016/j.micres.2018.04.008>
50. Hong JK, Hwang BK. Induction of enhanced disease resistance and oxidative stress tolerance by overexpression of pepper basic PR-1 gene in *Arabidopsis*. *Physiologia Plantarum*. 2005;124(2):267-277. DOI: <https://dx.doi.org/10.1111/j.1399-3054.2005.00515.x>
51. Sarowar S, Kim YJ, Kim EN, Kim KD, Hwang BK, Islam R, Shin JS. Overexpression of a pepper basic pathogenesis-related protein 1 gene in tobacco plants enhances resistance to heavy metal and pathogen stresses. *Plant cell reports*. 2005;24:216-224. DOI: <https://dx.doi.org/10.1007/s00299-005-0928-x>
52. Wu W, Ding YA, Wei W, Davis RE, Lee IM, Hammond RW, Zhao Y. Salicylic acid-mediated elicitation of tomato defence against infection by potato purple top phytoplasma. *Annals of Applied Biology*. 2012;161(1):36-45. DOI: <https://dx.doi.org/10.1111/j.1744-7348.2012.00550.x>
53. Wang W, Wang X, Huang M, Cai J, Zhou Q, Dai T, Cao W, Jiang D. Hydrogen peroxide and abscisic acid mediate salicylic acid-induced freezing tolerance in wheat. *Frontiers in plant science*. 2018;9:1137. DOI: <https://dx.doi.org/10.3389/fpls.2018.01137>
54. Krantev A, Yordanova R, Janda T, Szalai G, Popova L. Treatment with salicylic acid decreases the effect of cadmium on photosynthesis in maize plants. *Journal of plant physiology*. 2008;165(9):920-931. DOI: <https://dx.doi.org/10.1016/j.jplph.2006.11.014>
55. Metwally A, Finkemeier I, Georgi M, Dietz KJ. Salicylic acid alleviates the cadmium toxicity in barley seedlings. *Plant physiology*. 2003;132(1):272-281. DOI: <https://dx.doi.org/10.1104/pp.102.018457>
56. Bandurska H, Stroi ski A. The effect of salicylic acid on barley response to water deficit. *Acta Physiologiae Plantarum*. 2005;27:379-386. DOI: <https://dx.doi.org/10.1007/s11738-005-0015-5>
57. Bulgarelli D, Rott M, Schlaeppi K, Ver Loren van Themaat E, Ahmadinejad N, Assenza F, Rauf P, Huettel B, Reinhardt R, Schmelzer E, Peplies J. Revealing structure and assembly cues for *Arabidopsis* root-inhabiting bacterial microbiota. *Nature*. 2012;488(7409):91-95. DOI: <https://dx.doi.org/10.1038/nature11336>
58. Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S, Tremblay J, Engelbrektson A, Kunin V, Rio TG, Edgar RC. Defining the core *Arabidopsis thaliana* root microbiome. *Nature*. 2012;488(7409):86-90. DOI: <https://dx.doi.org/10.1038/nature11237>

59. Lebeis SL, Paredes SH, Lundberg DS, Breakfield N, Gehring J, McDonald M, Malfatti S, Glavina del Rio T, Jones CD, Tringe SG, Dangl JL. Salicylic acid modulates colonization of the root microbiome by specific bacterial taxa. *Science*. 2015;349(6250):860-864. DOI: <https://dx.doi.org/10.1126/science.aaa8764>
60. Tamás L, Mistrík I, Alemayehu A, Zelinová V, Bočová B, Huttová J. Salicylic acid alleviates cadmium-induced stress responses through the inhibition of Cd-induced auxin-mediated reactive oxygen species production in barley root tips. *Journal of Plant Physiology*. 2015;173:1-8. DOI: <https://dx.doi.org/10.1016/j.jplph.2014.08.018>
61. Agtuca B, Rieger E, Hilger K, Song L, Robert CA, Erb M, Karve A, Ferrieri RA. Carbon-11 reveals opposing roles of auxin and salicylic acid in regulating leaf physiology, leaf metabolism, and resource allocation patterns that impact root growth in *Zea mays*. *Journal of Plant Growth Regulation*. 2014;33:328-339. DOI: <https://dx.doi.org/10.1007/s00344-013-9379-8>
62. Shakirova FM, Allagulova CR, Maslennikova DR, Klyuchnikova EO, Avalbaev AM, Bezrukova MV. Salicylic acid-induced protection against cadmium toxicity in wheat plants. *Environmental and experimental botany*. 2016;122:19-28. DOI: <https://dx.doi.org/10.1016/j.envexpbot.2015.08.002>
63. Per TS, Khan MI, Anjum NA, Masood A, Hussain SJ, Khan NA. Jasmonates in plants under abiotic stresses: Crosstalk with other phytohormones matters. *Environmental and experimental botany*. 2018;145:104-120. DOI: <https://dx.doi.org/10.1016/j.envexpbot.2017.11.004>