

Abstract

Climate change causes various negative effects on plants, especially due to rapid changes in temperature, rainfall patterns, floods or droughts, and outbreaks of pests and diseases. Its change is predicted to cause widespread species expansion and extinction. Climate-induced local extinction has already occurred in hundreds of species. However, at the edge of the warm zone, an equal number of species were not exposed to local extinction, indicating that either phenotypic plasticity or genetic adaptations may allow some populations to persist in warmer conditions. This shows the importance of including intraspecific adaptations in climate change vulnerability assessments. In addition, in response to global climate change, the application of gene editing, also known as genome editing or genome engineering, has emerged as a technology to help organisms adapt to global climate change or mitigate climate impacts. Transforming agriculture by developing crops and livestock that can better withstand the effects of climate change is imperative. Gene editing allows precise changes to a plant's genome, speeding up the production of new crop varieties, including those better able to withstand the stress of a changing climate and those that capture and store excess atmospheric carbon dioxide. The precision and efficiency of creating changes has greatly improved with the introduction of CRISPR/Cas systems, although there is certainly more work to be done with other gene editing techniques

Key words : Biotechnology, Climate Change, CRISPER/Cas, Drought, Gene editing,

1. Introduction

“According to the Intergovernmental Panel on Climate Change (IPCC) group, climate change is the average change or variation in temperature, precipitation and wind patterns over the long term. According to the IPCC report, climate change is largely caused by anthropogenic causes, including human-induced changes in land use, as well as natural forces, such as solar cycles, volcanic eruptions, and continental drift” (IPCC, 2014).

Most effects of climate change concern soil and water quality, temperature for pests and pathogens, precipitation and weeds. Research data showed that agriculture is responsible for 25% of greenhouse gases and the main sources are methane (48%) and nitrous oxide (52%) from rice fields (Sallema and Mtui, 2008). “Greenhouse gases are both natural and man-made elements that prevent the reflection of radiation in the atmosphere and the warming of the environment. These gases are mainly emitted by industry and other activities, such as carbon dioxide (CO₂), methane (CH₄), nitrous oxide, hydrofluorocarbons (HFC) and sulfur dioxide (SF₆). In the long term, various activities increase their concentration in the atmosphere and lead to global climate change” (Kumar et al., 2015).

According to the United States Geological Survey, global warming is one of the aspects of climate change, and according to them, global warming is an increase in the temperature of the earth, which is mainly caused by an increase in the concentration of greenhouse gases in the atmosphere. Due to climate change, arable land is decreasing due to soil erosion, desertification and salinization. Drought threatens agriculture worldwide more than ever before. “The Food and Agriculture Organization of the United Nations (FAO) documents that between 2005 and 2015, drought caused USD 29 billion in direct damage to agriculture in developing countries. In addition, more than 70% of the world's available fresh water is used for irrigation (Organization for Economic Cooperation and Development, 2017)”. [140]

Climate change is predicted to cause widespread distribution and extinction of species (Urban, 2015), and climate-related local extinctions have already occurred in hundreds of species. However, an equal number of species at the edge of the warm zone did not experience local extinction (Wiens, 2016), indicating that either phenotypic plasticity or genetic adaptations may allow some populations to persist in warmer conditions. This shows that it is important to include intraspecies adaptations in the assessment of vulnerability to climate change (Savolainen et al., 2013; Bay et al., 2018). Biotechnology, especially genetic engineering, gives crops a greater advantage in response to stress than traditional breeding. In addition, genetically modified soybeans, corn and cotton produced for insect resistance and weed tolerance have made very impressive and dramatic progress in pest control and crop improvement worldwide since their first introduction in 1996.

Climate change mitigation strategies in generalis immediate goal to reduce the negative impact of environmental problems on land and water bodies. For example, reducing the concentration of greenhouse gases that can emitted into the atmosphere by limiting the emission of industrial sewage and radioactive substances can be some. measures that protect the earth from the effects of climate change. In addition, climate change can be contained through afforestation and other sinks (natural absorbers and adsorbents). Biotechnological methods currently used for mitigation purposes include tissue culture, bioremediation, biosorption, bioleaching, conventional breeding, molecular marker breeding, genetic engineering, and genome editing, Carbon reduction through biofuels, carbon sequestration, use of inorganic fertilizers etc. are other processes.

Hence, the purpose of this reviwie paper is give highlights the impact of biotechnology on climate change to address pressing environmental challenges through novel approaches that can operate at the scale and efficiency to describe the importance of biotechnology and its intervention in potential disasters resulting from climate change to avoid irreversible damage.

2. Therole of genetic variation for adaptation to climate change

“Genetic diversity is the most important requirement for all species to survive in the long term and adapt to environmental changes over evolutionary time” (Falk *et al.*, 2001; Frankham, 2005).

“Genetic structure is very important because it can provide insight into the history of a population, and the current levels and distribution of genetic variation can influence the future success of populations” (Erickson *et al.*, 2004).

“In some combination of natural selection and random genetic drift, a population separated by geographic distance may diverge due to gene flow and reduced population connectivity (isolation by geographic distance - IBD)” (Nosil and Rundle, 2012). “However, population differences can emerge when reproductive isolation develops between neighboring populations in different environments as a result of ecologically based differential selection from isolating environment (IBE)” (Wang and Bradburd, 2014).

“Global climate change has become one of the greatest threats to biodiversity” (Davis and Shaw, 2001; Parmesano, 2006). Species may respond to global climate change through local adaptation (Parmesano, 2006), individual migration (Breshears *et al.*, 2008; Lenoir *et al.*, 2008), range reduction (Thiller *et al.*, 2005), or a combination of these (Margaret *et al.*, 2005).

“Local adaptations have been found to be a traditional way for various plant species to respond to climate change” (Coop *et al.*, 2010; Gonzalez-Martinez *et al.*, 2006; Hancock *et al.*, 2011; Savolainen *et al.*, 2007).

In addition, projected global warming will have dramatic effects on mountain ecosystems (Kohler *et al.*, 2010), especially alpine plant communities (Gottfried *et al.*, 2012). “Vulnerability to climate change is most often assessed based on projected distributional changes using ecological niche modeling. This model can predict future changes in the distribution of suitable climatic conditions that characterize the current range of the species” (Bay *et al.*, 2017). “An important limitation of these models, which can lead to incorrect predictions and misplaced conservation efforts, is the neglect of adaptation to internal climate and the resulting differences in population response to climate change” (Shafer *et al.*, 2015). “Evidence of contrasting patterns of physiological variation in thermal tolerance between and within species underscores the importance of including within-species variation in ecological niche models (ENM) of climate adaptation” (Pacifi *et al.*, 2015). “However, such model improvements are limited by the scarcity of observational and experimental studies on local climate adaptation” (Hällfors, *et al.*, 2016).

“To date, studies attempting to incorporate genetic variation into an ecological niche model (ENM) have mostly used a neutral marker to identify phylogeographic structure and create separate models for each genetically distinct population. They led to more pessimistic projections than conventional ENMs, predicting an increased risk of climate change due to the loss of vulnerable populations, but did not affect projections of range size changes at the species level” (Slaty *et al.*, 2016 and Valladares *et al.*, 2014). “These experiments are limited in scope because neutral markers provide information about the evolutionary history of species and barriers to gene flow, but not about the ability of individuals to adapt and survive under changing conditions. In addition, range shifts caused by future climate change are expected to result in the

genetic homogenization of different species and the disappearance of historical and current population subdivisions” (Ikeda *et al.*, 2017).

“Recent studies integrate genomic adaptations into ENM projections to identify vulnerable populations that need to adapt to survive future climate change” (D’Amenet *et al.*, 2017;Paulsetal., 2013). “However, genetic information related to interspecific variation in climate adaptation has yet to be directly incorporated into ENMs. To overcome these challenges, plant breeders began to produce new crop varieties that increased yield, tolerate abiotic stresses and improved water and nutrient consumption efficiency” (Fita *et al.*, 2015).

3Adaptation of agriculture to climate change through plant breeding

“In agriculture, drought can generally be defined as a prolonged lack of water that affects plant growth and survival, ultimately reducing crop productivity. In botany, the broadest definition of drought stress is the same as water deficit, which occurs when the rate of evaporation exceeds water consumption” (Bray, 1997). “This can be caused by a lack of water, but also increased salinity or osmotic pressure. From a molecular biology point of view, the first event during drought stress is water loss from the cell or dehydration. Desiccation typically triggers osmotic signals and hormones mainly related to abscisic acid (ABA)” (Blum, 2015).

“Drought resistance is determined by how effectively and timely the plant senses changing environmental conditions and combining the environmentalstreesin response to reduced water availability. Plant breeders have identified physiological traits that result from the drought response and facilitate plant adaptation to limited water. Understanding the molecular and physiological mechanisms underlying these traits is important for crop improvement through biotechnology”. [140] Biotechnology is a promising way to mitigate the negative effects of climate change by reducing greenhouse gases through biofuels (Lybbertand Sumner D, 2010) and carbon sequestration (Kleteretal. , 2008), less fertilizer (Yan *et al.*, 2008), biotic tolerance (Hsieh *et al.*, 2002) and biotic stress (Barrows *et al.*, 2014)

4.The role of microbes in climate change resilience

“Microbes are various organisms found on the surface of the earth. Plants themselves consist of many microbes found in them and in the soil ecosystem. Microbes are known to perform

various ecological functions in nature. They regulate the concentration of greenhouse gases and influence the radiative forcing. Microbiota can influence both positive and negative feedbacks on climate tolerance. Several microbial species play important roles in carbon sequestration, carbon minimization and reduction of greenhouse gases such as CO₂, CH₄ and N₂O in the soil ecosystem”.(Singh et al., 2010) “Microbes such as bacteria and fungi effectively break down organic matter, which further stimulates global warming in the environment and the flow of carbon dioxide into the atmosphere. Microbial communities influence the biogeochemical cycle, nutrient cycling, carbon and methane cycle status in the atmosphere” (Abatenhet *al.*, 2018).

“Microbial respiration is a key pathway for carbon dioxide efflux that promotes the natural release of carbon dioxide. On the other hand, methanotrophs play an important role as biological sinks that reduce methane emissions to the atmosphere. Microbial respiration is an important carbon dioxide emission pathway that contributes to the natural release of carbon dioxide.. The plant microbiome also contributes to global food security by determining yield and climate resilience. Climate change mitigation is a necessary measure that can be achieved through several means. The use of biofertilizers composed of microorganisms such as bacteria and fungi can be an effective alternative to chemical fertilizers, as well as the use of biofuels instead of fossil fuels. Soil contains many microorganisms. Soil microbes play important roles in nutrient cycling, resistance to soil-borne pathogens, and regulation of climate change. Soil contains many microorganisms. Soil microbes participate in the decomposition of soil organic matter, regulate carbon supplies and nutrient cycling, and facilitate plant nutrient assimilation” (Delgado *et al.*; 2020).

“Modern green technologies such as biofertilizers composed of cyanobacteria, fungi (*arbuscular mycorrhiza*, AMF) and bacteria (plant growth-promoting *rhizobacteria*, PGPR) improve and restore soil fertility and ensure sustainable agricultural production. In addition, these microorganisms can reduce energy requirements in the form of synthetic fertilizers and have the ability to alleviate stressed agricultural ecosystems and desert lands” (Du *et al.*, 2007; Mohammadi and Sorabi, 2012).

“Sustainable agriculture includes soil, water and pest management, crop selection and soil conservation. This practice of sustainable agriculture combined with biotechnology can

increase productivity by creating new transgenic plants, microbes and animals” (Singh, 2000). “Cyanobacteria produce a number of valuable compounds such as ethanol, butanol, fatty acids and other organic acids and are promising candidates to continuously satisfy our energy needs. Recent advances in cyanobacterial genetic engineering, cultivation, and culture screening have enabled new ways to exploit the riches of these ancient microorganisms. Gene manipulation techniques are well developed for several cyanobacteria” (Rajineeshet *et al.*, 2017).

4.1. The role of cyanobacteria in climate change

Cyanobacteria play an important role in atmospheric nitrogen fixation and carbon fixation and sequestration (Mishra *et al.*; 2019; Schipper *et al.*; 2019), which are essential for plant nutrition and soil fertility.

Among nitrogen fixer cyanobacteria, *Oscillatoria*, *Nostoc*, *Anabaena*, etc., have a potential role in combating stress conditions in various plant species.

In terms of carbon dioxide capture, one of the most promising organisms is a cyanobacterium (Berla *et al.*, 2013; Nozzi *et al.*, 2013; Case and Atsumi, 2016). These photosynthetic bacteria also improve the activity and diversity of the microbial community through symbiotic associations (Adams *et al.*; 2013) and in addition to EPS, cyanobacteria can secrete several acids, hormones, amino acids and vitamins that promote plant growth and development (Mohan *et al.*; 2019). Compared to the growth of other plants with beneficial bacteria (PGPB), after their death and decomposition, cyanobacteria can increase the water holding capacity and soil biomass (Rady *et al.*; 2018) In addition, the ability of cyanobacteria to tolerate different salinities. reduces the fresh water for their cultivation, strengthening their values as NBS (Kirsch *et al.*; 2019) .

4.1.1. Cyanobacteria as a source of bioenergy

“First and second generation biofuels use raw materials such as rapeseed, soybeans, sunflower, wheat, grass, peanuts and sesame. Various energy sources such as ethanol, propanol, butanol and vegetable oils have been produced from these raw materials” (Quintana *et al.*, 2011). “However, energy crops used in the production of first and second generation biofuels compete with conventional food sources for water, nutrients and fertile

land. Therefore, the third generation using microalgae has emerged as an alternative to biofuels to avoid competition between food crops and energy crops for available natural resources, and in addition, cyanobacteria are one of the most promising raw materials for the production of third generation biofuels” (Quintana *et al.*, 2011; Al-Haj *et al.*, 2016; Rajneesh *et al.*, 2017).

“Rapid growth and cultivation in suitable indoor bioreactors and/or non-cultivable soils gives cyanobacteria an advantage over plants” (Singh, 2014; Sarma *et al.*, 2016). “In addition, cyanobacteria show higher photosynthetic efficiency (~10%) compared to land plants (maximum efficiency ~3-4%)” (Lewis and Nocera, 2006; Melis, 2009). “Blue-green algae are easier to genetically manipulate than other algae and are therefore better candidates for the production of chemicals and fuels compared to eukaryotic algae. The genome size of cyanobacteria is relatively small and the genomes of several genera have been sequenced so far” (Rajneesh *et al.*, 2017). “Therefore, cyanobacteria offer an exceptional opportunity for genetic and metabolic engineering research to improve biomass production, which is relatively difficult to do with eukaryotic algae” (Rittmann, 2008).

“Cyanobacteria contain significant amounts of lipids; located mainly in the thylakoids and plasma membranes and have greater growth and photosynthesis. Biofuel improvement of cyanobacteria using genetic engineering has been attempted mainly with *Synechocystis sp. PCC 6803* and *S. elongatus PCC 7942*, whose genomes were completely sequenced and established by molecular techniques” (Kaneko *et al.*, 1996). “Different fuels such as 2,3-butanediol, acetone-1-butanol, ethylene, ethanol, fatty acids, isobutyraldehyde, isobutanol, 2-methyl-1-butanol and isoprene can be produced in cyanobacteria using genetic engineering. Therefore, genetically engineered cyanobacteria can play a crucial role in reducing oil dependence and CO₂ emissions, since CO₂ is directly photosynthetically linked to biofuels and other valuable secondary metabolites” (Ducat *et al.*, 2011; Oliver *et al.*, 2016). “However, the use of cyanobacteria for biofuel production has some limitations. Production of valuable chemicals in photoautotrophic cyanobacteria is always lower than in sugar-using systems such as *S. cerevisiae* and *E. coli*” (Savakis and Hellingwerf, 2014).

In general, a photoautotrophic cyanobacterial body can only produce ~100 mg of biochemicals per liter of cell culture (Gao *et al.*, 2016), which is too little for a commercially

viable application. “Theoretical yields for the production of several chemicals under heterotrophic and autotrophic growth conditions were calculated for the cyanobacterial body to explain the limiting factors of the cyanobacterial metabolic network” (Gudmundsson and Nogales, 2015). “But the study suggests that the low performance is not due to the topology of the photosynthetic metabolic networks of the cyanobacteria. Therefore, it is important to optimize the natural biological framework to increase the yield of cyanobacteria-derived biochemicals. In recent years, several groups have emphasized the construction, design, and expression of biosynthetic pathways and the development of cyanobacterial metabolic engineering tools that can lead to economic viability by increasing the production of existing and new chemicals and biofuels” (Wang *et al.*, 2012; Berla *et al.*, 2013; Desai and Atsumi, 2013; Oliver and Atsumi, 2014; Gudmundsson and Nogales, 2015; Markley *et al.*, 2015).

4.1.2 Cyanobacteria as biofertilizer

“The production of inorganic nitrogen fertilizers is very expensive because it uses a lot of fossil fuel energy. This required the development of alternative, sustainable and cost-effective bioavailable nitrogen resources that can sustainably meet the nitrogen demand of agriculture” (Mahanty *et al.*, 2017). “To this end, biological systems capable of fixing atmospheric nitrogen have been identified” (Hegde *et al.*, 1999; Vaishampayan *et al.*, 2001).

Biological N fixation produces $\sim 2 \times 10^2$ Mt N per year (Guerrero *et al.*, 1981). According to Metting (1988), “total nitrogen fixation can be ~ 90 kg N ha⁻¹ y⁻¹. Symbiotic and free-living eubacteria, including cyanobacteria, are two groups of nitrogen-fixing organisms”. “Free-living cyanobacteria fix 10 kg N ha⁻¹ a⁻¹, but ~ 10 –30 kg N ha⁻¹ per year is fixed by dense cyanobacterial worms” (Aiyer *et al.*, 1972; Watanabe *et al.*, 1977). “Therefore, cyanobacteria are an important component in naturally available biofertilizers” (Vaishampayan *et al.*, 2001; Prasanna *et al.*, 2013). “Rice production in tropical countries depends mainly on biological N₂ fixation by cyanobacteria, which is a natural part of rice

fields” (Vaishampayan *et al.*, 2001). In these cultivated agricultural systems, ~32 Tg of N are fixed annually by biological N-fixers (Singh *et al.*, 2016), and cyanobacteria add approximately 20–30 kg of N fixed ha⁻¹ to rice fields along with organic matter (Subramanian and Sundaram, 1986; Issa *et al.*, 2014). “Cyanobacteria also form symbiotic associations with various photosynthetic and non-photosynthetic organisms such as algae, fungi, diatoms, hornworts, liverworts, mosses, pteridophytes and angiosperms” (Rai *et al.*, 2000; Sarma *et al.* 2006).

“Cyanobacteria show a higher photosynthetic efficiency ((~10%) compared to land plants (~3-4% maximum efficiency))” (Lewis and Nocera, 2006; Melis., 2009). “Cyanobacteria can be genetically manipulated more easily than other algae and thus, act as target chemicals and better than eukaryotic algae in fuel production. The genome size of cyanobacteria is relatively small and the genomes of several genera have been sequenced so far” (Rajneesh *et al.*, 2017).

“Therefore, cyanobacteria can develop genetic and metabolic technology to produce biomass, which is relatively difficult for eukaryotic algae to do” (Rittmann, 2008). This species contains significant amounts of lipids; especially is located in thylakoids and plasma membranes and shows faster growth and photosynthesis.

“The production of biofuels from cyanobacteria through genetic engineering was observed mainly in *synechocystis* sp. pcc6803 and *s.elongatus* pcc7942, whose genomic DNA was completely sequenced” (Kaneko *et al.*, 1996). “Thus, genetically enhanced cyanobacteria can play a crucial role in reducing oil dependence and carbon dioxide emissions, because carbon dioxide photosynthetically combines directly with biofuels and other valuable secondary metabolic products” (Ducat *et al.*, 2011; Oliver *et al.*, 2016).

“Genera of heterocytic cyanobacteria such as *Nostoc*, *Anabaena*, *Nodularia*, *Scytonema*, *Cylindrospermum*, *Mstigocladus*, *Calothrix*, *Anabaenopsis*, *Aulosira*, *Tolypothrix*, *Haplosiphon*, *Campytonema*, *Stigonema*, *Fischerella*, *Chlopsiofix*, *Westrogstiarog*” (Vaishampayan, 1993). **Lists of possible cyanobacteria that can be used as biofertilizers in agricultural fields (Vaishampayan *et al.*, 2001) are indicated Table 1.** Mixed cultures of free-living forms of cyanobacteria are used to propagate rice fields

(Venkataraman, 1972; Roger and Kulasooriya, 1980). “The water fern *Azolla* carries *Anabaena azollae* on its fronds, and the cyanobacterium releases ammonium into water when rice fields are inoculated with foam-immobilized strains of *A. atolli*” (Kannaiyan *et al.*, 1997). “Significant increases in grain yield, biomass and nutritive value of rice can be achieved by inoculating *Anabaena doliolum* and *A. fertilissima* in rice fields with or without urea” (Dubey and Rai, 1995). “Several species of cyanobacteria such as *Anabaena iyengarii* var. considered, *A. fertilissima*, *Nostoc community*, *N. elliposporum*, *N. linckia* and *Gloeotrichianatans* can contribute to the productivity of rice fields in Chile” (Pereira *et al.*, 2009). “In general, 12.5 kg ha⁻¹ cyanobacterial biofertilizers are recommended for quantitative and qualitative improvement of rice production” (Dubey and Rai, 1995).

Table 1. List of nitrogen-fixing cyanobacteria important for their application in biofertilizer industry (adapted from Vaishampayan *et al.*, 2001).

Filamentous		Unicellular
Heterocystous	Non-heterocystous	
<i>Anabaena</i> , <i>Anabaenopsis</i> ,	<i>Lyngbya</i> , <i>Microcoleus</i>	<i>Aphanothece</i> ,
<i>Aulosira</i> , <i>Calothrix</i> , <i>Camptylonema</i> ,	<i>chthonoplastes</i> ,	<i>Chroococciopsis</i> ,
<i>Chlorogloea</i> , <i>Chlorogloeopsis</i> ,	<i>Myxosarcina</i> ,	<i>Dermocarpa</i> ,
<i>Cylindrospermum</i> , <i>Fischerella</i> ,	<i>Oscillatoria</i> , <i>Plectonema</i>	<i>Gloeocapsa</i> ,
<i>Gloeotrichia</i> , <i>Haplosiphon</i> ,	<i>Boryanum</i> ,	<i>Myxosarcina</i> ,
<i>Mastigocladus</i> , <i>Nodularia</i> , <i>Nostoc</i> ,	<i>Pseudoanabaena</i> ,	<i>Pleurocapsa</i> ,
<i>Nostochopsis</i> , <i>Rivularia</i> ,	<i>Schizothrix</i> ,	<i>Synechococcus</i> ,
<i>Scytonema</i> , <i>Scytonematopsis</i> ,	<i>Trichodesmium</i>	<i>Xenococcus</i>
<i>Stigonema</i> , <i>Tolypothrix</i> , <i>Westiella</i> ,		
<i>Westiellopsis</i> , <i>Wollea</i>		

In addition to rice yield, cyanobacterial biofertilization can also improve wheat yield, shoot/root length and dry weight (Spiller and Gunasekaran, 1990; Obreht *et al.*, 1993; Karthikeyan *et al.*, 2009). Soil inoculation with different cyanobacterial strains such as *Nostoc carneum*, *N.piscinale*, *Anabaena doliolum* and *A.torulosa* results in significantly higher acetylene-reducing activity (Prasanna *et al.*, 2013).

Moreover, acetylene reducing activity is highest in harvest phase when wheat field is inoculated with *Anabaena-serratia* biofilm along with rock phosphate (Swarnalakshmi *et al.*, 2013). Biofertilizers based on cyanobacteria are one third more cost-effective than chemical fertilizers (Prasanna *et al.*, 2013). In addition to nitrogen fixation, cyanobacteria also contribute to the mobilization of inorganic phosphates through the secretion of organic acids and extracellular phosphatases (Bose and Nagpal, 1971; Rai and Sharma, 2006). Cyanobacteria solubilize and mobilize insoluble organic phosphatase and improve phosphorus availability to crops (Dorich *et al.*, 1985; Cameron and Julian, 1988). The humus content created after the death and decomposition of cyanobacteria creates strong reducing conditions in the soil, which improve soil structure and fertility (Abdel-Raouf *et al.*, 2012).

Various cyanobacterial strains can produce plant growth hormones and siderophores, and therefore cyanobacteria can affect crop development and productivity (Rodriguez *et al.*, 2006; Rastogi and Sinha, 2009). EPS secreted by cyanobacteria induces soil aggregation. items that improve soil structure and fertility by increasing accumulation. Together, these findings confirm the importance of cyanobacteria as biofertilizers, and methods have been developed for their cultivation and use in the fertilizer industry (Brouers *et al.*, 1987; Shi and Hall, 1988; Vaishampayan *et al.*, 2000).

4.1.3. The role of myco-biotechnology in climate change

Mycobiotechnology is the use of fungi to create various products. These methods use fungi to restore damaged ecology that endo- and ectomycorrhizal symbiotic fungi with actinomycetes have been used as inoculants to restore degraded forests. Mycorestoration tries to use mushrooms as an aid in the restoration of an ecologically weakened environment. Whether the environment has been damaged by a man-made or natural disaster, saprophytic and mycorrhizal fungi can help it recover. Afforestation would indirectly improve product safety and food security, as forests create a microclimate that can improve availability.


Fungal applications of biotechnology, called mycobiotechnology, are part of a broader trend of using living systems to solve environmental problems and restore degraded ecosystems((Saikia and Jain, 2007) .

Various fungal species are generally distinguished by their biochemical, physiological and metabolic capacities to metabolize or degrade various hazardous or persistent chemicals (Barrechet *al.*, 2018). Myco-remediation could be one of the ideal strategies to clean the contaminated soil and water. Myco-remediation is economically efficient and ecological sound strategy to counter the escalating crisis of aquatic and terrestrial pollution. The advantages of fungi are mainly due to robust growth, immense hyphal network, production of multipurpose extracellular enzymes, and increased surface area to volume ratio, confrontation capabilities towards complex pollutants, adaptability to fluctuating pH, temperature, and having metal-binding proteins (Kumar, 2021).

5.Increasing the uptake of carbon dioxide of organisms by genetic engineering

Plants, certain bacteria and algae continuously use photosynthesis to convert sunlight, water and atmospheric carbon dioxide (CO₂) into most of our food, furniture and fuel (Fischer *et al.*, 2016). However, this process has become more complicated over time. . Rubisco is an

enzyme that transforms CO₂ into organic molecules. In turn, slow uptake of CO limits the expansion of many plants, including crops such as rice and wheat (Long *et al.*, 2006). However, some organisms have developed ways to concentrate carbon dioxide around Rubisco, allowing the enzyme to work faster (Hennacy and Jonikas, 2020). Introducing such carbon-concentrating mechanisms to crops can increase yields by 60% while reducing the need for water and fertilizers (McGrath and Long, 2014). The simplest understood mechanism of carbon concentration is the mechanism found in bacteria, predicted in the protein structure of the so-called carboxysome, which contains Rubisco and other enzymes involved in carbon fixation. This species actively imports carbon in the form of bicarbonate (HCO₃⁻), which diffuses into the carboxysome and is converted to CO₂. The high CO₂ concentration achieved in the carboxysome maximizes Rubisco activity and thus increases total CO₂ absorption (Figure 1)



• **Figure 1. Engineering a carbon-concentration into *E. coli***

A. *Halothiobacillus neapolitanus* has carboxysome-based carbon concentration mechanisms. The cell imports carbon dioxide as bicarbonate (HCO₃⁻), which diffuses into the carboxysome. Previous work succeeded in assembling a carboxysome-like structure in the non-photosynthetic model bacterium *Escherichia coli* (Bobacciet *al.*, 2012). However, these cells required a lot of carbon dioxide to grow, indicating that additional components are needed to concentrate the carbon dioxide. Now in *eLife*, David Savage, Ron Milo and colleagues, including first author AviFlamholz, report how they require functional carbon concentration mechanisms in an organism that lacks one (Flamolzetal., 2020).

The team based at the University of California at Berkeley, the Weizmann Institute of Science and the Max Planck Institute for Molecular Plant Physiology chose the bacterium *Halothiobacillus neapolitanus* as the genetic donor for their experiment. The carboxysomes of this species are simple and well studied. In particular, Savgeabd workers previously identified 20 candidate genes likely to be necessary for the correct functioning of these structures (Desmarais *et al.*, 2019).

As a receiving species, Flamholzet *al.*,2020 chose *E. coli*, which they genetically engineered to rely on Rubisco for growth. Without the carbon concentration mechanism, this strain could not grow in ambient air, but required an additional concentration of carbon dioxide approximately 100 times greater than that in the atmosphere. Hoping to reintroduce a functional carbon concentration mechanism, the team transferred 20 candidate genes to *H. neapolitanus E. coli* strain. It is not surprising that the strain was initially unable to grow in ambient carbon dioxide, as simply adding genes is often not enough to form a complex pathway to a new organism (Antonovsky *et al.*, 2016). However, Flamholzet et al. could use genetically modified *E. coli* strain - its growth rate is proportional to Rubisco activity. This allowed the team to use a natural process experiment to identify mutations that cause the carbon concentration mechanism to malfunction, thereby increasing Rubisco activity. The experiment revealed a mutant that can grow at ambient CO₂ levels, apparently by regulating the expression level of proteins that cooperate in the carbon concentration process. This result indicated that the carboxysome-based carbon concentration mechanism of *H. neapolitanus* was successfully restored in their *E. coli* strain.

To further support this conclusion, electron microscopy was used to observe carboxysome-like structures in the engineered *E. coli* strain. To ensure the functionality of these constructs, they individually knocked out several genes known to be important for carboxysome function in the original host. These mutations had the same effect on *E. coli* than in *H. neapolitanus* - cells no longer grew at ambient CO₂ levels - confirming that the carboxysome functioned in the engineered strain as it did in the original host. These results of Flamholzet et al.2020 show that the carboxysome-based carbon concentration mechanism is transferable and

functional in another organism, providing a blueprint that paves the way for engineered plants with increased carbon absorption and thus higher yields.

6.The role of gene editing in combating climate change

Climate change is a major threat to the environment in the long term because it affects agriculture, biodiversity, human society and almost every part of our world. The first cause of climate change is the addition of greenhouse gases caused by humans to the atmosphere. Due to these human-caused emissions, the average temperature of the planet has increased by almost 1 °C since 1850 (IPCC, 2018; Nunez *et al.*, 2019).

In response to the challenges of global climate change, gene editing, also known as genome editing has emerged as a technology to help organisms adapt to global climate change or mitigate the effects of climate change. In agriculture, developing crops and livestock that can better withstand the effects of climate change.

Gene editing can be a method to insert DNA editing at precise genomic locations. These modifications can result in the deletion or destruction of one or more genes without the permanent addition of foreign DNA. Alternatively, genes from the genome of the organism or other organisms are inserted into precise locations in the genome to correct the trait. Transcription activator-like effector nucleases (TALEN), zinc finger nucleases (ZFN) and CRISPR/Cas systems have been used to achieve precise gene editing (Gajet *et al.*, 2016; Khalil, 2020).

The precision and efficiency of creating changes has greatly improved with the introduction of CRISPR/Cas systems, although there is certainly more work to be done with other gene editing techniques. Because gene editing programs make it possible to make precise changes to a plant's genome and accelerate the production of new varieties of organisms, including those that can better withstand the stress of a changing climate and those that capture and store excess atmospheric carbon dioxide.

6.1. Roles of gene editing in abiotic stress.

Future crops must be highly resistant to extreme heat and variable rainfall. In some species, such as rice, genes that confer tolerance to flooding have been identified at specific junctions, and once the genetic basis of tolerance is understood, gene editing can be used to propagate the trait more widely(*kuetal.*,2006).However, the tolerance of most plants to drought, heat and flooding is due to the influence of several genes(Wang *etal.*,2021) .Basic research in species such as sorghum and millet helps us understand how these genes work(Woldesemayat*etal.*,2017;Tan*etal.*, 2017).

6.2 Roles of gene editing in resistance to pests and pathogens

Increased abiotic stress makes plants more susceptible to biotic stresses such as pathogens caused by insects, fungi and bacteria(Coakley *etal.*,1999).In addition, warmer temperatures increase the abundance of some pathogens and change their geographic distribution. Crop varieties resistant to fungal and bacterial pathogens are usually developed by crossing resistance genes from non-elite varieties or wild relatives to commercial varieties(Scheben *etal.*,2017).

The hybrids are then crossed with a susceptible parent for several generations to eventually produce disease-resistant elite germplasm. Such breeding programs require significant time - up to ten years for some species - which is dangerously slow due to rapid climate change. In just one year, gene editing has conferred tolerance to fungal pathogens in wheat (Wan*etal.*,2014)and bacterial pathogens in rice (Li *etal.*,2012)by altering genes important for pathogen resistance and response.

6.3. Roles of gene editing for re-domestication.

Considerably few plant species provide the food that sustains humanity. During the process of domestication, changes in the expression and activity of a handful of regulatory genes led to a remarkable physical transformation of wild stem cells into the high-yielding modern crops we rely on for food(Doebley, 2006;Meyer and Purugganan,2013).

As the climate changes, we benefit from breeding new species that can withstand extreme weather conditions or thrive in marginal areas. Gene editing can greatly accelerate the domestication of these species by changing key regulatory genes to improve their productivity. One candidate for domestication is tef, a cereal crop from Ethiopia and Eritrea. Loss of seeds inhibits productivity.

6.4. Improvement of heat tolerance in cattle by genetic modification

In animals, gene editing has also been applied to mitigate abiotic stress caused by global climate change. Recombinetics Inc. subsidiary Acceligen launched an initiative to improve heat tolerance in cattle with the support of Food and Agriculture Research (FFAR) and Semex. Researchers initially focused on reproducing the SLICK phenotype identified in Senepoli cattle through gene editing. Variants of this phenotype in cattle contribute to heat tolerance (Dikmen *et al.*, 2014; Porto-Neto *et al.*, 2018). Conventionally bred cattle with SLICK genetics are more heat tolerant as shown by lower vaginal temperature, lower rectal temperature, lower respiratory rate and increased sweating, leading to increased milk production during the summer months (Dikmen *et al.*, 2014). Acceligen aims to reproduce SLICK genetics using gene editing techniques to increase heat tolerance in important cattle breeds (Bellini, 2018).

7. Roles of genetic engineering in carbon sequestration.

Plants naturally bind carbon from the atmosphere and fix it in the above and below ground parts of the plant. Unfortunately, this carbon storage is often temporary. When plants die and decompose, carbon is released back into the atmosphere (Mulligan *et al.*, 2018). Gene editing could be used to redirect captured carbon into compounds more resistant to degradation, such as suberin, a carbon-rich compound found in the roots of many plants (Schweitzer *et al.*, 2021).

In addition, root architecture could be changed to increase underground biomass and thus increase the amount of carbon stored (Ogura *et al.*, 2019). Even if there is a very small increase

in the amount of carbon stored in the soil by larger row crops, millions of tons of carbon can be washed out of the atmosphere(Mulligan*etal.*,2018).

All plants carry out photosynthesis to capture carbon dioxide from the atmosphere, but some plants have developed much more efficient photosynthetic mechanisms. So-called C4 plants, such as corn and sugarcane, are up to 50% more efficient photosynthetically than C3 plants, such as rice and wheat(Hibberd *etal.*,2008).

Attempts are being made to engineer C3 plants to carry out C4 photosynthesis(Hibberd *etal.*,2008;. Langdale,2011) .Enabling the C4 trait requires gene editing to alter the genome to produce changes in leaf architecture and metabolism. If successful, rice yields could be greatly increased, and since native C4 plants use less water, engineered C4 plants are likely to be drought tolerant

8.The role of synthetic biology in reducing atmospheric greenhouse gases

prospects and challenges ofsynthetic biology uses the concept of engineering design to engineer, modify, and even resynthesize target organisms at the molecular level, creating new organisms or transforming existing organisms (zaho, 2019). This process is usually driven by specific biological functions, including mining, designing, building and standardizing biological parts, devices, and genetic circuits, or sometimes complete chemical de novo DNA synthesis, building parts, devices, and circuits with new functions, forming .assembled, tested and optimized in networks and platform cells.

Enabling technologies related to synthetic biology generally include DNA sequencing, DNA synthesis, gene editing, genome design, synthesis and assembly, design of biological parts (including new protein design), gene circuit design, computational and biological informatics, data processing and modeling. Using these technologies, it is possible to study functional genes and make cheap and efficient computer simulations of the design, construction and metabolism of these organisms to enable the practical development and application of synthetic biology for various purposes, including climate change mitigation.

9. Conclusion

The long residence time of carbon dioxide in the atmosphere creates an urgent need to include atmospheric carbon reduction in carbon control strategies. Gene editing, genetic engineering and synthetic biology can provide powerful approaches to reduce atmospheric carbon and increase new opportunities. Possibilities include converting carbon dioxide from respiration into stable carbonate, engineering plants with a higher root-to-shoot ratio, creating plants capable of self-fertilization, using genetic engineering to trap carbon dioxide in organisms, editing genetic engineering to invent DNA modifications precisely. genomic locations, and reengineering of biological elements not normally found in nature. However, several important environmental and social challenges must be faced and resolved before such an application can be evaluated, implemented and deployed.

10. References

1. IPCC (2014) .Impacts, Adaptation, and Vulnerability. Intergovernmental Panel on Climate Change (Eds. J Houghton,). Cambridge University Press, Cambridge, UK.
2. Sallema RE, Mtui GYS (2008). Adaptation technologies and legal instruments to address climate change impacts to coastal and marine resources in Tanzania. *Afr J Environ Sci Technol* 2(9): 239-248.
3. Kumar, A., Gupta, T., Berzsenyi, S., Giangrande, A. (2015). N-cadherin negatively regulates collective Drosophila glial migration through actin cytoskeleton remodeling. *J. Cell Sci.* 128(5): 900--912.
4. FAO. (2018). The impact of disasters and crises on agriculture and food security”. food and agriculture organization of the united nations.
5. Organization for Economic Cooperation and Development (2017)
6. Falk DA, Knapp EE, Guerrant EO(2001).. *An Introduction to Restoration Genetics*. Plant Conservation Alliance, Bureau of Land Management, US Department of Interior, US Environmental Protection Authority. <http://www.ser.org/pdf/SER-restoration-genetics.pdf>. [[Google Scholar](#)]
7. Frankham, R. (2005). Stress and Adaptation in Conservation Genetics. *J. Evol. Biol* 18 (4), 750–755. doi:10.1111/j.1420-9101.2005.00885.x

8. Erickson, D. L., Hamrick, J. L., & Kochert, G. D. (2004). Ecological determinants of genetic diversity in an expanding population of the shrub *Myrica cerifera*. *Molecular Ecology*, 13(6), 1655–1664. 10.1111/j.1365-294X.2004.02139.x
9. Nosil, P., & Rundle, H. D. (2012). *Ecological speciation*. Oxford University Press.
10. Wang, I. J. and Bradburd, G. S. (2014). Isolation by environment. – *Mol. Ecol.* 23: 5649–5662.
11. Davis, M.B. and Shaw, R.G. (2001). Range Shifts and Adaptive Responses to Quaternary Climate Change. *Science*, 292, 673-679.
12. Parmesan C and Martens P. (2006). Climate change. In *Biodiversity, Health and the Environment: SCOPE/Diversitas Rapid Assessment Project*, ed. O Sala, L Meyerson, C Parmesan. Washington, DC: Island Press. In press
13. Breshears, D. D., Huxman, T. E., Adams, H. D., Zou, C. B., & Davison, J. E. (2008). Vegetation synchronously leans upslope as climate warms. *Proceedings of the National Academy of Sciences*, 105(33), 11591–11592. 10.1073/pnas.0806579105
14. Lenoir, J., Gegout, J. C., Marquet, P. A., de Ruffray, P., & Brisse, H. (2008). A significant upward shift in plant species optimum elevation during the 20th century. *Science*, 320(5884), 1768–1771. 10.1126/science.1156831
15. Thuiller, W., Lavorel, S., Araujo, M. B., Sykes, M. T., & Prentice, I. C. (2005). Climate change threats to plant diversity in Europe. *Proceedings of the National Academy of Sciences*, 102(23), 8245–8250. 10.1073/pnas.0409902102

16. Margaret, B. D., Ruth, G. S., & Julie, R. E. (2005). Evolutionary responses to changing climate. *Ecology*, 86(7), 1704–1714. 10.1890/03-0788
17. Coop, G., Witonsky, D., Di Rienzo, A., & Pritchard, J. K. (2010). Using environmental correlations to identify loci underlying local adaptation. *Genetics*, 185(4), 1411–1423. 10.1534/genetics.110.114819
18. Gonzalez-Martinez, S. C., Krutovsky, K. V., & Neale, D. B. (2006). Forest-tree population genomics and adaptive evolution. *New Phytologist*, 170(2), 227–238. 10.1111/j.1469-8137.2006.01686.x
19. Hancock, A. M., Brachi, B., Faure, N., Horton, M. W., Jarymowycz, L. B., Sperone, F. G., & Bergelson, J. (2011). Adaptation to climate across the *Arabidopsis thaliana* genome. *Science*, 334(6052), 83–86.
20. Savolainen, O., Pyhäjärvi, T., & Knürr, T. (2007). Gene flow and local adaptation in trees. *Annual Review of Ecology, Evolution, and Systematics*, 38(1), 595–619. 10.1146/annurev.ecolsys.38.091206.095646
21. Kohler, T., Giger, M., Hurni, H., Ott, C., Wiesmann, U., Wymann von Dach, S., & Maselli, D. (2010). Mountains and climate change: A global concern. *Mountain Research and Development*, 30(1), 53–55. 10.1659/MRD-JOURNAL-D-09-00086.1
22. Gottfried, M., Pauli, H., Futschik, A., Akhalkatsi, M., Barančok, P., Benito Alonso, J. L., Coldea, G., Dick, J., Erschbamer, B., Fernández Calzado, M. R., Kazakis, G., Krajčič, J., Larsson, P., Mallaun, M., Michelsen, O., Moiseev, D., Moiseev, P., Molau, U.,

- Merzouki, A., ... Grabherr, G. (2012). Continent-wide response of mountain vegetation to climate change. *Nature Climate Change*, 2, 111–115. 10.1038/nclimate1329
23. Urban MC (2015). Climate change. Accelerating extinction risk from climate change. *Science* **348**:571–573.
24. Wiens JJ (2016). Climate-related local extinctions are already widespread among plant and animal species. *PLoS Biol* **14**:e2001104.
25. Savolainen O, Lascoux M, Merilä J (2013). Ecological genomics of local adaptation. *Nat Rev Genet* **14**:807–820.
26. Bay RA, Harrigan RJ, Underwood VL, Gibbs HL, Smith TB, Ruegg K. Genomic signals of selection predict climate-driven population declines in a migratory bird. *Science*. 2018 Jan 5;359(6371):83-86. doi: 10.1126/science.aan4380. PMID: 29302012.
27. Bay RA, Rose N, Barrett R, Bernatchez L, Ghalambor CK, Lasky JR, Brem RB, Palumbi SR, Ralph P. Predicting Responses to Contemporary Environmental Change Using Evolutionary Response Architectures. *Am Nat*. 2017 May;189(5):463-473. doi: 10.1086/691233. Epub 2017 Mar 17. PMID: 28410032.
28. Shafer AB, Wolf JB, Alves PC, Bergström L, Bruford MW, Brännström I, Colling G, Dalén L, De Meester L, Ekblom R, Fawcett KD, Fior S, Hajibabaei M, Hill JA, Hoesel AR, Höglund J, Jensen EL, Krause J, Kristensen TN, Krützen M, McKay JK,

Norman AJ, Ogden R, Österling EM, Ouborg NJ, Piccolo J, Popović D, Primmer CR, Reed FA, Roumet M, Salmona J, Schenekar T, Schwartz MK, Segelbacher G, Senn H, Thaulow J, Valtonen M, Veale A, Vergeer P, Vijay N, Vilà C, Weissensteiner M, Wennerström L, Wheat CW, Zieliński P. Genomics and the challenging translation into conservation practice. *Trends EcolEvol.* 2015 Feb;30(2):78-87. doi: 10.1016/j.tree.2014.11.009. Epub 2014 Dec 17. PMID: 25534246.

29. Pacifici, Michela & Foden, Wendy & Visconti, Piero & Watson, James & Butchart, Stuart & Kovacs, Kit & Scheffers, Brett & Hole, David & Martin, Tara & Akcakaya, H. Resit & Corlett, R. & Huntley, Brian & Bickford, David & Carr, Jamie & Hoffmann, Ary & Midgley, Guy & Pearce-Kelly, Paul & Pearson, Richard & Williams, Stephen & Rondinini, Carlo. (2015). Assessing species vulnerability to climate change. *Nature Climate Change.* 5. 215-224. 10.1038/nclimate2448.

30. Hällfors MH, Liao J, Dzurisin J, Grundel R, Hyvärinen M, Towle K, Wu GC, Hellmann JJ. Addressing potential local adaptation in species distribution models: implications for conservation under climate change. *Ecol Appl.* 2016 Jun;26(4):1154-69. doi: 10.1890/15-0926. PMID: 27509755.

31. Slatyer RA, Nash MA, Hoffmann AA (2016). Scale-dependent thermal tolerance variation in Australian mountain grasshoppers. *Ecography* **39**:572–582.

32. Valladares F, Matesanz S, Guilhaumon F, Araújo MB, Balaguer L, Benito-Garzón M, Cornwell W, Gianoli E, van Kleunen M, Naya DE, Nicotra AB, Poorter H, Zavala MA. The effects of phenotypic plasticity and local adaptation on forecasts of species range shifts under climate change. *Ecol Lett*. 2014 Nov;17(11):1351-64. doi: 10.1111/ele.12348. Epub 2014 Sep 9. PMID: 25205436.
33. Ikeda DH, Max TL, Allan GJ, Lau MK, Shuster SM, Whitham TG. Genetically informed ecological niche models improve climate change predictions. *Glob Chang Biol*. 2017 Jan;23(1):164-176. doi: 10.1111/gcb.13470. Epub 2016 Sep 9. PMID: 27543682.
34. D'Amen M, Zimmermann NE, Pearman PB (2013). Conservation of phylogeographic lineages under climate change. *Glob Ecol Biogeogr* **22**:93–104.
35. Pauls SU, Nowak C, Bálint M, Pfenninger M (2013). The impact of global climate change on genetic diversity within populations and species. *Mol Ecol* **22**:925–946.
36. Razgour O, Taggart JB, Manel S, Juste J, Ibáñez C, Rebelo H, Alberdi A, Jones G, Park K. An integrated framework to identify wildlife populations under threat from climate change. *Mol Ecol Resour*. 2018 Jan;18(1):18-31. doi: 10.1111/1755-0998.12694. Epub 2017 Jul 25. PMID: 28649779; PMCID: PMC6849758.

37. Ruegg K, Bay RA, Anderson EC, Saracco JF, Harrigan RJ, Whitfield M, Paxton EH, Smith TB. Ecological genomics predicts climate vulnerability in an endangered southwestern songbird. *Ecol Lett.* 2018 Jul;21(7):1085-1096. doi: 10.1111/ele.12977. Epub 2018 May 9. PMID: 29745027.
38. Fita, A., Rodríguez-Burruezo, A., Boscaiu, M., Prohens, J., Vicente, O. (2015). Breeding and domesticating crops adapted to drought and salinity: A new paradigm for increasing food production. *Front. In Plant Sci.* 6, 978. doi: 10.3389/fpls.2015.00978
39. Bray, E. A. (1997). Plant responses to water deficit. *Trends Plant Sci.* 2 (2), 48–54. doi: 10.1016/S1360-1385(97)82562-9
40. Blum, A. (2015). Stress, strain, signaling, and adaptation –not just a matter of definition. *J. Exp. Bot.* 67 (3), 562–565. doi: 10.1093/jxb/erv497
41. Lybbert T, Sumner D (2010) Agricultural technologies for climate change mitigation and adaptation in developing countries: Policy options for innovation and technology diffusion. ICTSD-IPC Platform on Climate Change.
42. Kleter GA, Harris C, Stephenson G, Unsworth J (2008) .Comparison of herbicide regimes and the associated potential environmental effects of glyphosate-resistant crops versus what they replace in Europe. *Pest Manage Sci* 64(4): 479-488.

43. Yan Y, Yang J, Dou Y, Chen M, Ping S, et al. (2008). Nitrogen fixation island and rhizosphere competence traits in the genome of root associated *Pseudomonas stutzeri* A1501. *Proc Nat Acad Sci* 105(21): 7564-7569.
44. Hsieh TH, Lee JT, Yang PT, Chiu, LH, Charng YY, et al. (2002) .Heterogy expression of *Arabidopsis* C-repeat/dehydration response element binding factor I gene confers elevated tolerance to chilling and oxidative stresses in transgenic tomato. *Plant Physiol* 129: 1086-1094.
45. Barrows G, Sexton S, Zilberman D (2014). Agricultural Biotechnology: The Promise and Prospects of Genetically Modified Crops. *J Economic Perspectives* 28(1): 99-120.
46. Tester M, Bacic A. 2005. Abiotic stress tolerance in grasses. From model plants to crop plants. *Plant Physiol.* 137: 791–793
47. Ainsworth, E., Rogers, A. and Leakey, A.D.B. 2008. Targets for crop biotechnology in a future high-CO₂ and high-O₃ world. *Plant Physiology* 147:13-19
48. FAO, 2007
49. Howden, S.M., Soussana, J., Tubiello, F.N., Chhetri, N., Dunlop, M. and Meinke, H. (2007). Adapting agriculture to climate change. *Proceedings of the National Academy of Sciences of the United States of America (PNAS)*, 104: 19691–19696.
50. Slater R, Peskett L, Ludi E, Brown D (2007). Climate change, agricultural policy and poverty reduction: How much do we know? *Natural Resource Perspectives* 109. Overseas Development Institute, London.

51. Abatenh, Endeshaw & Gizaw, Birhanu & Tsegaye, Zerihun & Genene, Tefera. (2018). Microbial Function on Climate Change - A Review. Environment Pollution and Climate Change. 02. 10.4172/2573-458X.1000147.

UNDER PEER REVIEW

52. Delgado-Baquerizo, M. et al. Multiple elements of soil biodiversity drive ecosystem functions across biomes. *Nat. Ecol. Evol.* **4**, 210–220 (2020).
53. Du, Z., Li, H., and Gu, T. (2007). A state of the art review on microbial fuel cells: a promising technology for wastewater treatment and bioenergy. *Biotechnol. Adv.* **25**, 464–482. doi: 10.1016/j.biotechadv.2007.05.004
54. Mohammadi, K., and Sohrabi, Y. (2012). Bacterial biofertilizers for sustainable crop production: a review. *J. Agric. Biol. Sci.* **7**, 307–316. doi: 10.1186/1475-2859-13-66
55. Singh, R. B. (2000). “Biotechnology, biodiversity and sustainable agriculture—a contradiction?” in *Regional Conference in Agricultural Biotechnology Proceedings: Biotechnology Research and Policy -Needs and Priorities in the Context of Southeast Asia's Agricultural Activities*, ed E. M. T. Mendoza (Bangkok: SEARCA (SEAMEO)/FAO /APSA).
56. Rajneesh, Singh, S. P., Pathak, J., and Sinha, R. P. (2017). Cyanobacterial factories for the production of green energy and value-added products: an integrated approach for economic viability. *Renew. Sustain. Energy Rev.* **69**, 578–595. doi: 10.1016/j.rser.2016.11.110
57. Lewis, N. S., and Nocera, D. G. (2006). Powering the planet: chemical challenges in solar energy utilization. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 15729–15735. doi: 10.1073/pnas.0603395103
58. Melis, A. (2009). Solar energy conversion efficiencies in photosynthesis: minimizing the chlorophyll antennae to maximize efficiency. *Plant Sci.* **177**, 272–280. doi: 10.1016/j.plantsci.2009.06.005
59. Rittmann, B. E. (2008). Opportunities for renewable bioenergy using microorganisms. *Biotechnol. Bioeng.* **100**, 203–212. doi: 10.1002/bit.21875
60. Kaneko, T., Sato, S., Kotani, H., Tanaka, A., Asamizu, E., Nakamura, Y., et al. (1996). Sequence analysis of the genome of the unicellular cyanobacterium *Synechocystis* sp. 6803. II. Sequence determination of the entire genome and assignment of potential protein-coding regions. *Res.* **3**, 109–136.
61. Ducat, D. C., Way, J. C., and Silver, P. A. (2011). Engineering cyanobacteria to generate high-value products. *Trends biotechnol.* **29**, 95–103. doi: 10.1016/j.tibtech.2010.12.003

- 62..Mishra, A.K.; Kaushik, M.S.; Tiwari, D.N. Nitrogenase and Hydrogenase: Enzymes for Nitrogen Fixation and Hydrogen Production in Cyanobacteria. In Cyanobacteria; Mishra, A.K., Tiwari, D.N., Rai, A.N., Eds.; Academic Press: Cambridge, MA, USA, 2019; pp. 173–191. ISBN 978-0-12-814667-5.
- 63.Schipper, K.; Al Muraikhi, M.; Alghasal, G.S.H.S.; Saadaoui, I.; Bounnit, T.; Rasheed, R.; Dalgamouni, T.; Al Jabri, H.M.S.J.; Wijffels, R.H.; Barbosa, M.J. Potential of novel desert microalgae and cyanobacteria for commercial applications and CO₂ sequestration. *J. Appl. Phycol.* **2019**, 31, 2231–2243.
- 64.Adams, D.G.; Bergman, B.; Nierzwicki-Bauer, S.A.; Duggan, P.S.; Rai, A.N.; Schüßler, A. Cyanobacterial-Plant Symbioses. In *The Prokaryotes*; Rosenberg, E., DeLong, E.F., Lory, S., Stackebrandt, E., Thompson, F., Eds.; Springer: Berlin/Heidelberg, Germany, 2013; pp. 359–400. ISBN 978-3-642-30194-0.
- 65.Mohan, A.; Kumar, B. Plant Growth Promoting Activities of Cyanobacteria Growing In Rhizosphere of Agriculturally Fertile Soil. *J. Biotechnol. Biochem.* **2019**, 5, 28–36.
- 66..Rady, M.M.; Taha, S.S.; Kusvuran, S. Integrative application of cyanobacteria and antioxidants improves common bean performance under saline conditions. *Sci. Hortic. Amst.* **2018**, 233, 61–69.
- 67..Kirsch, F.; Klähn, S.; Hagemann, M. Salt-Regulated Accumulation of the Compatible Solutes Sucrose and Glucosylglycerol in Cyanobacteria and Its Biotechnological Potential. *Front. Microbiol.* **2019**, 10.
- 68.Qiang, H., and Richmond, A. (1994). *Optimizing the population density in Isochrysisgalbana grown outdoors in a glass column photobioreactor.* *J. Appl. Phycol.* 6, 391–396.
- 69.Al-Haj , L. , Lui , Y.T. , Abed , R.M.M. , Tire , M.A. and Purton , S. . (2016). Cyanobacteria as a platform for industrial biotechnology: progress and future perspectives. *Life* 6:42. doi: 10.3390/life6040042

70. Sarma, M.K., Kaushik, S. and Goswami, P. (2016). Cyanobacteria: metabolic power to harvest solar energy for bioelectricity and biofuel production. *Biomass. Bioenergy.* 90, 187–201. doi: 10.1016/j.biombioe.2016.03.043.
71. Lewis, N.S. and Nocera, DG (2006). Powering the Planet: The Chemical Challenges of Harnessing Solar Energy (The Chemical Challenges of Utilizing Solar Energy). *Proc. Natl. Acad. To know US* 103, 15729-15735. doi: 10.1073/pnas.0603395103
72. Melis, A. (2009). Solar energy conversion efficiency in photosynthesis: minimizing chlorophyll antennas to maximize efficiency. *Plant Sci.* 177, 272–280. doi: 10.1016/j.plantsci.2009.06.005
73. Rittmann, B.E. (2008). Possibilities of renewable bioenergy with microorganisms. *Biotechnology. Bioeng.* 100, 203–212. doi: 10.1002/bit.21875
74. Kaneko, T., Sato, S., Kotani, H., Tanaka, A., Asamizu, E., Nakamura, Y. et al. (1996). The unicellular cyanobacterium *Synechocystis* sp. 6803. II. Whole genome sequencing and identification of potential protein coding regions. *Res.* 3, 109–136.
75. Ducat, D. C., Way, J. C. kaj Silver, P. A. (2011). Development of blue-green algae for the production of valuable products. *Trends in Biotechnology.* 29, 95–103. doi: 10.1016/j.tibtech.2010.12.003
76. Oliver, N. J., Rabinovitch-Deere, C. A., Carroll, A. L., Nozzi, N. E., Case, A. E. kaj Atsumi, S. (2016). Metabolic technology of blue-green algae for the production of biofuels and chemicals. *Curr. I learned Chem. Biol.* 35, 43–50. doi: 10.1016/j.cbpa.2016.08.023
77. Savakis, P. and Hellingwerf, K. J. (2014). Engineered cyanobacteria for direct biofuel production from CO₂. *Curr. I learned Biotechnology.* 33C, 8-14. doi: 10.1016/j.copbio.2014.09.007

78. Gao, X., Sun, T., Pei, G., Chen, L., and Zhang, W. (2016). Blue-green algae substrate design for enhanced production of biofuels and chemicals. *Application Microbiol. Biotechnology*. 100, 3401–3413. doi: 10.1007/s00253-016-7374-2
79. Gudmundsson, S. and Nogales, J. (2015). Cyanobacteria as photosynthetic biocatalysts: a systems biology perspective. *Mol. Biosist.* 11, 60–70. doi: 10.1039/C4MB00335G
80. Wang, B., Wang, J., Zhang, W. kaj Meldrum, D. R. (2012). Applications of synthetic biology in cyanobacteria and algae. *Before. Microbiol.* 3:44 in the morning. doi: 10.3389/fmicb.2012.00344
81. Berla, B. M., Saha, R., Immethun, C. M., Maranas, C. D., Moon, T. S. and Pakrasi, H. (2006). B. (2013). Cyanobacterial synthetic biology: unique challenges and opportunities. *Before. Microbiol.* 4:246. doi: 10.3389/fmicb.2013.00246
82. Desai, S.H. kaj Atsumi, S. (2013). Photosynthetic Approaches al Chemical Biotechnology. *Curr. Arvamus. Bioteknologio.* 24, 1031-1036. doi: 10.1016/j.copbio.2013.03.015
83. Mahanty, T., Bhattacharjee, S., Goswami, M., Bhattacharyya, P., Das, B., Ghosh, A. et al. (2017). Biofertilizers: a potential approach to sustainable agricultural development. *The neighborhood. To know Contaminate Race. Rev.* 24, 3315–3335.
84. Hegde, D. M., Dwiwedi, B. S. and Babu, S. (1999). N.S. (1999). Biofertilizers for cereal production in India: a review. *Indian J. Agriculture. Sci.* 69, 73–8
85. Vaishampayan, A., Sinha, R. P., Hader, D.-P. K., Bhan, U. et al. (2001). Biostercous cyanobacteria in recultivated. *Ship. Rev. Fr.* 67, 453–516. doi: 10.1007/BF02857893
86. Metting, B. (1988). "Mikroalgot en agrikultur", in *Micro-Algal Biotechnology*, ed. M.A. Borowitzka and L. J. Borowitzka (Cambridge: Cambridge University Press), 288-304.
87. Aiyer, R. S., Sulahudean, S. and Venkataraman, G. S. (1972). A long-term propagation field experiment with high-yielding rice varieties. *Indian, J. Agric. Theo* 42, 380-383.

88. Watanabe, I., Spinas, C.R., Berja, N.S. V. (1977). Uzoof Azolla-Anabaena complex kielnitrogensterko by curl. IRRI Res. Paper Ser. 11, 1-15.
89. Prasanna, R., Sharma, E., Sharma, P., Kumar, A., Kumar, R., Gupta, V., et al. (2013). Soil fertility and potential for inoculated cyanobacteria in rice crops grown under non-flooded conditions. Paddy Water Environment. 11, 175–183. doi: 10.1007/s10333-011-0302-2
90. Subramanian, G. and Sundaram, S.S. (1986). Evolution of ammonia from the nitrogenfixing cyanobacterium Anabaena. FEMS Microbiol. Lett. 37, 151–154. doi: 10.1111/j.1574-6968.1986.tb01784.x
91. Rai, A. K. kaj Sharma, N. K. (2006). Phosphate metabolome in the cyanobacterium Anabaena doliolum sub saline. Curr. Microbiology. 52, 6–12. doi: 10.1007/s00284-005-0043-9
92. Sarma, M.K., Kaushik, S. and Goswami, P. (2016). Cyanobacteria: metabolic power to harvest solar energy for bioelectricity and biofuel production. BiomasoBioenergio. 90, 187–201. doi: 10.1016/j.biombioe.2016.03.043
93. Venkataraman, G.S. (1993). "Bluverday algae (cyanobacteria)," in Biological Nitrogen Fixation, ed. S. N. Tata, A.M. Wadhvani ja M. S. Mehdi (Nov-Delhio: Indian Council Agric. Res.), 45-76.
94. Venkataraman, G. S. (1972). Algal biofertilizer and rice cultivation. New Delhi: Printers and Publishers of Today and Tomorrow.
95. Roger, P. A. and Kulasooriya, S.A. (1980). Blue green algae and rice. Manila: International Rice Research Institute. Kannaiyan, S., Aruna, S.J., Kumari, S.M.P. and Hall, D.O. (1997). Immobilized cyanobacteria as biofertilizer for rice cultivation. J. Appl. Phycol. 9, 167–174. doi: 10.1023/A:1007962025662
96. Dubey, A. K. and Rai, A. K. (1995). Use of algal biofertilizers (Aulosirafertilissimatenuis and Anabaena doliolumBhardwaja) for long-term rice cultivation in North India. Israel J. Plant. To know 43, 41–51. doi: 10.1080/07929978.1995.10676589

97. Pereira, I., Rodrigo, O., Barrientos, L., Moya, M., Reyes, G. and Kramm, V. (2009). Development of a biofertilizer based on filamentous nitrogen-fixing cyanobacteria for rice cultivation in Chile. *J. Apliko de Phycol.* 21, 135–144. doi: 10.1007/s10811-008-9342-4
98. Spiller, H. and Gunasekaran, M. (1990). An ammonia-secreting mutant strain of the cyanobacterium *Anabaena variabilis* supports wheat growth. *Application Microbiol. Biotechnology.* 33, 477–480. doi: 10.1007/BF00176670

99. Obreht, Z., Kerby, N.W., Gantar, M., & Rowell, P. (1993). Effects of root-associated N₂-fixing cyanobacteria on growth and nitrogen content of wheat (*Triticum vulgare* L.) seedlings. *Biol. Fekunda. Grundoj* 15, 68–72. doi: 10.1007/BF00336292
100. Karthikeyan, N., Prasanna, R., Sood, A., Jaiswal, P., Nayak, S. and Kaushik, B.D. (2009). Physiological characterization and electron microscopic studies of wheat rhizosphere-associated cyanobacteria. *Leaf Microbiol.* 54, 43–51. doi: 10.1007/s12223-009-0007-8
101. Swarnalakshmi, K., Prasanna, R., Kumar, A., Pattnaik, S., Chakravarty, K., Shivay, Y.S. et al. (2013). Evaluation of the effects of novel cyanobacterial biofilmed biofertilizers on wheat soil fertility and plant nutrition. *Euros J. Soil Biol.* 55, 107–116. doi: 10.1016/j.ejsobi.2012.12.008
102. Bose, P. and Nagpal, USA (1971). Tricalcium phosphate solution of cyanobacteria. *Curr. Sci.* 40, 165–167. doi: 10.1007/s00284-005-0043-9
- Rai, A.K. and Sharma, N.K. K. (2006). Phosphate metabolome in the cyanobacterium *Anabaena doliolum* sub saline. *Curr. Microbiology.* 52, 6–12. doi: 10.1007/s00284-005-0043-9
103. Dorich, RA, Nelson, DW and Sommers, LE. (1985). Estimation of algal available phosphorus in suspended sediments using chemical extraction. *J. Environment. Quality* 14, 400–405. doi: 10.2134/jeq1985.00472425001400030018x
104. Cameron, H. J. and Julian, G. R. (1988). Utilization of hydroxyapatite by cyanobacteria as a sole source of phosphate and calcium. *Plant Earth* 109, 123–124. doi: 10.1007/BF02197589
105. Abdel-Raouf, N., Al-Homaidan, A.A. and Ibrahim, I.B. (2012). The importance of algae in agriculture. *African J. Biotechnol.* 11, 11648–11658. doi: 10.5897/AJB11.3983
106. Rodriguez, AA, Stella, AM, Storni, MM, Zulpa, G and Zaccaro, M. C. (2006). Effects of cyanobacterial extracellular products and gibberellic acid on salinity tolerance of *Oryza sativa* L. *Saline Syst.* 2:7. doi: 10.1186/1746-1448-2-7

107. Rastogi, R.P. and Sinha, R.P. (2009). Biotechnological and industrial importance of cyanobacterial secondary metabolites. *Biotechnology. Adv.* 27, 521–539. doi: 10.1016/j.biotechadv.2009.04.009
108. Brouers, M., De Jong, H., Shi, D.J., Rao, K. K. and Hall, D.O. (1987). Continuous production of ammonia by immobilized cyanobacteria. *Program Photosynth. Res.* 2, 645–647. doi: 10.1007/978-94-009-3535-8_153
109. Shi, D.-J. and Hall, D.O. (1988). The Azolla-Anabaena association: a historical perspective, symbiosis and energy metabolism. *Bot. Rev.* 54, 253–386. doi: 10.1007/BF02858416
110. Antonovsky N, Gleizer S, Noor E, Zohar Y, Herz E, Barenholz U, Zelbuch L, Amram S, Wides A, Tepper N, Davidi D, Bar-On Y, Bareia T, Wernick D, GShani I, Malitsky J, Ona G, Bar-Even A, Milo R (2016) Sugar synthesis from CO₂ in *Escherichia coli* *Cell* **166**:115–125.
111. Bonacci
 WTeng PKAfonso BNiederholtmeyer HGrob PSilver PASavage DF (2012) Modularity of a carbon-fixing protein organelle *PNAS* **109**:478–483. <https://doi.org/10.1073/pnas.1108557109>
112. Desmarais
 JJFlamholz AIBlikstad CDugan EJLaughlin TGOltrogge LMChen AWWetmore KDiamond SWang JYSavage DF (2019) DABs are inorganic carbon pumps found throughout prokaryotic phyla *Nature Microbiology* **4**:2204–2215.
<https://doi.org/10.1038/s41564-019-0520-8>
113. Saikia, S. P., and Vanita Jain. “Biological Nitrogen Fixation with Non-Legumes: An Achievable Target or a Dogma?” *Current Science* 92, no. 3 (2007): 317–22.
<http://www.jstor.org/stable/24096726>.

114. Fischer, W.W., Hemp, J., Johnson, J.E (2016) Evolution of oxygenic photosynthesis *Annual Review of Earth and Planetary Sciences* **44**:647–683.

<https://doi.org/10.1146/annurev-earth-060313-05481>

115. Flamholz, A.I., Dugan, E., Blikstad, C., Gleizer, S., Ben-Nissan, R., Amram, S., Antonovsky, N., Ravishankar, S., Noor, E., Bar-Even, A., Milo, R., Savage, D.F. (2020). Functional reconstitution of a bacterial CO₂ concentrating mechanism in *Escherichia coli* *eLife* **9**:e59882. <https://doi.org/10.7554/eLife.59882>

116. Hennacy, J.H., Jonikas, M.C. (2020). Prospects for engineering biophysical CO₂ concentrating mechanisms into land plants to enhance yields *Annual Review of Plant Biology* **71**:461–485.

<https://doi.org/10.1146/annurev-arplant-081519-040100>

117. Long, S.P., Zhu, X., Naidu, S., Ort, D.R. (2006). Can improvement in photosynthesis increase crop yields? *Plant, Cell and Environment* **29**:315–330. <https://doi.org/10.1111/j.1365-3040.2005.01493.x>

118. McGrath, J., Long, S.P. (2014). Can the cyanobacterial carbon-concentrating mechanism increase photosynthesis in crop species? A theoretical analysis *Plant Physiology* **164**:2247–2261.

<https://doi.org/10.1104/pp.113.232611>

- 119..IPCC (2018). “Summary for policymakers,” in *Global Warming of 1.5°C. An IPCC Special Report on the Impacts of Global Warming of 1.5°C Above Pre-industrial Levels and Related Global Greenhouse Gas Emission Pathways, in the Context of Strengthening the Global Response to the Threat of Climate Change, Sustainable Development, and Efforts to Eradicate Poverty*, eds V. Masson-Delmotte, P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P. R. Shukla, et al. (Geneva: World Meteorological Organization),
120. Nunez, S., Arets, E., Alkemade, R., Verwer, C., and Leemans, R. (2019). Assessing the impacts of climate change on biodiversity: is below 2 °C enough? *Clim. Change* 154, 351–365. doi: 10.1007/s10584-019-02420-x
- 121.Gaj, T., Sirk, S. J., Shui, S., and Liu, J. (2016). Genome-editing technologies: principles and applications. *Cold Spring Harb. Perspect. Biol.* 8:a023754. doi: 10.110
122. Xu, K., Xu, X., Fukao, T., Canlas, P., Maghirang-Rodriguez, R., et al., 2006. Sub1A is an ethylene-response-factor-like gene that confers submergence tolerance to rice. *Nature* 442, 705–708.
- <https://doi.org/10.1038/nature04920>
123. Wang, J., Li, C., Li, L., Reynolds, M., Mao, X., Jing, R., 2021. Exploitation of Drought Tolerance-Related Genes for Crop Improvement. *International Journal of Molecular Sciences* 22.
- <https://doi.org/10.3390/ijms221910265>

124. Woldesemayat, A.A., Van Heusden, P., Ndimba, B.K., Christoffels, A., 2017. An integrated and comparative approach towards identification, characterization and functional annotation of candidate genes for drought tolerance in sorghum (*Sorghum bicolor* (L.) Moench). *BMC Genetics* 18, 119. <https://doi.org/10.1186/s12863-017-0584-5>
125. Tang, S., Li, L., Wang, Y., Chen, Q., Zhang, W., et al., X., 2017. Genotype-specific physiological and transcriptomic responses to drought stress in *Setaria italica* (an emerging model for Panicoideae grasses). *Scientific Reports* 7, 10009. <https://doi.org/10.1038/s41598-017-08854-61/cshperspect.a023754>
126. Coakley, S.M., Scherm, H., Chakraborty, S., 1999. Climate change and plant disease management. *Annu. Rev. Phytopathol.* 37, 399–426. <https://doi.org/10.1146/annurev.phyto.37.1.399>
127. Scheben, A., Wolter, F., Batley, J., Puchta, H., Edwards, D., 2017. Towards CRISPR/Cas crops – bringing together genomics and genome editing. *New Phytologist* 216, 682–698. <https://doi.org/10.1111/nph.14702> Potential impacts of gene editing on climate adaptation and mitigation 8
128. Wang, Y., Cheng, X., Shan, Q., Zhang, Y., Liu, J., Gao, C., Qiu, J.-L., 2014. Simultaneous editing of three homoeoalleles in hexaploid bread wheat confers heritable resistance to powdery mildew. *Nature Biotechnology* 32, 947–951.

<https://doi.org/10.1038/nbt.2969>

129. Li, T., Liu, B., Spalding, M.H., Weeks, D.P., Yang, B., 2012. High-efficiency TALEN-based gene editing produces disease-resistant rice. *Nature Biotechnology* 30, 390–392. <https://doi.org/10.1038/nbt.2199>
130. Mulligan, J., Ellison, G., Gasper, R., Rudee, A., 2018. Carbon Removal in Forests and Farms in the United States. Working Paper. Washington, DC: World Resources Institute. <https://www.wri.org/publication/land-carbon-removal-us>
131. Khalil, A. M. (2020). The genome editing revolution: review. *J. Genet. Eng. Biotechnol.* 18:68. doi: 10.1186/s43141-020-00078-y
132. Schweitzer, H., Aalto, N.J., Busch, W., Chat Chan, D.T., Chiesa, M., et al., 2021. Innovating carbon-capture biotechnologies through ecosystem-inspired solutions. *One Earth* 4, 49–59. <https://doi.org/10.1016/j.oneear.2020.12.006>
133. Ogura, T., Goeschl, C., Filiault, D., Mirea, M., Slovak, R., et al., 2019. Root System Depth in Arabidopsis Is Shaped by EXOCYST70A3 via the Dynamic Modulation of Auxin Transport. *Cell* 178, 400-412.e16. <https://doi.org/10.1016/j.cell.2019.06.021>
134. Mulligan, J., Ellison, G., Gasper, R., Rudee, A., 2018. Carbon Removal in Forests and Farms in the United States. Working Paper. Washington, DC: World Resources Institute. <https://www.wri.org/publication/land-carbon-removal-usa>

135. Hibberd, J.M., Sheehy, J.E., Langdale, J.A., 2008. Using C4 photosynthesis to increase the yield of rice—rationale and feasibility. *Current Opinion in Plant Biology* 11, 228–231. <https://doi.org/10.1016/j.pbi.2007.11.002>
136. Langdale, J.A., 2011. C4 Cycles: Past, Present, and Future Research on C4 Photosynthesis. *The Plant Cell* 23, 3879–3892. <https://doi.org/10.1105/tpc.111.092098>
137. Meyer, R.S., Purugganan, M.D., 2013. Evolution of crop species: genetics of domestication and diversification. *Nature Reviews Genetics* 14, 840–852. <https://doi.org/10.1038/nrg3605>
138. Doebley J.F., Gaut B.S., Smith B.D., 2006. The molecular genetics of crop domestication. *Cell*. 127, 1309-1321. <https://doi.org/10.1016/j.cell.2006.12.006>
139. Shi, J., Gao, H., Wang, H., Lafitte, H. R., Archibald, R. L., Yang, M., et al. (2017). ARGOS8 Variants generated by CRISPR-Cas9 improve maize grain yield under field drought stress conditions. *Plant Biotechnol. J.* 15, 207–216. [doi: 10.1111/pbi.1260](https://doi.org/10.1111/pbi.1260)
140. Jainendra Pathak, Rajneesh, Pankaj K. Maurya, Shailendra P. Singh, Donat-P. Häder and Rajeshwar P. Sinha. *Cyanobacterial Farming for Environment Friendly Sustainable Agriculture Practices: Innovations and Perspectives. Front. Environ. Sci., 28 February 2018 Sec. Environmental Toxicology. Volume 6 - 2018*

UNDER PEER REVIEW