

# Antifungal Activities of Non-volatile Compounds of *Trichoderma harzianum* And *Trichoderma longibrachiatum* against Rot Fungi Associated with *Ananas comosus* Linn. (Pineapple)

## ABSTRACT

**Aim:** This study investigated the inhibitory effect of non-volatile compounds of *Trichoderma harzianum* and *Trichoderma longibrachiatum* against fungal pathogens associated with spoiled pineapple.

**Study design:** This experiment was laid out in a randomized complete block design.

**Place and duration of study:** Department of Biology, The Polytechnic, Ibadan, Nigeria, between June 2023 and September 2023

**Methodology:** Pineapples exhibiting symptoms of rot were purchased from Oje market within Ibadan, Nigeria. Isolation of rotten fungi was done by direct inoculating samples showing symptoms of rot on sterilized Potato Dextrose agar (PDA). Fungal isolates were identified using molecular techniques. The role of the isolates obtained in a formation of symptoms was determined by the pathogenicity test. Evaluation of non-volatile antifungal compounds of *T. harzianum* and *T. longibrachiatum* against the fungal isolates were carried out *in-vitro* at 15%, 20% and 25% concentration. All the data obtained were subjected to statistical analysis.

**Results:** The mycoflora isolated was identified as *Penicillium funiculosum*, *Aspergillus niger* and *Meyerozyma carribica*. The isolates reproduced the same rot symptoms, while *A. niger* and *P. funiculosum* had a high level of virulence with increasing incubation times. The non-volatile compounds of both biocontrol agents exert fungistatic effect on *A. niger*, *M. carribica* and *P. funiculosum* at 15%, 20% and 25% concentrations.

**Conclusion:** Therefore, non-volatile compounds of both biological agents inhibited growth of pathogenic fungi of pineapple *in vitro*. Thus, it could serve as a better alternative to synthetic chemicals.

**Key words :** Fungistatic effect, Mycoflora, Non-volatile compounds, Pathogenicity test.

## 1. INTRODUCTION

Pineapple (*Ananas comosus* L.) is a fruit with high nutritional contents, commonly grown in the tropics. It is among the ten most cultivated fruits globally and it contributes to more than 20% of the world production of tropical fruits (Liu *et al.*, 2017; Chillet *et al.*, 2020). Nigeria is ranked seventh among the world's leading producers and the leading producer in Africa (FAO, 2019). The pineapple value chain from farm to consumption ensures food security, and creates employment while improving rural livelihoods. This identifies pineapple as a crop of economic value (Quijandria *et al.*, 1997; All Africa, 2011).

However, postharvest fruit loss is estimated to account for more than 50% of the farm produce losses including pineapple in Nigeria (Moss, 2002; Zhang *et al.*, 2017; Zakawa *et al.*, 2019). Rot is a common problem of fleshy fruits such as pineapple. It is caused by phytopathogens such as fungi which enter the host tissue through mechanical wound and induce spoilage. The intrinsic low pH values, high sugar contents and the presence of other nutrients in pineapple are among the factors that predispose it to fungal deterioration (Onuorah *et al.*, 2013). In order to limit spoilage of the produce during storage, pineapple is treated with synthetic fungicides (Chillet *et al.*, 2020) which pose detrimental effects on human health; this necessitates the need to search for an alternative means.

In addition to being eco-friendly, biological control agents such as *Trichoderma* sp. are considered a good alternative to hazardous synthetic chemicals because they are host-specific, non-resistant to pathogens and effective in the short, medium and long terms. Moreover, they do not have a detrimental effect on humans or the environment (Carmon-Hernandez *et al.*, 2019). *Trichoderma* sp. shows diverse antagonistic mechanisms towards fungal phytopathogens. These could include competition, antibiosis and

mycoparasitism which involves coiling of hyphae and secretion of cell wall degrading hydrolytic enzymes (Bhardwaj and Kumar, 2017). Therefore, this study aimed to investigate effects of non-volatile compound of *Trichoderma* against fungal pathogens in pineapple.

## 2. MATERIALS AND METHODS

### 2.1 Sample collection

Samples of pineapples with obvious lesions were purchased and transported in sterile plastic bags to the laboratory for further analysis.

### 2.2 Isolation of fungi

The modified method of Odebode *et al.* (2006) was used for isolation of fungi. The tissues of pineapple were sectioned from their healthy portions to diseased portions and cut into small fragments, soaked in 5% sodium hypochlorite solution for 5 minutes for surface sterilization and then rinsed thrice in distilled water. The infected tissues were aseptically blotted using sterile filter paper. These were aseptically inoculated onto freshly prepared plates of acidified potato dextrose agar (PDA). The plates were incubated at  $30 \pm 2^\circ\text{C}$  for 3-5 days. Mixed cultures were subcultured by repeated culturing on freshly prepared agar until a pure culture was obtained.

### 2.3 Molecular characterization of Isolates

#### 2.3.1 DNA extraction

The fungal isolates obtained and stocked in agar slants were incubated at  $25^\circ\text{C}$  for 7 days. Lysis buffer (400 mM Tris-HCL (pH 8), 60 mM EDTA- pH 8.0, 150 mM NaCl and 1% sodium dodecyl sulphate) was put in an Eppendorf tube and a small quantity of mycelium was taken from each of the slants and introduced into the tube. The tube was vortexed to disrupt the mycelia and maintained at room temperature for 10 minutes. Potassium acetate (150  $\mu\text{l}$ ) was added into the Eppendorf tube and this was followed by brief vortexing. The tube and contents were finally centrifuged at  $13,000 \times g$  for 1 min. The supernatant obtained was decanted, centrifuged again and decanted into another Eppendorf tube, to which an equal volume of isopropyl alcohol was added. After mixing by inversion, the contents of the tube were centrifuged, the supernatant was discarded and the residual pellets obtained were washed with ethanol (300  $\mu\text{l}$ ; 70%). This was centrifuged at 10,000 rpm for 1 min followed by decantation of the supernatant. The DNA pellet was air dried and dissolved in  $1\times$ Tris- EDTA (50 $\mu\text{l}$ ). Purified DNA (1  $\mu\text{l}$ ) was used in 24  $\mu\text{l}$  of PCR mixture (Liu *et al.* 2000)

#### 2.3.2 PCR amplification

The extracted DNA was amplified by the Polymerase Chain Reaction. Each PCR mixture contained 10  $\mu\text{l}$  of Red TAQ READY MIX, 0.5  $\mu\text{l}$  of each primer pair, 8  $\mu\text{l}$  of analytical grade sterile water and 5  $\mu\text{l}$  of genomic DNA in a total volume of 24  $\mu\text{l}$ . The thermocycling program used was an initial denaturation ( $94^\circ\text{C}$  for 5 minutes), 30 cycles of denaturation ( $94^\circ\text{C}$  for 1 min), annealing ( $60^\circ\text{C}$  for 1 minute) and elongation ( $72^\circ\text{C}$  for 1 minute), then a stabilization ( $72^\circ\text{C}$  for 5 minutes) (Michaelsen *et al.*, 2006). Amplified genetic materials were electrophoresed on 2% agarose gel in Tris acetate- EDTA buffer and the gel was stained with ethidium bromide before observation in UV detector (Okoro *et al.*, 2009).

#### 2.3.3 DNA sequencing

The sequence was determined at the Laboratory of Mycology, Department of Environmental Science, University of Pavia, Pavia, Italy. The identities of the isolates were determined against known sequences in the GenBank using BLAST (Basic Local Alignment Search Tool).

### 2.4 Pathogenicity Test

The method of Chukwura *et al.* (2010) was adopted. Mature and fresh pineapple fruits were washed with distilled water, surface sterilized with 75% ethanol and blotted dry with filter paper. A sterile 4mm cork borer was used to bore hole in each fruit, where the isolated fungi were inoculated and the cores were aseptically replaced to cover the hole in the fruits. The holes were sealed with petroleum jelly. All experiments including

the control were placed in clean polythene bag and incubated in a humid environment at ambient temperature for ten days. All experiments were carried out in triplicate. The fruits were examined through the inoculated site for lesion development. Infected parts were aseptically transferred onto sterilized PDA to identify microorganisms responsible for the infections.

## 2.5 Assessment of antifungal properties of non-volatile metabolites of *Trichoderma* sp.

This was carried out according to the procedure described by Dennis and Webster (1971a). *Trichoderma* was cultivated in 250 mL conical flasks containing 100 mL of potato dextrose broth (PDB) and incubated for 10 days with periodic shaking. The set up was filtered with Whatman no. 42 filter paper. The culture was further centrifuged at 6000 rpm for 10 minutes and sterilized using a 0.4 µm pore Millipore membrane filter. A final concentration of 15%, 20% and 25% (v/v) was obtained by mixing the culture filtrate with requisite amount of molten PDA. The fortified medium was poured into Petri dishes. A 5mm diameter mycelial plug of test pathogen was inoculated on the plates and incubated for 7 days at 30 ± 2°C. A control experiment was set up without inoculation of the antagonist. The radial mycelial growth of treatment and control was measured (to the nearest milliliter) for 7 days.

## 2.6 Statistical analysis

The data obtained were analyzed using SPSS software and subjected to analysis of variance. The means were differentiated at 5% confidence interval using Duncan's Multiple Range Test.

## 3. RESULTS

### 3.1 Isolation of fungi

Three isolates obtained from the sample were identified as *Aspergillus niger*, *Penicillium funiculosum* and *Meyerozyma carribica*.

### 3.2 Phylogenetic Relationships between the Isolated *Aspergillus niger*, *Penicillium funiculosum* and *Meyerozyma carribica*. and Otherstrains Already Documented in the National Center for Biotechnology Information (NCBI), USA.

It was observed that these strains showed high similarities with other fungus strains that were already documented on NCBI. It was observed that Strain tok1 is very similar to other five strains. AJ876876 was the most similar organism to strain1. Strain1 share a common ancestral lineage with AJ876876 and KU877217. These fungi also share common ancestor with other three fungi [MH553376, MF422165, MF422153]. It was discovered that strain tok 2 is mostly similar to *P.funiculosum* JX469422. However, they both share common ancestral lineage with other fungal strains. *Talaromyces stollii* JX965246 and *Talaromyces funiculosus* KJ728703. All these strains uniquely originated from the same ancestor which is similar to *Talaromyces funiculosus* HG964290. The strain tok 3 was identified as *Meyerozyma caribbica* and phylogeny reveals that this strain is very similar to *M. Caribbica* LC422336 whom they both share similar attributes with *M. Caribbica* LC422337 which in turn serves as an ancestor to *M. Caribbica* LC422338. Interestingly, these four strains share a strikingly similar ancestral lineage (0.92 pasimonial similarities) with other three *candida* strains viz: *Candida smithsoni* KY495770 which is most similar to *C. smithsonii* MG385075 but farther to *M. caribbica* MG976725.




Fig. 1. Phylogenetic relationship between the isolated *Aspergillus niger* strain tok 1 and similar fungal strains already documented in National Center for Biotechnology Information (NCBI), USA




Fig. 2. Phylogenetic relationship between the isolated *Penicillium funiculosum* strain tok 2 and similar fungal strains already documented in National Center for Biotechnology Information (NCBI), USA.




Fig. 3. Phylogenetic relationship between the isolated *Meyerozyma caribbica* ostrain FM and similar fungal strains already documented in National Center for Biotechnology Information (NCBI), USA.

### 3.3 Pathogenicity of pathogenic fungi causing rots in pineapple

*Aspergillus niger*, *Meyerozyma carribica* and *Penicillium funiculosum* isolated from rotten pineapples were able to incite similar rot symptoms on healthy pineapple fruits. It was observed that these pathogens caused varying levels of rot symptoms based on their varying virulence (Table 1). The symptoms observed on pineapple after re-infection with these three pathogens were more intense on days 7-8 as compared to the control. There was no visible rot symptom at days 1-3 of inoculation in the treatments, minimal between days 5 and 6 and progressing as time increased till the tenth day of incubation where virulence, measured by the length of the lesion formed was  $6.07 \pm 1.79$  cm,  $4.47 \pm 0.57$  cm and  $6.10 \pm 1.51$  cm for *A. niger*, *M. carribica* and *P. funiculosum* respectively. *A. niger* was observed to be more virulent than *Penicillium funiculosum* but both were more virulent in causing pineapple rot than *Meyerozyma carribica*.

**Table 1: Virulence of the isolated pineapple rot fungi as observed from the pathogenicity test**

<b>Rot formation (cm)</b>				
<b>Days</b>	<i>Aspergillus niger</i>	<i>Meyerozyma carribica</i>	<i>Penicillium funiculosum</i>	<b>Control</b>
Day1	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Day2	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Day3	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Day4	0.00±0.00	0.00±0.00	0.27±0.46	0.00±0.00
Day5	0.37±0.15	0.57±0.51	0.33±0.58	0.00±0.00
Day6	1.87±0.12	0.80±0.69	0.83±1.01	0.00±0.00
Day7	2.7±0.46	1.90±0.10	1.90±0.79	0.00±0.00
Day8	3.2667±0.40	2.27±0.38	2.53±0.74	0.00±0.00
Day9	4.90±1.31	3.73±0.46	5.27±1.45	0.00±0.00
Day10	6.07±1.79	4.47±0.57	6.10±1.51	0.00±0.00
<b>TOTAL MEAN</b>	<b>1.92<sup>a</sup></b>	<b>1.37<sup>d</sup></b>	<b>1.72<sup>b</sup></b>	<b>0.00<sup>e</sup></b>

Total means having the same superscript letter are not significantly different ( $\alpha_{0.05}$ ) according to Duncan Multiple Range Test (DMRT).

### **3.4 Effect of Non-Volatile Compounds of *Trichoderma harzianum* and *Trichoderma longibrachiatum* on Pathogenic Fungi Causing Pineapple Rot**

All the pathogens isolated from pineapple were susceptible to the biocontrol agents (Tables 2 and 3). *M. carribica* was the least susceptible to both *T. harzianum* and *T. longibrachiatum*. There was no inhibition of *M. carribica* at 15% concentration of *T. harzianum* inoculum while inhibition at 20% and 25% for both biocontrol agents was statistically similar. Though the bioagents were effective in inhibiting *A. niger*, no significant difference was observed at 15% and 20% concentrations of both *T. harzianum* and *T. longibrachiatum* which improved at 25% concentration. However, *T. harzianum* significantly inhibited *A. niger*. The inhibition of the pathogen *P. funiculosum* by *T. harzianum* was most effective at 20% and 25% .

UNDER PEER REVIEW

**Table 2: Fungistatic activities of non-volatile compounds *Trichoderma harzianum* against pathogenic fungi causing pineapple rot disease**

Pathogens	Treatments	Day1	Day2	Day3	Day4	Day5	Day6	Day7	Total Mean
<i>Aspergillus niger</i>	15%	0.62±0.06	0.62±0.06	0.62±0.06	0.62±0.06	0.67±0.03	0.67±0.03	0.67±0.03	0.64 <sup>b</sup>
	20%	0.55±0.17	0.57±0.20	0.57±0.20	0.58±0.19	0.70±0.13	0.70±0.13	0.70±0.13	0.62 <sup>b</sup>
	25%	0.53±0.08	0.53±0.08	0.53±0.08	0.53±0.08	0.67±0.20	0.67±0.20	0.67±0.20	0.59 <sup>c</sup>
	Control	0.65±0.30	1.37±0.48	2.18±0.64	3.12±1.03	4.33±0.08	4.38±0.03	4.42±0.08	2.92 <sup>a</sup>
<i>Meyerozyma carribica</i>	15%	2.37±0.88	2.38±0.90	2.57±0.79	2.72±0.79	2.73±0.77	2.73±0.77	2.73±0.77	2.30 <sup>a</sup>
	20%	1.70±0.67	1.70±0.67	1.77±0.71	1.85±0.72	1.85±0.72	1.85±0.72	1.85±0.72	1.80 <sup>b</sup>
	25%	1.23±0.08	1.23±0.08	1.32±0.13	1.33±0.15	1.33±0.15	1.33±0.15	1.33±0.15	1.30 <sup>b</sup>
	Control	1.30±0.17	1.40±0.13	1.75±0.10	2.78±0.41	2.87±0.46	3.07±0.81	3.17±0.75	2.63 <sup>a</sup>
<i>Penicillium foniculosum</i>	15%	0.60±0.13	0.60±0.13	0.60±0.13	0.60±0.13	0.72±0.21	0.72±0.21	0.72±0.21	0.65 <sup>b</sup>
	20%	0.43±0.06	0.43±0.06	0.58±0.08	0.58±0.08	0.63±0.10	0.63±0.10	0.63±0.10	0.56 <sup>c</sup>
	25%	0.42±0.25	0.42±0.25	0.47±0.25	0.48±0.24	0.55±0.22	0.55±0.22	0.55±0.22	0.49 <sup>c</sup>
	Control	0.55±0.05	1.28±0.30	3.07±0.28	3.92±0.71	4.48±0.12	4.52±0.14	4.52±0.13	3.19 <sup>a</sup>

Values are means of three replicates, Total means having the same superscript letter are not significantly different ( $\alpha_{0.05}$ ) according to Duncan Multiple Range Test (DMRT).

**Table 3: Fungistatic activities of non-volatile compound *Trichoderma longibrachiatum* against pathogenic fungi causing pineapple rot disease**

Pathogens	Treatments	Day1	Day2	Day3	Day4	Day5	Day6	Day7	Total Mean
<i>Aspergillus niger</i>	15%	0.68±0.18	0.78±0.25	0.82±0.28	0.87±0.30	0.92±0.38	0.92±0.38	0.92±0.38	0.84 <sup>b</sup>
	20%	0.58±0.06	0.83±0.24	0.85±0.23	0.87±0.23	0.88±0.23	0.88±0.23	0.88±0.23	0.83 <sup>b</sup>
	25%	0.52±0.06	0.57±0.06	0.57±0.06	0.57±0.06	0.63±0.08	0.63±0.08	0.63±0.08	0.59 <sup>c</sup>
	Control	0.65±0.30	1.37±0.48	2.18±0.64	3.12±1.03	4.33±0.08	4.38±0.03	4.42±0.08	2.92 <sup>a</sup>
<i>Meyerozyma carribica</i>	15%	2.25±0.13	2.37±0.25	2.37±0.25	2.37±0.25	2.53±0.15	2.53±0.15	2.53±0.15	2.42 <sup>b</sup>
	20%	0.97±0.25	0.97±0.25	1.03±0.29	1.03±0.29	1.03±0.29	1.05±0.26	1.05±0.26	1.02 <sup>c</sup>
	25%	0.88±0.23	0.90±0.26	0.90±0.26	0.90±0.26	0.95±0.23	0.97±0.21	0.97±0.21	0.92 <sup>c</sup>
	Control	1.30±0.17	1.40±0.13	1.75±0.10	2.78±0.41	2.87±0.46	3.07±0.81	3.17±0.75	2.63 <sup>a</sup>
<i>Penicillium foniculosum</i>	15%	0.72±0.13	0.78±0.12	0.88±0.10	0.88±0.10	0.92±0.08	0.92±0.08	0.92±0.08	0.86 <sup>b</sup>
	20%	0.55±0.10	0.55±0.10	0.85±0.26	0.90±0.23	0.92±0.21	0.92±0.21	0.92±0.21	0.80 <sup>b</sup>
	25%	0.48±0.03	0.52±0.03	0.73±0.08	0.77±0.10	0.78±0.12	0.78±0.12	0.78±0.12	0.69 <sup>bc</sup>
	Control	0.55±0.05	1.28±0.30	3.07±0.28	3.92±0.71	4.48±0.12	4.52±0.14	4.52±0.13	3.19 <sup>a</sup>

Values are means of three replicates, Total means having the same superscript letter are not significantly different ( $\alpha_{0.05}$ ) according to Duncan

Multiple

Range

Test

## 4. DISCUSSION

Analysis of the rot associated fungi of *Ananas comosus* (pineapple) revealed the composition of the mycobiome as *Penicillium funiculosum*, *Aspergillus niger* and *Meyerozyma carribica*. The report by Oniah and Tawose (2018) also confirmed that *Penicillium* spp. and *Aspergillus niger* were among the organisms causing deterioration of pineapple fruits. Likewise, Ogaraku *et al.* (2016) discovered similar fungi in deteriorated pineapple fruit. *M. caribbica* has been found in many samples including in association with fruits such as sugarcane and the Brazilian fruit locally called “camu camu” (*Myrciaria dubia*) (Limtong *et al.* 2014). However, this is arguably the first time *Meyerozyma carribica* has been isolated as a causal agent of pineapple rot. It has been ascertained that fruits are rotten by phytopathogens such as fungi which infect the produce through mechanical injuries during harvesting. The isolated fungi produced similar disease symptoms when inoculated on healthy pineapple fruits. This confirms them as the causal agents of rot of the pineapple fruits, similarly, *Penicillium* spp, *Alternaria alternata* and *Aspergillus niger* have been previously reported as causative pathogens of heart rot of pomegranate fruits (Michailides *et al.* 2012; Zhang *et al.* 2012). It is also in agreement with the submission of Bastein *et al.* (2020) that the genus *Fusarium* and *Talomyces* which share a common ancestor with *Penicillium funiculosum* are causal agents of pineapple rot.

Different inhibitory level was observed among the various pathogens. Non-volatile compounds of *Trichoderma harzianum* and *Trichoderma longibrachiatum* exerted fungistatic effect on *A. niger*, *P. funiculosum* and *M. caribbica* at concentrations of 15%, 20% and 25% in an ascending order of effectiveness. Hanan and Mohamed (2014) reported antagonistic activities of non-volatile compounds of *Trichoderma* sp. against many phytopathogens. The fungistatic effects of non-volatile compounds of *T. harzianum* have been reported by Khaledi and Taheri (2016) where it was reported that non-volatile compounds can decrease the mycelia growth of plant pathogens. Also, non-volatile compound of culture filtrate of *Trichoderma* strain SQR-T037 had significant effect on radial growth of *F. oxysporum* f. sp. *niveum* (Waseem *et al.*, 2013).

## 5. Conclusion

This study revealed the potential of using biopesticides which are biodegradable and nontoxic as a substitute to the use of expensive synthetic fungicides and other methods for plant disease control. The suppression of the fungal pathogens strongly suggests that the active components of the non-volatile compounds of the biological control agents are suitable for the control of fungi causing pineapple rot. Thus, *Trichoderma harzianum* and *Trichoderma longibrachiatum* can be used as biological control agents which could serve as a better alternative to the use of synthetic fungicides.

## REFERENCE

All Africa. Horticulture Can Provide Three Million Jobs in Few Year. 2011. Available:

<http://allafrica.com/stories/201111180784.html?page=2>

Bastien B, Marc C, Anna D, Maeva G, Laetitia R, Mathieu L, Noel D, Lindy JR, Altus V, Sabine Schorr-G. Diversity and Toxigenicity of Fungi that Cause Pineapple Fruitlet Core Rot. *Toxins* 2020; (12)5: 339-342.

Bhardwaj NR, Kumar J. Characterization of volatile secondary metabolites from *Trichoderma asperellum*. *Journal of Applied and Natural Science*. 2017;9:954-959

Chillet M, Hoareau A, Hoarau M, Minier J. Potential Use of Essential Oils to Control Fruitlet Core Rot (FCR) in Pineapple (Queen Victoria Variety) in Reunion Island. *American Journal of Plant Sciences* 2020;11:1671-1681.

Chukwuka KS, Okonko IO, Adekunle AA. Microbial ecology of organisms causing pawpaw (*Carica papaya* L.) Fruit decay in Oyo State, Nigeria. *American-Eurasian J. Toxicol. Sci.* 2010;2:43-50.

Dennis C, Webster J.. Antagonistic properties of species-groups of *Trichoderma*, I. Production of non-volatile antibiotics. *Transactions of the British Mycological Society.* 1971a;57: 41-48

Food and Agricultural Organization of United State. Yearbook. 2019. Website: [www.fao.org](http://www.fao.org).

Hanan IM, and Mohamed OI. The efficacy of non-volatile compound of *Trichoderma* species and *Bacillus* isolates in the control of chickpea wilt pathogens. *Agriculture, Forestry and Fisheries.* 2014;3: 346-351

Ítalo TSRM, Vanderly AS, Giovana DD, Spartaco AF, Edson JC, Marcos JSV. Yeasts with Fermentative Potential Associated with Fruits of Camu-Camu (*Myrciaria dubia*, Kunth) from North of Brazilian Amazon. *The Scientific World Journal.* 2021; 5:1-6.

Khaledi N, Taheri P. Biocontrol mechanisms of *Trichoderma harzianum* against soybean charcoal rot caused by *Macrophomina phaseolina*. *Journal of plant protection research.* 2016; 56: 21-31.

Limtong S, Kaewwichian R, Yongmanitchai W, Kawasaki H. Diversity of culturable yeasts in phylloplane of sugarcane in Thailand and their capability to produce indole-3-acetic- acid. *World J. Microbiol Biotechnol.* 2014; 30:1785-1796

Liu D, Coloe S, Baird R, Pedersen J. Rapid mini-preparation of fungal DNA for PCR. *Journal of Clinical Microbiology.* 2017; 381: 471.

Michaelsen A, Pinzari F, Ripka K, Lubitz W, Pinar G. Application of molecular techniques for identification of fungal communities colonizing paper material. *International Biodeterioration and Biodegradation.* 2006;58: 33-141.

Michailides TJ, Morgan DP, Quist M, Reyes HC. Infection pomegranate by *Alternaria* sp. causing blackheart. <http://ucanr.edu/sites/Pomegranates/files/122813.pdf>. 2012

Moss MO. Mycotoxin review journal on *Aspergillus penicillium*. *Mycologist.* 2002;16: 116-119.

Odebode AC, Joseph CC, Jonker SA, Wachira. SW. Antifungal activities of constituents from *Uvaria schefferi* and *Abotrys brachpetalas*. *Journal of Agricultural Science.* 2006; 51: 79-86.

Okoro CC, Amund OO, Eliora R. 2009. Plasmids of phenanthrene and dibenzothiophene-degrading bacteria isolated from produced water samples in oil production. *Int. J. Biol. Chem Sci.* 2009;3:186-191.

Oniah T, Tawose FO. Fungi associated with black rot disease of pineapple (*Ananas comosus* L.) fruits and the effects of the disease on nutritional value o the fruits. *ISABB Journal of Food and Agricultural Science* 2018; 8(3):18-24.

Onuoroh S C, Udemezue OF, Uche JC, Okoli IC. Fungi Association with the spoilage of Pineapple Fruits in Eke Awka Market Anambra State. *The Bioscientist.* 2013; 1: 22-27

Quijandria G, Berrocal J, Lawrence P. "La industria de la pina en Costa Rica: Analisis de sostenibilidad. Centro Centroamericano dr desarrollo Sostenible" 1997. Available: <http://www.incae.edu/EN/clads/publicaciones/pdf/Cen707.pdf>.

Raza W, Faheem M, Yousuf S, Rayar UF, Yameen M. Volatile and non-volatile antifungal compounds produced by *Trichoderma harzianum* SQR-T037 suppressed the growth of *Fusarium oxysporum* f. sp. *niveum*. *Science Letters.* 2013;1:21-24

Waseem R, Muhammad F, Sohail Y, Faheem UR, Muhammad Y. Volatile and nonvolatile antifungal compounds produced by *Trichoderma harzianum* SQR-T037 suppressed the growth of *Fusarium oxysporum* f. sp. *niveum*. Science letters.2013;1(1): 21-24.

Zakawa NN, Timon D, Yusuf CS, Tizhe TD, Bala UJ, Isa A, Waja S, Aphonsus G. Isolation and Control of Fungal Rot Pathogen of Tomato Fruit Using Aqueous Leaf Extracts of *Azadirachta indica* in Mubi, Adamawa State. Research Journal of Plant Pathology. 2019; 2: 1-5

Zhang LU, McCarthy MJ. Black heart characterization and detection in pomegranate using NMR relaxometry and MR imaging. Postharvest Biol. Technol. 2017;67:96–101.

UNDER PEER REVIEW