

# **A REVIEW ON THE RELATIONSHIP OF CLIMATE VARIABILITY AND EXTREMES WITH CROP PRODUCTION**

## **ABSTRACT**

Risks are always present in agriculture for a variety of reasons. Climate-related risks are the most significant among them since they can occur unexpectedly and cannot be avoided. The main climatic factors influencing crop productivity are rising temperatures, altered precipitation patterns, and rising atmospheric CO<sub>2</sub>. The average global temperature is expected to rise by 2°C until 2100, which would result in significant global economic losses. The average global temperature is currently rising steadily. The concentration of CO<sub>2</sub>, which makes up a large amount of greenhouse gases, is rising alarmingly. Climate change, with rising temperatures and increased greenhouse gas emissions, impacts agriculture significantly, leading to varied crop yields and potentially catastrophic economic consequences. While some regions may experience favorable effects, overall, climate variability poses challenges such as reduced crop productivity, increased pest activity, and elevated food costs, particularly affecting underdeveloped nations. Environmental policies must be adaptable and flexible to mitigate these impacts effectively. Farmers' responses to climate change, influenced by perceptions and data accessibility, drive both mitigation and adaptation efforts. Sustainable agriculture, conservation practices, and technological innovations play key roles in mitigating climate change impacts and enhancing resilience in farming communities, though success depends on multiple factors including local perceptions, technical feasibility, and economic viability. This paper reviews the causes of climate change, the climate variables affecting crop production and the mitigation and adaptation strategies against climate change.

Keywords: Climate Change, Agriculture, Climate variability, Mitigation, Adaptation.

## **INTRODUCTION**

There's mounting evidence that the earth's temperature has increased during the past century, leading to unequalled and unpredictable variations in the climate and negative effects on people's lives everywhere. (IPCC (Intergovernmental Panel on climate change, 2022). Due to agriculture's heavy reliance on weather, climate change has slowed down agricultural growth globally and is predicted to have a major impact on crop production. (IPCC, 2022). The numerous documented instances of ongoing crop yield declines worldwide serve as evidence of the severe effects of the global climate on crop production. (Bednar-Friedl et al. 2022, Trisos et al. 2022).

The rise in global air temperature brought on by higher greenhouse gas concentrations is an impending change in the climate (Mall et al. 2006). Numerous studies have demonstrated that India has experienced an unparalleled rise in surface temperature over the past century. Compared to the previous century, the average global temperature has risen by 0.8°C (IPCC, 2013). Approximately 7.3 billion people live on Earth today; by 2050, that number is predicted to rise to 9.7 billion, and by 2100, it will reach 11.2 billion (UN, 2015). By 2100, food production will need to increase several times over current levels in order to ensure food security for the world's

rapidly growing population. Both the Indian and South Asian regions are heavily populated, have poor economies, and rely heavily on agriculture—which is susceptible to the effects of climate change. India's population could reach 1.66 billion by 2050 (UN, 2015).

In South Asia, climate change poses a threat to sustainable development because of the region's high population density, extreme poverty, and lack of resources for adaptation. Thus, climate change is expected to severely harm the region's economy, society, and environment, undermining prospects for growth and initiatives to combat poverty (Ahmed et al. 2014). Because climate change affects crop growth and output, hydrologic balances, input sources, and other management methods, agriculture is particularly sensitive (Knox et al. 2012). The impacts of climate change on agriculture are numerous and include variations in sea level, pest and disease conditions, average temperatures, rainfall, and weather extremes, among other things (Niang et al. 2014; Porter et al. 2014). Increased temperatures cause early flowering and a shorter grain-filling period, which shorten crop cycles and lower yield per unit area ( Chattopadhyay, 2011). High temperatures during the day and at night are expected to increase in the near future and pose a serious environmental threat to food production and security worldwide (Lobell et al. 2011; Cairns et al. 2012; Hijjoka et al. 2014). According to Randhawa et al. (2014), future climatic variability will also result in more frequent extreme weather events, such as unpredictable monsoons and an increase in the frequency and severity of drought and flooding, which will have an impact on both rainfed and irrigated agriculture systems.

With the minimum temperature rising at twice the rate of the maximum temperature, the diurnal temperature range has likewise shrunk throughout India (Roy and Balling, 2005). Over the Indian subcontinent, climate forecasts suggest an annual mean temperature increase between 3.5°C and 5.5°C by the 2080s, with the relative increase being smaller in the rabi (winter) season than in the kharif (monsoon) season (Lal et al. 2001).

Rainfall is expected to increase in the kharif season by 2050, despite a tendency to decline in the rabi season (Lal et al. 2001; Prabhakar et al. 2008; Randhawa et al. 2014). Consequently, it is anticipated that rising temperatures and erratic rainfall will have a variety of effects on agriculture in India, endangering the food security and means of subsistence for over 700 million rural residents (Chattopadhyay, 2011; Dagar et al. 2012). Global climate change has occurred, although its effects frequently differ from place to place and even from region to region (Trajkovic, 2009; Hadgu et al., 2014).


Therefore in this paper we discuss the relationship between effect climate variability and extremes on crop production. The relationship is discussed under the following headings; the unpredictability of climate change and its causes, the relationship between agriculture and climate change, climate variability versus crop production, climate change related stressors and their effects on crop adaptation, effect of climate change on physiological, metabolic and morphological processes in plants and climate change mitigation and adaptation. The paper therefore helps to understand the effect of climate change on agriculture and the mitigation strategies as well.

### **The unpredictability of the climate and its causes**

Many studies have been conducted on the significant climate variability (mean computation and variability of linked parameters of variables, such as temperature, rainfall, and wind during certain periods of time). According to estimates, climatic variability is rising as a result of rising temperatures. In many respects, the physiology

of plants is significantly impacted by climate fluctuation. While most tropical and subtropical regions have seen a drop in precipitation, high latitudes have seen an increase. Animals as well as crop plants are under more stress due to climate variability and extreme weather events (Thornton et al. 2014).


The implications of climate change, especially its biological effects, have become abundantly clear in the last several years. Due to human activity and natural sources, there is an increase in the emission of gases such as carbon dioxide, methane, nitrous oxide, and halocarbons. These gases absorb solar radiation and contribute to the greenhouse effect (Valizadeh et al. 2014). Drought is caused by high temperatures, little precipitation, salt stress, and intense light (Salehi-Lisar and Bakhshayeshan, 2016).



Ahmad Khan, 2012

### The relationship between agriculture and climate change

Due to its vast scope and extreme weather sensitivity, agriculture is the industry most susceptible to climate change, with potentially catastrophic economic consequences (Mendelsohn, 2009). The amount of crop output is greatly impacted by variations in meteorological events like temperature and rainfall. The crop, location, and degree of parameter change all affect the effects of rising temperatures, fluctuating precipitation, and CO<sub>2</sub> fertilization. It has been discovered that rising temperatures lower yields, while rising precipitation is likely to neutralize or lessen the effects of rising temperatures (Hatfield et al. 2011). Crop productivity is dependent on crop type, climate scenario, CO<sub>2</sub> fertilization effect, and adaptability skills when it comes to climate variables as observed in Iran (Karimi et al. 2018).



Impact of climate change on agriculture. (Ullah et al. 2021)

Depending on the region and irrigation technique, agricultural yields are affected differently by climate change. Extending irrigated regions can boost crop yields, but doing so may have negative environmental effects (Kang et al. 2009). As a result of their shorter lifetime, several crops are expected to yield less due to temperature rise (Mahato, 2014). If both the temperate and tropical regions undergo a warming of 2°C, the total production of wheat, rice, and maize is anticipated to decline (Challinor et al. 2014). Because tropical crops stay closer to their high-temperature optima and so face high-temperature stress under rising levels of temperature, climate change generally has a greater impact on tropical regions.

Furthermore, humid and warmer regions tend to have higher rates of disease and insect pests (Malhi et al. 2021). Crop yields are also affected by other factors like humidity, wind speed, temperature, and rainfall; in the absence of these factors, there has been a potential for overestimating the cost of climate change. Reduced crop yields have the potential to drive up food costs and have a disastrous impact on agricultural wellbeing worldwide, with an estimated 0.3% yearly loss of future global GDP by 2100 (Stevanovic et al. 2016).

However, (Bosello and Zhang, 2005) discovered that while climate change has little impact on the world's food supply, underdeveloped nations will suffer greatly as a result. The agriculture industry in India may suffer as a result of expected temperature increases of 2.33 to 4.78°C, a doubling of CO<sub>2</sub> concentration, and longer heat waves (Kumar and Gautam, 2014).

Temperature and precipitation variations have a significant impact on plant-water relations, and physiological changes are more likely to be affected by abrupt shifts in these variables than by variations in the average climate (Reyer et al. 2013). The way that different plant species and developmental stages react to climate change is evident. Different plant species have different thresholds, and different plant species respond differently—for example, by elongating their roots, altering their root development angle, or producing less of a crop (Grey and Brady, 2016). Plant

transpiration was shown to be reduced with increased CO<sub>2</sub> level in the environment, resulting in an increase of  $0.42 \pm 0.02$  K in air temperature. Land surfaces can warm by  $3.33 \pm 0.03$  K due to a direct radiative effect and the indirect physiological effect of increased CO<sub>2</sub> (Cao et al. 2010). Harvestable crop yields are predicted to rise in response to rising atmospheric CO<sub>2</sub> levels, and plant developmental changes vary depending on the kind of crop. While higher yields are anticipated for C3 crops, decreased water requirements are anticipated for both C3 and C4 crops in the absence of stressful situations. However, higher temperatures and changed precipitation are expected to counterbalance these positive effects of increased CO<sub>2</sub> (DaMatta et al. 2010).

On the other hand, some regions also show that climate change has a favourable effect on agricultural productivity. However, these regional variations—whether they were increases or decreases—would not cause significant alterations, and they would only become more noticeable in a few low latitudes. On the other hand, significant economic losses may result from temperature increases greater than those caused by doubling CO<sub>2</sub> (Aydinalp and Cresser, 2008). The tropical regions of developing countries will bear a heavy burden from climate change, however the exact impact will primarily depend on local climate conditions. According to Zilberman et al. (2012), environmental policies need to be dynamic and executed with adaptability and flexibility, as the pace of climate change influences its impact and, in turn, the cost of adjustment.

### **Climatic variables versus crop production**

Research clearly demonstrated that an increase in global temperature would have a negative impact on agriculture in the Indian subcontinent. According to Joshi and Amalkar (2009), temperature fluctuations, precipitation, carbon dioxide fertilization, short-term weather variability, and surface water runoff are the primary climatic elements that could impact agricultural output. Both good and negative effects on agricultural productivity are attributed to these factors. Temperature and precipitation trends in the climate have had, and will continue to have, a major influence on agriculture. Individuals of the same or different species will respond differently to changes in light, nutrients, water, or CO<sub>2</sub> (Fernando et al. 2016); as a result, heterogeneity within or between species can be expected when atmospheric CO<sub>2</sub> levels rise globally. With the right fertilization, variety selection, and optimal nutrient supply, adverse agro-ecological weather effects can be mitigated.

**Temperature and crop production:** Higher average temperatures in all major cropping zones have been and will continue to be the most pervasive aspect of climate change. In most places, the temperature thresholds for crops, such as 35°C or 40°C, will be exceeded on more days due to this increase in average temperatures alone. Global agriculture will typically suffer from temperature increases exceeding 2.5°C (IPCC, 2001). Compared to precipitation, the growing trend towards global warming over the 20<sup>th</sup> century has been remarkably noticeable. Global warming is predicted to cause a 9–21% decrease in total agricultural productivity in developing nations, according to Cline (2007). Increases in temperature in mid- to high-latitude regions lead to higher yields; however, the impacts become less pronounced when temperature fluctuations above 30°C. Temperature variations may result in air vapour pressure deficits, which may have an effect on how much water is used in agricultural landscapes (Kirschbaum, 2004).

This may result in changes in temperature and water loss, as well as an impact on the rate of transpiration. According to reports, soil warming may have a deleterious impact on the growth, shape, and longevity of roots (McCormack et al. 2014). Numerous studies have shown that variations in soil temperature can affect the way that  $\text{NH}_4^+$ ,  $\text{NO}_3^-$ ,  $\text{P}_2\text{O}_5$ , and  $\text{K}^+$  are transported by roots, hence having a direct impact on plant nutrient absorption. Soil warming may promote plant nutrient acquisition by improving root uptake kinetics, as seen by improved plant nutritional status when root development and morphological traits are severely affected (BassiriRad, 2000). High temperatures impair the processes that allow light to be absorbed and transformed into energy, which causes photorespiration to rise.

It is possible for the light-harvesting chlorophyll protein complex to irreversibly split apart from the nucleus of the photochemical reaction centre, and there may also be harm to the oxygen-producing water splitting mechanism. Higher temperatures so typically result in lower yields since they accelerate a plant's growth to maturity earlier, thereby shortening the time available for yield generation. They also frequently exacerbate the stress on water resources, which are crucial for crop growth. Since the air's capacity to store water increases non linearly with temperature, higher average temperatures will also result in higher rates of evapo-transpiration (ET). Higher ET rates will therefore tend to dry up the soil, increasing the frequency of low moisture extremes. If the yearly variability of the climate were to rise, extremes might also become more frequent in addition to variations in the average temperature. Pests, illnesses, and weeds are also more likely to be present in areas that are warmer and wetter.

**Radiation and crop production:** The only known source of energy for the universe's ecology, either directly or indirectly, is solar radiation. Photosynthesis and crop productivity are significantly impacted by radiation (Gholipour and Shahsavani, 2008). Crop development and production are derived from photosynthesis and are reliant on the receipt and absorption of solar energy, as demonstrated by Reynolds et al. 2012. According to Richards (2006), seasonal fluctuations in solar radiation have a significant impact on yield. This notable crop response trend to solar radiation could be the result of a change in photosynthesis' duration. Due to the unequal partitioning of light, sowing time can be used to introduce variability in solar radiation, which eventually affects the duration of crop growth.

A longer period of time spent exposing a crop to favourable environmental conditions may also result in good seed establishment and yield. On the other side, owing of cloud cover's reduction of solar radiation, growth and development would turn negative. Solar radiation is a significant environmental element that modifies light partitioning and leaf architecture, promoting favourable alterations in crop growth. Any decrease in solar radiation will significantly lower agricultural productivity because it is closely linked to crop growth. The concomitant rise in minimum temperatures causes the crops to require more respiration, which lowers net growth and productivity (Aggarwal, 2003). Additional research points to a declining trend in solar radiation. Recently, it has been shown that gases and particulate matter (aerosols) released into the atmosphere by human activity have disrupted the earth's surface's ability to intercept solar radiation (Yamasoe et al. 2006).

**Precipitation and crop Production:** Thirty percent of the world's population lives in areas of the globe that are water stressed (McCarthy, 2001). With global warming, the hydrological regimes in which crops flourish will undoubtedly alter. The availability

of water is a significant factor in agricultural output fluctuation in various situations (Keating et al. 2010). Variations in the amount of precipitation throughout the year, within-season patterns, and seasonal variations may also have an impact on the crop water regime. Stronger convection cells and more air moisture are expected to result in increased convective rainfall, especially in the tropics. Over the course of the 20<sup>th</sup> century, precipitation has most likely grown by 5–10% over the majority of the mid- and high-latitude continents in the northern hemisphere, whereas it has declined by 3% on average over the majority of the subtropical land area. Precipitation is widely acknowledged to be a major element influencing crop productivity, particularly for rainfed crops (Izaurre et al. 2003). While insufficient precipitation can harm crop output, particularly if dry spells happen during crucial developmental stages, excessive precipitation can lead to disease infestation in crops. Changes in the precipitation and seasonal and yearly evapotranspiration regimes will have an impact on the availability and quantity of water held in the soil, which is an essential component of crop growth. The need for irrigation water is projected to increase due to global climate change (Jones et al. 2017).

**CO<sub>2</sub> increase and crop production:** Since carbon dioxide is a fundamental resource for plant growth, the continuous rise in its concentration would enable breeders to start choosing the best cultivars from the existing crop lines. It is the belief of breeders and agronomists that empirical selection for the fastest-growing, highest-yielding cultivars under current environmental conditions will lead to CO<sub>2</sub> responsiveness; these conditions will eventually reflect changes in the background concentration of CO<sub>2</sub> (Bouras et al. 2019).

According to projections made by the Intergovernmental Panel on Climate Change (IPCC, 2001), atmospheric CO<sub>2</sub> concentrations will rise from 368 µmol/mol in 2000 to a range of 540 to 970 µmol/mol in 2100. Temperature rises are predicted to coincide with increases in the atmospheric concentration of CO<sub>2</sub>. According to IPCC predictions, air temperatures will rise by 1.4 to 5.8°C between 2000 and 2100 as a result of greenhouse gas accumulation in the atmosphere. These worldwide trends will be supported by significant regional variances (White et al. 2004). Higher CO<sub>2</sub> concentrations cause plants to develop at faster rates of net photosynthesis and/or smaller stomatal diameters (Kobiljski and Dencic, 2001). Positively, some crops are predicted to benefit from the increased atmospheric concentration of carbon dioxide. Reduced transpiration per unit leaf area caused by partial stomatal closure, when combined with increased photosynthesis, frequently results in improved WUE (Neenu et al. 2013).

As a result, higher CO<sub>2</sub> concentrations can boost productivity while lowering water consumption. Under conditions of doubled CO<sub>2</sub>, these C<sub>3</sub> crops can experience growth rates of up to 50% (Dixon. 2009). Compared to tropical crops, temperate crops might gain more from higher CO<sub>2</sub>. It has been demonstrated that CO<sub>2</sub> enrichment reduces photorespiration, or the quick oxidation of recently generated sugars in the light, in crop species that have the C<sub>3</sub> pathway characteristic of non-tropical plants (such as wheat, soybean and cotton). This reduces the efficiency of photosynthesis as a whole. Because C<sub>4</sub> crops fix CO<sub>2</sub> into malate in their mesophyll cells before transferring it to the RuBP enzyme in the bundle-sheath cells, they are more photosynthetically efficient than C<sub>3</sub> plants under current CO<sub>2</sub> levels. C<sub>4</sub> crops are particularly characteristic of tropical and warm arid regions (e.g., maize, sorghum, and millet). Experimental studies indicate that C<sub>4</sub> plants are less susceptible to CO<sub>2</sub>

enrichment because of this CO<sub>2</sub>-concentrating and photorespiration-avoiding mechanism (Prior et al. 2011).

### **Climate change-related stressors and their effects on crop adaptation**

Because of climate fluctuations, the impacts of abiotic pressures on crop adaptability and productivity have been reported (Nowsherwan et al. 2017). Abiotic variables cause abiotic stressors in plants. Extremes in temperature (stress from heat or cold), light, radiation, water (stress from flooding and drought), chemicals (metals and pH), and gaseous pollutants (ozone, sulphur dioxide) are some of these causes. According to Pereira (2016), heat and drought are two common field stressors that significantly affect plant performance and that of their progeny. They can also cause a plant's persistent problems.

Because plants need an ideal range of temperatures for both growth and adaptation, the range of temperatures surrounding any species' plants determines how much of that species' output there is. Temperature changes have an equally important impact on plant phenology (Hatfield and Prueger, 2015). Drought stress negatively impacts plant physiology, morphology, and biology. Heat stress decreases grain production and grain-filling duration. Frost causes sterility and abortion of grains (Barlow et al. 2015).

While climate change may benefit agriculture in certain parts of the world where the temperature is above 55°C, many plants, particularly those that are native to warm environments, are negatively impacted by these changes. Severe summertime weather events pose major risks to crop adaptability and productivity. For example, a rise in temperature, evaporation, and transpiration causes a wheat plant to shorten its flowering period from planting to emergence. A 1°C temperature increase shortens flowering by 5 days. Temperature increases also affect the grain filling period, which ultimately results in a decrease in biomass (Valizadeh et al. 2014).

### **Plant Growth Affected by Various Stresses Caused by Climate Change**

Stress	Induced Stresses	Secondary	Effects in Plant	References
Chilling/freezing stress	Nutritional imbalance, osmotic and oxidative stress		<ul style="list-style-type: none"> <li>• Buildup of reactive oxygen species (ROS) and oxidative damage; suppression of enzyme function and imbalance in metabolism.</li> <li>• Reduced photosynthetic activity, senescence, delayed maturity, increased cell starvation and dehydration, and damage to PS II.</li> <li>• Reduced development and efficiency</li> </ul>	Esim and Atici, 2014; Megha et al. 2018
Drought	Osmotic, heavy metal, and oxidative stress		<ul style="list-style-type: none"> <li>• Increased ROS production and ion leakage; induced dehydration and turgor loss.</li> <li>• Decrease in absorption and translocation of mineral</li> </ul>	Ansari and Lin, 2010; Iqbal et al. 2020

		<p>nutrients.</p> <ul style="list-style-type: none"> <li>• Protein denaturation, loss of enzyme activities, reduced photosynthetic activity due to abridged chlorophyll content and CO<sub>2</sub> assimilation.</li> <li>• Elevated leaf temperature, necrosis, early abscission, and slowed plant development.</li> </ul>	
Flooding/ waterlogging	Water and nutrient deficiency stress, oxidative stress	<ul style="list-style-type: none"> <li>• Reduced absorption and translocation of mineral nutrients; increased generation of reactive oxygen species (ROS) and ion leakage; induced dehydration and turgor loss.</li> <li>• Denaturation of proteins, loss of enzyme activity, and decreased photosynthetic activity as a result of lower CO<sub>2</sub> absorption and chlorophyll concentration.</li> <li>• Stunted plant development, early abscission, necrosis, and elevated leaf temperatures.</li> </ul>	Bailey-Serres and Voeselek 2008; Ashraf, 2012
Heat stress	Water scarcity, osmotic and oxidative stress	<ul style="list-style-type: none"> <li>• Increased oxidative damage, denaturation, misfolding of proteins, and generation of ROS.</li> <li>• Reduced CO<sub>2</sub> fixation, foliar senescence and abscission, growth inhibition, fruit and leaf discoloration, disturbance of PS I and PS II, and disrupted ion transport.</li> </ul>	Nievolala et al. 2017; Sarkar et al. 2018; Demirel et al. 2020
Light/radiation stress	Oxidative stress	<ul style="list-style-type: none"> <li>• Increased oxidative damage and ROS generation, interrupted photosynthesis ETC, and/or increased activity of membrane-bounded NADPH oxidase, degraded chlorophyll, decreased photosynthetic activity, and suppression of epidermal cell growth.</li> <li>• Condensed inflorescence stem with an increased number of flowering stems, decreased rosette diameter, and leaf senescence.</li> </ul>	Noshi et al. 2016
Nutrient imbalance	Oxidative stress	<ul style="list-style-type: none"> <li>• Reduced antioxidants leading to a buildup of reactive oxygen species (ROS); increased ion and</li> </ul>	Bisht et al. 2019; Saleem et

		<p>solute leakage; decreased metalloenzyme activity; and decreased photosynthesis.</p> <ul style="list-style-type: none"> <li>• Vulnerability to additional biotic and abiotic factors.</li> <li>• Growth retardation, chlorosis, necrosis, little fruit and flowering, and decreased output</li> </ul>	al. 2020
Ozone (O <sub>3</sub> ) stress	Oxidative stress	<ul style="list-style-type: none"> <li>• The generation of ROS results in oxidative damage, reduced stomatal conductance, decreased photosynthesis, slowed enzyme activity, and degradation of chlorophyll and xanthophyll.</li> <li>• Reduced plant biomass and productivity, early senescence, and chlorosis and necrosis of the leaves</li> </ul>	Ainsworth et al. 2012; Ueda et al. 2013
Salinity	Water scarcity, ionic imbalance, nutrient, osmotic and oxidative stress	<ul style="list-style-type: none"> <li>• The generation of ROS results in oxidative damage, decreased K<sup>+</sup>, Ca<sup>2+</sup>, and Mg<sup>2+</sup> content, restricted water and mineral nutrient uptake and translocation leading to Na<sup>+</sup> toxicity, and decreased soil water potential.</li> <li>• Reduced photosynthesis, disordered thylakoid ultrastructure and decreased stomatal opening.</li> <li>• Decreased seed germination, premature leaf withering, as well as reduced productivity and growth</li> </ul>	Kumar et al. 2017; Rahman et al. 2016; Jalil et al. 2020

### Effects of climate change on physiological, metabolic, and morphological processes in plants

Plants are experiencing unique ecological settings that fall outside of their optimal range for adaptation as a result of the rapid changes in climate conditions. The significant variations in temperature and precipitation patterns may make plant migration an impractical strategy. Despite the fact that plants have modified their physiology for benefit in unique settings, but according to Becklin et al. (2016), climate change might be just as harmful as pushing plants over their tolerance limits. Many plant species' morpho-biochemical processes are significantly impacted by abiotic stressors (Jan et al. 2016). However, according on agricultural physiology responses to anticipated climatic conditions for the next few years, crops will grow more quickly and, depending on the crop variety and specie, may see modest changes in blooming and fruiting (DaMatta et al. 2010). For plants, 10 to 35°C is the ideal temperature range. According to Tkemaladze and Makhashvili (2016), plants can only

create a limited amount of energy as temperatures rise. After that, photosynthesis in the leaves drastically declines and is irreversibly lost.

Plant turgor pressure is lowered during drought, which eventually restricts cell development. Lack of water affects photosynthetic enzyme activity, lowers metabolic process efficiency and ultimately causes failure in the photosynthetic apparatus (Zargar et al. 2017). A rise in CO<sub>2</sub> concentration brought by shifting weather patterns causes a decrease in plant respiration and an increase in temperature. Temperature increases cause crop respiration to rise up to a maximum of 15–40°C before declining (Amedie, 2013). The enzyme Rubisco is a component of the carbon fixation process, which transforms atmospheric carbon dioxide into molecules with high energy content in photosynthetic organisms. Rubisco activase removes metabolites to maintain Rubisco active within typical temperature ranges. Rubisco deactivates at low temperatures because it produces chemicals that are inhibitory (xylulose-1, 5-bisphosphate). Rubisco activase also denatures over the optimal range of species temperature, forming insoluble aggregates that are unable to remove inhibitors and activate Rubisco. Temperature elevation has been shown to cause a very quick drop in the activation state of Rubisco in cotton (Nasim et al. 2017). Enzymatic and non-enzymatic antioxidant defence mechanisms regulate reactive oxygen species (ROS), which are byproducts of cellular metabolisms and include hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), super oxides (O<sub>2</sub><sup>-</sup>), hydroxyl ions (OH), and singlet oxygen (O<sub>2</sub><sup>1</sup>). ROS are predominantly produced by cells and organelles at low concentrations under normal environmental settings, but their concentration rises under stressful circumstances (Sage et al. 2008). Plants that produce too much reactive oxygen species (ROS) pose a threat to proteins, lipids, and DNA, ultimately causing cellular damage and death (You and Chan 2015).

### **Climate Change Mitigation and Adaptation**

Farmers' perceptions of the severity and threat of climate change are the primary drivers of voluntary mitigation. However, the adaptation is contingent upon the accessibility of pertinent data (Semenza et al. 2011). The anticipated rise in extreme weather events, alongside diminishing soil health and biodiversity, may elevate the cost of food production. The United Nations advocates for global intervention through 17 sustainable development goals (SDGs), four of which pertain to food production and security. These include tackling biodiversity decline (SDG 15), addressing ecosystem service loss and agroecosystem instability due to heightened stress from intensified food production and climate change (SDG 13), combatting soil health deterioration stemming from agricultural methods (SDGs 2 and 6), and reducing reliance on synthetic fertilizers and pesticides for enhanced productivity (SDG 2). Achieving these SDGs necessitates the agricultural sector's proactive engagement in reversing prevalent negative environmental trends. This entails implementing significant changes in agricultural practices to ensure resilience and adaptation to climate change by 2030 and beyond. (Shahmohamadloo et al. 2022).

Additionally, fewer individuals will be subjected to water stress thanks to mitigation methods; nevertheless, those who remain will still require adaptation strategies because their stress levels will be elevated (van Vuuren et al. 2011). Farmers can embrace climate-resilient technologies with the assistance of agroecological and traditional management techniques, such as biodiversification, soil management, and water collection (Altieri and Nicholls, 2017). According to Lal et al. (2011), these management techniques guarantee improved soil health, reduced soil erosion, enhanced carbon sequestration, and resilient agricultural systems that will ultimately

guarantee food security in the face of climate change. The most effective educational interventions for promoting climate change awareness for ecological development are those that centre on local, palpable, and doable issues that can be observed through individual behaviour (Anderson, 2012). The majority of farmers supported adaptations, but only a small percentage supported reducing greenhouse gas emissions. This indicates the need to concentrate on programmes that include mitigation and adaptation elements (Arbuckle et al. 2015). Ventakeswarlu and Shanker (2009) have grouped the primary adaptation strategies of mitigation into three categories: resource-conservation technologies, cropping-system technologies, and socio-economic or policy initiatives. Small and marginal farmers are particularly vulnerable to losses because they lack the awareness necessary to adapt to climate change (Baul and McDonald, 2013). A few easy ways to reduce greenhouse gas emissions are rice that is alternately dried and drained in the middle of the season, better animal nutrition, increased N-use efficiency, and soil carbon. It may be possible to lessen the effects of climate change by implementing straightforward adaptation techniques such as altering planting dates and types (Aggarwal, 2008). Technology diffusion has a significant impact on how farmers react to climate change. According to Lybbert and Sumner (2012), the primary areas of focus are capacity building, market integration, and support for public research. Because conservation agriculture promotes minimal soil disturbance, crop diversity, and soil cover management, it has the ability to undo the damage caused by conventional ploughing over time. Additionally, conservation agriculture increases terrestrial carbon absorption, decreases fertilizer consumption, and reduces greenhouse gas emissions (Pagnani et al. 2019). Sustainable agriculture approaches are made possible by conservation agriculture, which is based on the fundamental concepts of minimum soil disturbance, crop rotation, and soil cover. Farmers in south Asia are cultivating wheat with zero tillage mainly due to a 15–16% decrease in cultivation costs. Furthermore, according to Erenstein et al. (2012), zero tillage produces higher yields of wheat and maize with less variability. No-till farming was also promoted as a substitute for traditional tillage, which reduces the effects of climate change by sequestering carbon. However, this claim is overstated because no-till farming adds very little additional organic carbon to the soil (Powlson et al. 2014). The adoption of conservation agriculture (CA) can be attributed to a number of factors, including the perception of personal benefits, the development of farmer organizations to support local adaptation, the use of functional market exchange techniques to supply the necessary resources for CA implementation, and the alliances between farmer organizations and institutions to create appropriate environments. (Brown et al. 2018).

Modified farming methods are the primary means of adapting to climate change. These practices are heavily influenced by policy decisions that take into account social, political, and economic factors as well as climate extremes and variability (Smit and Skinner, 2002). Nutrient management is crucial since the conventional intensification of agriculture results in enormous economic losses, of which over 80% are attributable to nutrient mismanagement (Lu et al. 2015). Soil restoration, nitrogen management, cover crops, manuring, no-till farming, and agroforestry can all promote carbon sequestration, or an increase in soil organic carbon (SOC). One irrigation method that is being promoted to lessen groundwater overdraft and shocks brought on by climate change is drip irrigation. It lessens the need for groundwater for irrigation and has the potential to be climate change resilient. However, farmers are employing drip irrigation in their intensive farming practices, which is producing more groundwater over extraction and the Jevons dilemma (Birkenholtz, 2017). Sprinkler

and drip irrigation are two examples of water-saving irrigation methods that can both reduce and adapt to climate change and offer long-term economic benefits. Reduced N application can be achieved through agricultural practices that take into account site-specific information without compromising profitability. For this reason, precision farming is thought to be more profitable than field management (Bongiovanni and Lowenberg-Deboer, 2004). One strategy for adjusting to environmental challenges in plants is to breed them to create new types. In order to test a variety's appropriateness for the target environment, multilocation experiments, breeding cycle shortening, and germplasm selection will be necessary (Atlin et al. 2017; Chhogyell et al. 2016).

It is crucial to create stress-tolerant cultivars as a mitigating measure because it is anticipated that climate change will increase the frequency and severity of abiotic stress. By accumulating evidence, boosting the efficacy of local institutions, advocating for climate-smart agricultural policies, and connecting agricultural financing to climate, climate-smart agriculture increases resilience to climate change (Lipper et al. 2014).

The most effective climate-smart technologies are those that either support soil structure or supply water or nutrients. These mitigation techniques offer enormous potential for both adaptation and mitigation. But they also rely on factors like people's perceptions, technical complexity, economic viability, and a technology's suitability for the area. Furthermore, these tactics function best when several interventions are applied in tandem and in support of one another.

## **Conclusion**

The world's most serious issue right now is climate change. Climate has a well-known impact on crop quality and quantity, making it a significant and independent component in agriculture. The burden of maintaining global food and nutritional security due to population growth has placed significant strain on agriculture, a situation that is made worse by climate change. In a typical climate, the weather during the growing season mostly determines crop growth, development, and production. Little variations from the typical weather have a significant negative impact on food output, applied input efficiency, and weather. Climate change will reduce agricultural productivity in the coming years, according to a number of studies, notwithstanding the uncertainty surrounding the future climate scenario and its potential effects. The primary determinants of climate, which are temperature, precipitation, and greenhouse gas emissions, have a substantial negative impact on plant physiology, pest infestation, soil fertility, irrigation resources, and metabolic processes. Agronomic choices, tactics, and interventions that result in climate-resilient farming systems can be guided by an understanding of how the climate is changing and will change locally, as well as the degree of the change and the opportunities and hazards it presents.

## **REFERENCES**

Aggarwal PK. Impact of climate change on Indian agriculture. *Journal of Plant Biology-new Delhi*. 2003;30(2):189-98.

Aggarwal PK. Global climate change and Indian agriculture: impacts, adaptation and mitigation. *Indian Journal of Agricultural Sciences*. 2008;78(11):911.

Ahmed M. Assessing the costs of climate change and adaptation in South Asia. Asian Development Bank; 2014.

Ahmed N, Khan TI, Augustine A. Climate change and environmental degradation: a serious threat to global security. *European Journal of Social Sciences Studies*. 2018.

Ainsworth EA, Yendrek CR, Sitch S, Collins WJ, Emberson LD. The effects of tropospheric ozone on net primary productivity and implications for climate change. *Annual review of plant biology*. 2012;63:637-61.

Altieri MA, Nicholls CI. The adaptation and mitigation potential of traditional agriculture in a changing climate. *Climatic change*. 2017;140:33-45.

Amedie FA. Impacts of climate change on plant growth, ecosystem services, biodiversity, and potential adaptation measure. MasterThesis. Program Study of Biological and Environmental Science, University of Gothenburg, Sweden. 2013.

Anderson A. Climate change education for mitigation and adaptation. *Journal of education for sustainable development*. 2012;6(2):191-206.

Ansari MI and Lin TP. Molecular analysis of dehydration in plants. *Int Res J Plant Sci*. 2010;1(2):21-5.

Arbuckle Jr JG, Morton LW, Hobbs J. Understanding farmer perspectives on climate change adaptation and mitigation: The roles of trust in sources of climate information, climate change beliefs, and perceived risk. *Environment and behavior*. 2015;47(2):205-34.

Ashraf MA. Waterlogging stress in plants: A review. *African Journal of Agricultural Research*. 2012;7(13):1976-81.

Atlin GN, Cairns JE, Das B. Rapid breeding and varietal replacement are critical to adaptation of cropping systems in the developing world to climate change. *Global food security*. 2017;12:31-7.

Aydinalp C, Cresser MS. The effects of global climate change on agriculture. *American-Eurasian Journal of Agricultural & Environmental Sciences*. 2008;3(5):672-6.

Bailey-Serres J, Voisenek LA. Flooding stress: acclimations and genetic diversity. *Annu. Rev. Plant Biol.*. 2008;59:313-39.

Barlow KM, Christy BP, O'Leary GJ, Riffkin PA, Nuttall JG. Simulating the impact of extreme heat and frost events on wheat crop production: A review. *Field crops research*. 2015;171:109-19.

Bassirirad H. Kinetics of nutrient uptake by roots: responses to global change. *The New Phytologist*. 2000;147(1):155-69.

Baul, Tarit Kumar, and Morag McDonald. "Integration of Indigenous knowledge in addressing climate change." 2015.

Becklin KM, Anderson JT, Gerhart LM, Wadgyamar SM, Wessinger CA, Ward JK. Examining plant physiological responses to climate change through an evolutionary lens. *Plant physiology*. 2016;172(2):635-49.

Bednar-Friedl, B., Biesbroek, R., Schmidt, D.N., Alexander, P., Børsheim, K.Y., Carnicer, J., Georgopoulou, E., Haasnoot, M., Le Cozzanet, G., Lionello, P. and Lipka, O., 2022. Europe (Chapter 13).

Birkenholtz, Trevor. "Assessing India's drip-irrigation boom: efficiency, climate change and groundwater policy." *Groundwater and Climate Change*. Routledge, 2018. 23-37.

Bisht N, Tiwari S, Singh PC, Niranjana A, Chauhan PS. A multifaceted rhizobacterium *Paenibacillus lentimorbus* alleviates nutrient deficiency-induced stress in *Cicer arietinum* L. *Microbiological research*. 2019;223:110-9.

Bongiovanni R, Lowenberg-DeBoer J. Precision agriculture and sustainability. *Precision agriculture*. 2004;5:359-87.

Bosello, Francesco, and Jian Zhang. "Assessing climate change impacts: agriculture." 2005.

Bouras E, Jarlan L, Khabba S, Er-Raki S, Dezetter A, Sghir F, Trambly Y. Assessing the impact of global climate changes on irrigated wheat yields and water requirements in a semi-arid environment of Morocco. *Scientific reports*. 2019;9(1):19142.

Brown B, Llewellyn R, Nuberg I. Global learnings to inform the local adaptation of conservation agriculture in Eastern and Southern Africa. *Global Food Security*. 2018;17:213-20.

Cairns JE, Sonder K, Zaidi PH, Verhulst N, Mahuku G, Babu R, Nair SK, Das B, Govaerts B, Vinayan MT, Rashid Z. Maize production in a changing climate: impacts, adaptation, and mitigation strategies. *Advances in agronomy*. 2012;114:1-58.

Cao L, Bala G, Caldeira K, Nemani R, Ban-Weiss G. Importance of carbon dioxide physiological forcing to future climate change. *Proceedings of the National Academy of Sciences*. 2010;107(21):9513-8.

Challinor AJ, Watson J, Lobell DB, Howden SM, Smith DR, Chhetri N. A meta-analysis of crop yield under climate change and adaptation. *Nature climate change*. 2014;4(4):287-91.

Chauhan BS, Mahajan G, Randhawa RK, Singh H, Kang MS. Global warming and its possible impact on agriculture in India. *Advances in agronomy*. 2014;123:65-121.

Chhogyell N, Pradhan N, Ghimiray M, Bajgai Y. Evaluation of short duration rice (*Oryza sativa*) varieties as a strategy to cope with climate change. Proc Bhutan Ecol Soc. 2016;1:91-103.

Cline WR. Global warming and agriculture: Impact estimates by country. Peterson Institute; 2007.

Dagar JC, Singh AK, Singh R, Arunachalam AA. Climate change vis-a-vis Indian agriculture. Annals of Agricultural Research. 2012;33(4).

DaMatta FM, Grandis A, Arenque BC, Buckeridge MS. Impacts of climate changes on crop physiology and food quality. Food Research International. 2010;43(7):1814-23.

Demirel U, Morris WL, Ducreux LJ, Yavuz C, Asim A, Tindas I, Campbell R, Morris JA, Verrall SR, Hedley PE, Gokce ZN. Physiological, biochemical, and transcriptional responses to single and combined abiotic stress in stress-tolerant and stress-sensitive potato genotypes. Frontiers in plant science. 2020;11:169.

Dixon, Geoffrey R. "The impact of climate and global change on crop production." *Climate Change*. Elsevier, 2009. 307-324.

Erenstein O, Sayre K, Wall P, Hellin J, Dixon J. Conservation agriculture in maize- and wheat-based systems in the (sub) tropics: lessons from adaptation initiatives in South Asia, Mexico, and Southern Africa. Journal of sustainable agriculture. 2012;36(2):180-206.

Esim N, Atici O. Nitric oxide improves chilling tolerance of maize by affecting apoplasmic antioxidative enzymes in leaves. Plant Growth Regulation. 2014;72:29-38.

Fernando N, Manalil S, Florentine SK, Chauhan BS, Seneweera S. Glyphosate resistance of C3 and C4 weeds under rising atmospheric CO<sub>2</sub>. Frontiers in plant science. 2016;7:190495.

Gholipoor, M., and S. Shahsavani. "Simulation study of past climate change effect on chickpea phenology at different sowing dates in Gorgan, Iran." 2008.

Gray SB and Brady SM. Plant developmental responses to climate change. Developmental biology. 2016;419(1):64-77.

Hadgu G, Tesfaye K and Mamo G. Analysis of climate change in Northern Ethiopia: implications for agricultural production. Theoretical and applied climatology. 2015;121:733-47.

Hatfield, J.L. and Prueger, J.H. Temperature extremes: Effect on plant growth and development. *Weather and climate extremes*. 2015;10,4-10..

Hatfield JL, Boote KJ, Kimball BA, Ziska LH, Izaurralde RC, Ort D, Thomson AM, Wolfe D. Climate impacts on agriculture: implications for crop production. Agronomy journal. 2011;103(2):351-70.

Hijioka Y, Lin E, Pereira JJ, Corlett RT, Cui X, Insarov G, Lasco R, Lindgren E, Surjan A, Aizen EM, Aizen VB. IPCC Fifth Assessment Report: Chapter 24 Asia. Climate Change 2014-Impacts, Adaptation and Vulnerability: Part B: Regional Aspects Working Group II Contribution to the IPCC Fifth Assessment Report. 2014:1327-70.

IPCC. 2013. Summary for policymakers. In: Stocker TF, Qin D, Plattner GK, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM (eds) Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, pp 3–29

IPCC. 2022. “Climate change 2022: Impacts, adaptation and vulnerability”. In: Pörtner, H.-O., Roberts, D.C., Tignor, M., Poloczanska, E.S., Mintenbeck, K., Alegría, A., Craig, M., Langsdorf, S., Löschke, S., Möller, V., Okem, A. and Rama, B (Eds.), Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on climate change, Cambridge University Press. Cambridge University Press, Cambridge, UK and New York, NY, USA, 3056 pp.

IPCC. 2001. Climate Change 2001: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Third assessment report of the Intergovernmental Panel on Climate Change.

Iqbal MS, Singh AK, Ansari MI. Effect of drought stress on crop production. New frontiers in stress management for durable agriculture. 2020:35-47.

Izaurrealde RC, Rosenberg NJ, Brown RA, Thomson AM. Integrated assessment of Hadley Center (HadCM2) climate-change impacts on agricultural productivity and irrigation water supply in the conterminous United States: Part II. Regional agricultural production in 2030 and 2095. Agricultural and Forest Meteorology. 2003;117(1-2):97-122.

Jalil SU, Ansari MI. Physiological role of Gamma-aminobutyric acid in salt stress tolerance. Salt and Drought Stress Tolerance in Plants: Signaling Networks and Adaptive Mechanisms. 2020:337-50.

Jan SA, Shinwari ZK, Rabbani MA. Morpho-biochemical evaluation of Brassica rapa sub-species for salt tolerance. Genetika. 2016;48(1):323-38.

Jones JW, Antle JM, Basso B, Boote KJ, Conant RT, Foster I, Godfray HC, Herrero M, Howitt RE, Janssen S, Keating BA. Brief history of agricultural systems modeling. Agricultural systems. 2017;155:240-54.

Joshi NL, Kar A. Contingency crop planning for dryland areas in relation to climate change. Indian Journal of Agronomy. 2009;54(2):237-43.

Kang Y, Khan S, Ma X. Climate change impacts on crop yield, crop water productivity and food security—A review. Progress in natural Science. 2009;19(12):1665-74.

- Karimi V, Karami E, Keshavarz M. Climate change and agriculture: Impacts and adaptive responses in Iran. *Journal of Integrative Agriculture*. 2018;17(1):1-5.
- Keating BA, Carberry PS, Bindraban PS, Asseng S, Meinke H, Dixon J. Eco-efficient agriculture: Concepts, challenges, and opportunities. *Crop science*. 2010;50:S-109.
- Kirschbaum MU. Direct and indirect climate change effects on photosynthesis and transpiration. *Plant Biology*. 2004;6(03):242-53.
- Knox J, Hess T, Daccache A, Wheeler T. Climate change impacts on crop productivity in Africa and South Asia. *Environmental research letters*. 2012;7(3):034032.
- Kumar R, Gautam HR. Climate change and its impact on agricultural productivity in India. *Journal of Climatology & Weather Forecasting*. 2014;2(1):1-3.
- Kumar V, Khare T, Sharma M, Wani SH. ROS-induced signaling and gene expression in crops under salinity stress. *Reactive oxygen species and antioxidant systems in plants: role and regulation under abiotic stress*. 2017:159-84.
- Lal M, Nozawa T, Emori S, Harasawa H, Takahashi K, Kimoto M, Abe-Ouchi A, Nakajima T, Takemura TU, Numaguti A. Future climate change: Implications for Indian summer monsoon and its variability. *Current science*. 2001:1196-207.
- Lal R, Delgado JA, Groffman PM, Millar N, Dell C, Rotz A. Management to mitigate and adapt to climate change. *Journal of Soil and Water Conservation*. 2011;66(4):276-85.
- Lipper L, Thornton P, Campbell BM, Baedeker T, Braimoh A, Bwalya M, Caron P, Cattaneo A, Garrity D, Henry K, Hottle R. Climate-smart agriculture for food security. *Nature climate change*. 2014;4(12):1068-72.
- Lobell DB, Schlenker W, Costa-Roberts J. Climate trends and global crop production since 1980. *Science*. 2011;333(6042):616-20.
- Lu Y, Chadwick D, Norse D, Powlson D, Shi W. Sustainable intensification of China's agriculture: the key role of nutrient management and climate change mitigation and adaptation. *Agriculture, Ecosystems & Environment*. 2015;209:1-4.
- Lybbert TJ, Sumner DA. Agricultural technologies for climate change in developing countries: Policy options for innovation and technology diffusion. *Food policy*. 2012;37(1):114-23.
- Mahato A. Climate change and its impact on agriculture. *International journal of scientific and research publications*. 2014;4(4):1-6.
- Malhi GS, Kaur M, Kaushik P. Impact of climate change on agriculture and its mitigation strategies: A review. *Sustainability*. 2021;13(3):1318.

Mall RK, Singh R, Gupta A, Srinivasan G, Rathore LS. Impact of climate change on Indian agriculture: a review. *Climatic change*. 2006;78:445-78.

McCarthy JJ, editor. *Climate change 2001: impacts, adaptation, and vulnerability: contribution of Working Group II to the third assessment report of the Intergovernmental Panel on Climate Change*. Cambridge University Press; 2001.

McCormack ML, Adams TS, Smithwick EA, Eissenstat DM. Variability in root production, phenology, and turnover rate among 12 temperate tree species. *Ecology*. 2014;95(8):2224-35.

Megha S, Basu U, Kav NN. Regulation of low temperature stress in plants by microRNAs. *Plant, Cell & Environment*. 2018;41(1):1-5.

Mendelsohn, R. The impact of climate change on agriculture in developing countries. *J. Nat. Res. Policy Res*. 2009, 1, 5–19.

Nations, U. World population prospects: The 2015 revision. *United Nations Econ Soc Aff*, 2015;33(2),1-66.

Neenu S, Biswas AK, Rao AS. Impact of climatic factors on crop production-a review. *Agricultural Reviews*. 2013;34(2):97-106.

Niang I, Ruppel OC, Abdrabo MA, Essel A, Lennard C, Padgham J, Urquhart P. *Climate change 2014: impacts, adaptation, and vulnerability. Part B: regional aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. 2017:1199-265.

Nievola CC, Carvalho CP, Carvalho V, Rodrigues E. Rapid responses of plants to temperature changes. *Temperature*. 2017;4(4):371-405.

Noshi M, Hatanaka R, Tanabe N, Terai Y, Maruta T, Shigeoka S. Redox regulation of ascorbate and glutathione by a chloroplastic dehydroascorbate reductase is required for high-light stress tolerance in *Arabidopsis*. *Bioscience, biotechnology, and biochemistry*. 2016;80(5):870-7.

Nowsherwan I, Shabbir G, Malik SI, Ilyas M, Cheema NM. Selection of wheat genotype (S) for drought stress based on physiological traits. *Int. J. Plant Soil Sci*. 2017;17(3):1-7.

Pagnani G, Galieni A, D'Egidio S, Visioli G, Stagnari F, Pisante M. Effect of soil tillage and crop sequence on grain yield and quality of durum wheat in Mediterranean areas. *Agronomy*. 2019;9(9):488.

Pereira A. Plant abiotic stress challenges from the changing environment. *Frontiers in plant science*. 2016;7:218865.

Porter, John R., Liyong Xie, Andrew J. Challinor, Kevern Cochrane, S. Mark Howden, Muhammad Mohsin Iqbal, David B. Lobell, and Maria Travasso. "Food security and food production systems." 2014.

- Powlson DS, Stirling CM, Jat ML, Gerard BG, Palm CA, Sanchez PA, Cassman KG. Limited potential of no-till agriculture for climate change mitigation. *Nature climate change*. 2014;4(8):678-83.
- Prabhakar SV, Shaw R. Climate change adaptation implications for drought risk mitigation: a perspective for India. *Climatic Change*. 2008;88(2):113-30.
- Prior SA, Runion GB, Marble SC, Rogers HH, Gilliam CH, Torbert HA. A review of elevated atmospheric CO<sub>2</sub> effects on plant growth and water relations: implications for horticulture. *HortScience*. 2011;46(2):158-62.
- Rahman A, Hossain MS, Mahmud JA, Nahar K, Hasanuzzaman M, Fujita M. Manganese-induced salt stress tolerance in rice seedlings: regulation of ion homeostasis, antioxidant defense and glyoxalase systems. *Physiology and molecular biology of plants*. 2016;22:291-306.
- Reyer CP, Leuzinger S, Rammig A, Wolf A, Bartholomeus RP, Bonfante A, De Lorenzi F, Dury M, Gloning P, Abou Jaoudé R, Klein T. A plant's perspective of extremes: terrestrial plant responses to changing climatic variability. *Global change biology*. 2013;19(1):75-89.
- Reynolds M, Foulkes J, Furbank R, Griffiths S, King J, Murchie E, Parry M, Slafer G. Achieving yield gains in wheat. *Plant, cell & environment*. 2012;35(10):1799-823.
- Richards RA. Physiological traits used in the breeding of new cultivars for water-scarce environments. *Agricultural water management*. 2006;80(1-3):197-211.
- Roy SS, Balling Jr RC. Analysis of trends in maximum and minimum temperature, diurnal temperature range, and cloud cover over India. *Geophysical Research Letters*. 2005;32(12).
- Sage RF, Way DA, Kubien DS. Rubisco, Rubisco activase, and global climate change. *Journal of experimental botany*. 2008;59(7):1581-95.
- Saleem MH, Fahad S, Khan SU, Din M, Ullah A, Sabagh AE, Hossain A, Llanes A, Liu L. Copper-induced oxidative stress, initiation of antioxidants and phytoremediation potential of flax (*Linum usitatissimum* L.) seedlings grown under the mixing of two different soils of China. *Environmental Science and Pollution Research*. 2020;27:5211-21.
- Salehi-Lisar SY, Bakhshayeshan-Agdam H. Drought stress in plants: causes, consequences, and tolerance. *Drought stress tolerance in plants, Vol 1: physiology and biochemistry*. 2016:1-6.
- Sarkar J, Chakraborty B, Chakraborty U. Plant growth promoting rhizobacteria protect wheat plants against temperature stress through antioxidant signalling and reducing chloroplast and membrane injury. *Journal of plant growth regulation*. 2018;37(4):1396-412.

Semenza JC, Ploubidis GB, George LA. Climate change and climate variability: personal motivation for adaptation and mitigation. *Environmental Health*. 2011;10:1-2.

Shahmohamadloo RS, Febria CM, Fraser ED, Sibley PK. The sustainable agriculture imperative: A perspective on the need for an agrosystem approach to meet the United Nations Sustainable Development Goals by 2030. *Integrated Environmental Assessment and Management*. 2022;18(5):1199-205.

Smit B, Skinner MW. Adaptation options in agriculture to climate change: a typology. *Mitigation and adaptation strategies for global change*. 2002;7(1):85-114.

Stevanović M, Popp A, Lotze-Campen H, Dietrich JP, Müller C, Bonsch M, Schmitz C, Bodirsky BL, Humpenöder F, Weindl I. The impact of high-end climate change on agricultural welfare. *Science advances*. 2016;2(8):e1501452.

Tesfaye K, Aggarwal PK, Mequanint F, Shirsath PB, Stirling CM, Khatri-Chhetri A, Rahut DB. Climate variability and change in Bihar, India: Challenges and opportunities for sustainable crop production. *Sustainability*. 2017;9(11):1998.

Thornton PK, Ericksen PJ, Herrero M, Challinor AJ. Climate variability and vulnerability to climate change: a review. *Global change biology*. 2014;20(11):3313-28.

Trajkovic S, Kolakovic S. Wind-adjusted Turc equation for estimating reference evapotranspiration at humid European locations. *Hydrology research*. 2009;40(1):45-52.

Trisos, C.H., Adelekan, I.O., Totin, E., Ayanlade, A., Efitre, J., Gameda, A., Kalaba, K., Lennard, C., Masao, C., Mgaya, Y., Ngaruiya, G., Olago, D., Simpson, N. P. and Zakieldean, S. (2022). "Africa". In: Pörtner, H.-O., Roberts, D.C., Tignor, M., Poloczanska, E.S., Mintenbeck, K., Alegría, A., Craig, M., Langsdorf, S., Löschke, S., Möller, V., Okem, A. and Rama, B (Eds.), *Climate change 2022: Impacts, adaptation and vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on climate change*. Cambridge University Press, Cambridge, UK and New York, NY, USA, 1285– 1455.

Ueda Y, Uehara N, Sasaki H, Kobayashi K, Yamakawa T. Impacts of acute ozone stress on superoxide dismutase (SOD) expression and reactive oxygen species (ROS) formation in rice leaves. *Plant Physiology and Biochemistry*. 2013;70:396-402.

Ullah A, Bano A, Khan N. Climate change and salinity effects on crops and chemical communication between plants and plant growth-promoting microorganisms under stress. *Frontiers in Sustainable Food Systems*. 2021;5:618092.

Valizadeh J, Ziaei SM, Mazlounzadeh SM. Assessing climate change impacts on wheat production (a case study). *Journal of the Saudi society of agricultural sciences*. 2014;13(2):107-15.

van Vuuren DP, Isaac M, Kundzewicz ZW, Arnell N, Barker T, Criqui P, Berkhout F, Hilderink H, Hinkel J, Hof A, Kitous A. The use of scenarios as the basis for

combined assessment of climate change mitigation and adaptation. *Global Environmental Change*. 2011;21(2):575-91.

Venkateswarlu B, Shanker AK. Climate change and agriculture: adaptation and mitigation strategies. *Indian Journal of Agronomy*. 2009;54(2):226-30.

White JW, McMaster GS, Edmeades GO. Genomics and crop response to global change: what have we learned?. *Field crops research*. 2004;90(1):165-9.

Yamasoe MA, Von Randow C, Manzi AO, Schafer JS, Eck TF, Holben BN. Effect of smoke and clouds on the transmissivity of photosynthetically active radiation inside the canopy. *Atmospheric chemistry and Physics*. 2006;6(6):1645-56.

You J, Chan Z. ROS regulation during abiotic stress responses in crop plants. *Frontiers in plant science*. 2015;6:165102.

Zargar SM, Gupta N, Nazir M, Mahajan R, Malik FA, Sofi NR, Shikari AB, Salgotra RK. Impact of drought on photosynthesis: Molecular perspective. *Plant gene*. 2017;11:154-9.

Zilberman D, Zhao J, Heiman A. Adoption versus adaptation, with emphasis on climate change. *Annu. Rev. Resour. Econ.* 2012;4(1):27-53.