

## Effect of sowing dates on performance of different mustard varieties (*Brassica juncea* L.) in western Uttar Pradesh

### Abstract

An experiment was conducted to study the “Effect of sowing dates on performance of different mustard varieties (*Brassica juncea* L.) in western Uttar Pradesh” during *rabi* season of 2021-22 at Crop Research Centre of Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut (U.P.). The experiment consisted of 12 treatment combinations with Date of sowing and Mustard varieties. The treatment consisted of three dates of sowing and four mustard varieties were tested in RCBD (Factorial) with three replications. Results revealed that the sowing of mustard on D<sub>1</sub> (7<sup>th</sup> October) gave higher plant height (190.2 cm), LAI (4.06), branches plant<sup>-1</sup>, dry matter accumulation (69.40 g plant<sup>-1</sup>), seed (21.11 q ha<sup>-1</sup>), stover (71.76 q ha<sup>-1</sup>) and biological yield (92.88 q ha<sup>-1</sup>) of mustard crop. Similarly, the variety V<sub>1</sub> (Pusa Vijay) gave highest plant height (197.5 cm), LAI (4.22), branches plant<sup>-1</sup>, dry matter accumulation (69.65 g plant<sup>-1</sup>), seed (22.27 q ha<sup>-1</sup>), stover (69.59 q ha<sup>-1</sup>) and biological yield (91.87 q ha<sup>-1</sup>) of mustard crop. Thus, it may be concluded that the Pusa Vijay variety of mustard sown on 7<sup>th</sup> October gave significantly higher values of growth, yield attributes and yield of mustard.

### **Introduction**

Indian mustard (*Brassica juncea* L.) is a winter season crop grown in different agro-climatic conditions, ranging from northern hills to down south under irrigated and rainfed conditions under vast range of temporal dimensions, namely early, timely and late-sown under irrigated and rainfed conditions. It is a member of the *Cruciferae* family and good source of high-quality edible oil (33-49 %). After groundnut (*Arachis hypogea* L.) oil and soybean (*Glycine max* L.) oil, mustard is the third most significant oilseed crop in the world. The world's tropical and subtropical regions are where it is grown. Globally, the total area under Rapeseed-Mustard crop is 35.95 Mha with 71.49 MT production and 1990 kg ha<sup>-1</sup> productivity. However, India continues to be rank second after Canada in acreage 6.86 Mha (19.81%) and rank fourth after Canada, European Union and China in production 9.12 MT (10.37%) with productivity 1331 kg ha<sup>-1</sup>. In India, Rajasthan have rank first with 2.84 Mha (41.44%) area and 4.10 MT (45.03%) production. Uttar Pradesh contributed 0.77 Mha (11.29%) to the total area and 0.98 MT (10.79%) to the production with 1090 kg ha<sup>-1</sup> productivity. **Anonymous (2021).**

Oilseeds cultivation is undertaken across the country in about 26 million ha, covering 72% under rain fed areas and producing around 30 million tons of oil seed. Out of the nine major oilseed crops grown in India, soybean (39%), Groundnut (26%) and Mustard (24%) add more than 88% of total oilseed production in the country. It has a protein content of 17–25%, a fiber content of 8-10%, a moisture content of 6-10%, and an extractable content of 10–12%. There is a significant level of erucic acid in mustard oil (38-57%). However, poisonous glucosinolates make mustard cake, which is also contained in manure and bovine feed, unsuitable as a source of human protein. The period of sowing seems crucial for crop production since different sowing dates offer varying climatic conditions for crop growth and development in the same location. Because mustard is a thermos-sensitive crop, the delayed seeding reduced grain yield by synchronizing the silique filling period with high temperatures, reducing assimilates production, increasing the rate of drought stress, and accelerating plant maturity. These crops are cultivated in India during the *rabi* season, which runs from September to October through February to March. The genotypes of Indian mustard, however, might vary its output capacity. It is a reality that certain genotypes do not always display the same phenotypic traits depending on the environment. The choice of genotypes for sustaining the higher yield ultimately depends on the genotypes' growth responses to the environment and their relative ranking. One of the main input variables influencing the development and production of crops is time. It has an impact on the mustard plants vegetative, reproductive, and maturation periods. Because agro-ecological circumstances can vary from one place to another and from one variety to another, the best time to grow mustard may differ (**Sarkar *et al.*, 2004**).

Selection of suitable sowing time with a suited varieties is equally important. Different sowing dates give various climatic conditions within the same area for crop growth and yield stability. Sowing at the correct time allows a crop to grow and develop sufficiently to provide a reasonable amount (**Pandey *et al.*, 1981**). Late sowing of mustard reduces duration due to high temperatures during the reproductive phase, which also results in a decrease in yield (**Kumari *et al.*, 2004**). Growth characters, yield attributes and yield of mustard were significantly influenced by 4 different sowing dates. Higher plant height, length of silique, seed/silique, and seed and stover yields were obtained when the crop was sown on 15 October than when it was sown on 30 October (**Awasthi *et al.*, 2007**).

Some studies showed that mustard crops seeded in the second week of September produced considerably more yield than those sown in the first fortnight of October (**Iraddi, 2008**). Generally, it was found that mustard crops planted after October 30th produced lesser yields (**Panwar et al., 2000; Singh et al., 2002a; Sonaniet al., 2002 and Panda et al., 2004b**).**Bharat et al. (2022)** recoded that sowing of mustard (15 October) significantly higher seed yield (1.30 t/ha) than 25 October (10.29 q/ha) and 4 November (0.86 t/ha). Similarly, during *winter* season of 2015-16, again statistically higher seed yield was recorded by the crop sown on 15 October (1.32t/ha) and was significantly higher than when crop was sown on 25 October (0.96 t/ha) and 4 November (0.80 t/ha).**Awasthi et al. (2007)** reported that various important growth characters and seed yield of mustard were significantly influenced by different varieties. They found that the Variety ‘Vaibhav’ excelled over ‘Urwash’ in terms of plant height, number of siliquae, length of siliqua, seeds/siliqua, seed yield and stover yield.

### **Material and methods**

The experiment was carried out at Crop Research Centre, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut (U.P.) located in Indo-Gangnetic plains of Western Uttar Pradesh at 29 13’ 96” N latitude and University is situated at 77 68’ 43” E longitudes with an elevation of 228 meters above the mean sea level. Meerut lies on national highway 58 and is at a distance of 70 km from Delhi. Meerut predominantly enjoys semi-arid and sub-tropical climate with extremely hot summer and cold winter. Minimum and maximum temperatures both exhibit a gradual decrease starting from first week of October and reach their minimum during December and January. An increase in the temperature is recorded with effect from first week of February and peak values noticed during end of May. Occasional frost is also experienced during second fortnight of December and first fortnight of January. The mean weekly minimum temperature records as low as 4.89<sup>0</sup>C in last week of December, whereas mean weekly maximum temperature reached 38.7<sup>0</sup> C in April. The area receives mean annual rainfall of 845 mm, of which more than 80% is in the month of July-September through South west monsoon. A few winter showers are also received. April and May are the driest months with mean relative humidity of 50 to 55%, Whereas the recorded of high humidity 92 % is in the month of August. Mean weekly temperature, relative humidity, sunshine hours, evaporation and rainfall as recorded at nearby located meteorological observatory of Indian Institute of Farming Systems Research, Meerut. The maximum and minimum temperatures recorded were 38.70°C and 4.7°C during the crop growth period. The

upward trend was recorded in temperature, as the crop approached towards maturity. Maximum temperature ranged from 16.6°C to 38.7°C during maturity phase of the crop. Relative humidity varied from 28.4% to 92.6% during crop growth period.

The experiment comprised of 12 treatment combinations of three dates of sowing i.e., 7<sup>th</sup> October, 19<sup>th</sup> October, 31<sup>st</sup>October and four Mustard varieties i.e., Pusa Vijay, RH-0749, Pioneer-45S46 and Pro-Agro-5222 were tested in RCBD (Factorial) with three replications. The height of the mustard plant was measured from the base to the top of the plant at harvesting, with the help of meter scale. The mean of plant height was worked out on the basis of total height of five randomly selected plants in each plot which was divided by the number of plants. Number of primary and secondary branches per plant were counted from five plants selected randomly at harvest and their mean value was taken. Five plants were randomly selected from border rows at harvest stage and after sun drying, materials were dried in oven (65 °C ±1) till the constant weight was attained. The average value obtained was recorded as the dry matter of the plant (g plant<sup>-1</sup>). The samples collected for dry matter estimation, leaves of five plants was packed at 90 DAS and leaf area was measured with the help of leaf area meter.

$$\text{Leaf Area Index} = \frac{\text{Leaf area (cm}^2\text{)}}{\text{Ground area (cm}^2\text{)}}$$

Number of primary and secondary branches per plant were counted from five plants selected randomly at 60, 90 DAS and at harvest and their mean value was taken. Similarly, five plants were randomly selected from border rows at 30, 60, 90 DAS and at harvest stage and after sun drying, materials were dried in oven (65±1 °C) till the constant weight was attained. The average value was recorded as the dry matter of the plant (g plant<sup>-1</sup>). Mean crop growth rate (g m<sup>-2</sup> day<sup>-1</sup>) of a plant for a time "t" is defined as the increase in dry weight of plant material from a unit area per unit of time. It was calculated with the following formula **Radford(1967)** from periodic dry matter recorded at different stages:

$$\text{Crop growth rate} = \frac{W_2 - W_1}{t_2 - t_1} \times \frac{1}{A}$$

Where,

$W_1$  = Total dry weight of plant at time  $t_1$  (g),  $W_2$  = Total dry weight of plant at time  $t_2$  (g),  $T_1$  = Time at first observation (days),  $T_2$  = Time at second observation (days),  $A$  = represents the ground area ( $m^2$ ).

From the individual plot, the net plot area was harvested separately and produce was sun dried. After sun drying, the crop was threshed and produce cleaned. The final weight was recorded in  $kg\ plot^{-1}$  and finally converted into  $q\ ha^{-1}$ . Stover yield of mustard was calculated with subtraction of seed yield from biological yield and reported in  $q\ ha^{-1}$ . After 3-4 days sun drying, all above the ground plant parts of the net plot were dried and weighed in  $kg\ plot^{-1}$  to represent the biological yield and finally converted into  $q\ ha^{-1}$ . Harvest index is an economic yield expressed as percentage of biological yield and calculated as formula given by **Donald and Hamblin (1976)**

$$HI (\%) = \frac{\text{Economic yield (q/ha)}}{\text{Biological yield (q/ha)}} \times 100$$

Mustard seeds were sown in line at the distance of 45 cm row to row and plant to plant 15 cm with the help of Kudal. The seed rate was used @5  $kg\ ha^{-1}$  and fertilizer nitrogen, phosphorus and potassium were applied in the forms of Urea, DAP and Muriate of potash @120:60:60:40  $kg\ ha^{-1}$ , respectively. Full dose of phosphorus, potassium and half dose of nitrogen were applied as basal dressing at the time of sowing and rest half dose of nitrogen was applied as two split doses at the time of first irrigation and second irrigation.

## **Result and discussion**

### **Growth and yield attributes**

Growth and various yield attributes of mustard were significantly influenced by different sowing dates and varieties (Table 1). 7<sup>th</sup> October sown crop resulted in significantly higher plant height, number of primary and secondary branches at harvest over 19<sup>th</sup> October and 31<sup>st</sup> October. Plant height in 7<sup>th</sup> October sown crop was significantly higher (190.2 cm) over 19<sup>th</sup> October (180.9 cm) and 31<sup>st</sup> October (172.1 cm). Similarly, Plant height was significantly influenced by varieties. Among the varieties, 'Pusa Vijay' recorded the highest plant height (197.5 cm) which was significantly superior to RH-0749(182.2 cm) and Pro-Agro-5222 (154.5 cm) but at par with Pioneer-45S46 (189.9 cm). Similar findings have also been reported by **Chand *et al.* (2000), Jadhav and Singh. (1992), Yadav *et al.* (1994), Gare**

*et al.* (1996) and Surekha and Reddy. (1996) which supports the findings of the present investigation.

The highest number of primary branches plant<sup>-1</sup> (9.80) was produced in 7<sup>th</sup> October sowing which got significantly reduced with delay in sowing i.e., 19<sup>th</sup> October (7.88) and 31<sup>st</sup> October (6.69). Similarly, the highest number of secondary branches plant<sup>-1</sup> (15.76) was recorded in 7<sup>th</sup> October sown crop which got reduced by 14.36 in 19<sup>th</sup> October and 11.60 in 31<sup>st</sup> October sowing. Similar results were also obtained by **Kumari *et al.* (2012)**, **Kumar *et al.* (2003)**. In case of varieties the highest number of primary and secondary branches plant<sup>-1</sup> was recorded in Pusa Vijay (9.63, 15.39) followed by Pioneer-45S46 (8.52, 14.39), RH-0749 (7.73, 13.61) and Pro-Agro-5222 (6.62, 123.24), respectively which might be due to genotype response and capability of higher dry matter production. Similar results were also obtained by **Awasthi *et al.* (2007)**, **Singh and Singh (2002)**.

Higher dry matter accumulation (69.40 g plant<sup>-1</sup>) was recorded under 7<sup>th</sup> October sowing over 19<sup>th</sup> October (66.05 g plant<sup>-1</sup>) and 31<sup>st</sup> October (63.53 g plant<sup>-1</sup>). Accumulation of dry matter in the plant is directly related to plant height, leaf area index and number of branches plant<sup>-1</sup> which were appreciably reduced as sowing delayed. Similar findings have been reported by **Singh and Singh (2002)** and **Lallu *et al.* (2010)**. Among the varieties Pusa Vijay recorded significantly higher dry matter accumulation (69.65 g plant<sup>-1</sup>) over RH-0749 (65.29 g plant<sup>-1</sup>), Pro-Agro-5222 (63.41 g plant<sup>-1</sup>) but *at par* with Pioneer-45S46 (66.96 g plant<sup>-1</sup>) and its probable reason might be attributed to genetic characters of Pusa Vijay which has higher capacity to utilize the photosynthates more efficiently for maximum leaf area index, number of primary and secondary branches plant<sup>-1</sup> and ultimately the dry matter accumulation. The increasing of yield per unit area could be due to efficient utilization of thermal energy by mustard which improving the early sowing and adoption of optimum plant population, whereas sufficient spacing recorded maximum utilization of photo and thermal units by the individual mustard plants thereby leading to significantly higher dry matter accumulation, yield attributes, yield and harvest index, but lower productivity due to lower plant population. Similar findings have been reported by **Panda *et al.* (2004)**, **Patel *et al.* (1994)**.

Crop sown on 7<sup>th</sup> October recorded significantly higher leaf area index (4.06) over 19<sup>th</sup> October (3.51) and 31<sup>st</sup> October (2.95) Similar trends were recorded at successive stages.

Delayed, sowing by 36 days achieved lower leaf area index (2.95) at all the stages of mustard crop which might be due to less vegetative growth because of less favourable environmental conditions when crop was sown too late sowing conditions. Different varieties have no significant influence on the leaf area index. Among the varieties Pusa Vijay recorded maximum leaf area index (4.22) as compared to RH-0749 (3.32), Pro-Agro-5222 (2.82) and Pioneer-45S46 (3.65), respectively. Similar results were also obtained by **Bhuiyan et al. (2008)**, **Tyagi (1994)** and **Kumar et al. (2008)**.

Crop growth rate ( $\text{g m}^{-2} \text{ day}^{-1}$ ) was significantly influenced by various sowing dates and varieties. CGR increased with the advancement of crop age and reached to maximum between 60-90 DAS and thereafter decreased sharply. In case of different dates of sowing, the maximum crop growth rate ( $4.26 \text{ g m}^{-2} \text{ day}^{-1}$ ) was recorded in mustard sown on 7<sup>th</sup> October over 19<sup>th</sup> October ( $4.22 \text{ g m}^{-2} \text{ day}^{-1}$ ) and minimum crop growth rate ( $4.17 \text{ g m}^{-2} \text{ day}^{-1}$ ) was recorded in 31<sup>st</sup> October date of sowing. Similar findings have been reported by **Lallu et al. (2010)**. Similarly, in case of different varieties, the maximum crop growth rate ( $4.26 \text{ g m}^{-2} \text{ day}^{-1}$ ) was recorded in variety Pusa Vijay over RH-0749 ( $4.21 \text{ g m}^{-2} \text{ day}^{-1}$ ), Pro-Agro-5222 ( $4.17 \text{ g m}^{-2} \text{ day}^{-1}$ ) but *statistically at par* with Pioneer-45S46 ( $4.23 \text{ g m}^{-2} \text{ day}^{-1}$ ) among these varieties at crop growth stages. Similar results were also obtained by **Tyagi (1994)** and **Kumar et al. (2008)**.

**Table 1. Effect of different date of sowing and varieties on plant height (cm), primary branches plant<sup>-1</sup>, secondary branches plant<sup>-1</sup>, dry matter accumulation ( $\text{g plant}^{-1}$ ), leaf area index and crop growth rate ( $\text{g m}^{-2} \text{ day}^{-1}$ ) of mustard at different growth stages**

Treatments		Plant height (cm)	Primary branches plant <sup>-1</sup>	Secondary branches plant <sup>-1</sup>	DMA ( $\text{g plant}^{-1}$ )	LAI	CGR ( $\text{g m}^{-2} \text{ day}^{-1}$ )
<b>Factor A (Date of sowing)</b>							
D1	7 <sup>th</sup> October	190.2	9.80	15.76	69.40	4.06	4.26
D2	19 <sup>th</sup> October	180.9	7.88	14.36	66.05	3.51	4.22
D3	31 <sup>st</sup> October	172.1	6.69	11.60	63.53	2.95	4.17
<i>SEM ±</i>		4.16	0.17	0.27	1.15	0.06	0.01
<i>CD (P=0.05)</i>		12.27	0.51	0.81	3.39	0.18	0.05
<b>Factor B (Varieties)</b>							
V1	Pusa Vijay	197.5	9.63	15.39	69.65	4.22	4.26

V2	RH-0749	182.2	7.73	13.61	65.29	3.32	4.21
V3	Pioneer-45S46	189.9	8.52	14.39	66.96	3.65	4.23
V4	Pro-Agro-5222	154.5	6.62	12.24	63.41	2.82	4.17
<i>SEm</i> ±		4.80	0.20	0.32	1.32	0.07	0.02
<i>CD(P=0.05)</i>		14.17	0.59	0.94	3.92	0.21	0.05

**Table 2. Effect of different date of sowing and varieties on seed, stover, biological yield ( $q\ ha^{-1}$ ) and harvest index (%) of mustard**

Treatments		Yield( $qha^{-1}$ )			
		Seed yield	Stover yield	Biological yield	H.I. (%)
<b>FactorA(Date of sowing)</b>					
D1	7 <sup>th</sup> October	21.11	71.76	92.88	22.67
D2	19 <sup>th</sup> October	19.40	65.61	85.01	22.51
D3	31 <sup>st</sup> October	17.51	54.29	71.80	24.56
<i>SEm</i> ±		0.96	1.05	1.50	0.89
<i>CD(P=0.05)</i>		2.84	3.11	4.44	NS
<b>FactorB(Varieties)</b>					
V1	Pusa Vijay	22.27	69.59	91.87	24.06
V2	RH-0749	19.65	62.75	82.40	23.81
V3	Pioneer-45S46	21.02	66.33	87.36	24.01
V4	Pro-Agro-5222	14.42	56.87	71.29	21.11
<i>SEm</i> ±		1.11	1.21	1.73	1.02
<i>CD(P=0.05)</i>		3.28	3.60	5.13	NS

### Biological, seed and stover yield

The data reveal that date of sowing and mustard varieties significantly influenced the seed, stover and biological yield (Table 2.) of mustard. Different dates of sowing under investigation brought significant influence on the seed yield. Crop sown on 7<sup>th</sup> October recorded significantly higher seed yield (21.11) over 31<sup>st</sup> October (17.51), while *at par* with

19<sup>th</sup> October (19.40) date of sowing. Seed yield was significantly influenced due to various varieties. Among the varieties Pusa Vijay recorded significantly higher seed yield (22.27) as compared to Pro-Agro-5222, while which was *at par* with Pioneer-45S46 (21.02) and RH-0749 (19.65). Similar findings have been reported by **Mondal *et al.* (2011) and Kumari *et al.* (2012).**

The stover yield was also affected by sowing dates because it is depended on plant growth and dry matter accumulation. Crop sown on 7<sup>th</sup> October recorded significantly higher stover yield (71.76) over 19<sup>th</sup> October (65.61) and lowest 31<sup>st</sup> October (54.29) sowing of mustard. Similarly, among the varieties Pusa Vijay recorded significantly higher stover yield (69.59) as compared to RH-0749 (62.75), Pro-Agro-5222 (56.87) and *at par* with Pioneer-45S46 (66.33). All the growth and yield attributes which determined the seed and stover yield of mustard crop, were adversely influenced when the sowing was done on too early and late sowing, which might be resulted to poor growth and translocation of photosynthates from source to sink and ultimately lower yield was recorded. Similar findings were reported by **Panda *et al.* 2004.**

The biological yield of mustard significantly influenced with the date of sowing and mustard varieties. The highest biological yield (92.88 q ha<sup>-1</sup>) was recorded on the date of sowing 7<sup>th</sup> October, followed by 19<sup>th</sup> October (85.01 q ha<sup>-1</sup>) and the lowest (71.80 q ha<sup>-1</sup>) on 31<sup>st</sup> October. Similarly, the highest value of biological yield (91.87 q ha<sup>-1</sup>) was recorded in variety Pusa Vijay which was *at par* with Pioneer-45S46 (87.36 q ha<sup>-1</sup>) and the lowest biological yield (71.29 q ha<sup>-1</sup>) in variety Pro-Agro-5222. Similar results were also obtained by **Singh *et al.* (2002a), Kumar *et al.* (2008) and Kumari *et al.* (2012).**

#### *Harvest index (%)*

It is clear that with different dates of sowing, the higher value of the harvest index (24.56%) was recorded in 31<sup>st</sup> October because the ratio of seed and stover yield is less as compared to other date of sowing therefore, the harvest index is higher of this date and the lowest (22.51%) in 19<sup>th</sup> October. Similarly, the different varieties of mustard did not bring any significant variation in harvest index. However, the maximum harvest index (24.06%) was recorded in variety Pusa Vijay and the minimum harvest index (21.11%) in variety Pro-Agro-5222. Data analysis revealed that the variety Pusa Vijay had a 1.04, 0.20 and 13.97% higher harvest index than the varieties RH-0749, Pioneer-45S46 and Pro-Agro-5222, respectively.

## Conclusion

Thus, it may be concluded that sowing of mustard on 7th October with the introduction of variety 'Pusa Vijay' at a spacing of 45×10 cm can achieve higher productivity and better economic returns from mustard in Western Uttar Pradesh.

## References

- Anonymous. 2021. Directorate of Rapeseed and Mustard Research, Bharatpur, Rajasthan. *Annual report-2021*.
- Awasthi, U.D., Singh, R.B. and Dubey, S.D. 2007. Effect of sowing date and moisture-conservation practice on growth and yield of Indian mustard (*Brassica juncea*) varieties. *Indian Journal of Agronomy* **52**(2): 151-153.
- Bharat, R., Gupta, M., Gupta, V. and Gupta, R. 2022. Effect of sowing schedules and crop geometry on photo-thermal energy utilization and productivity of Indian mustard (*Brassica juncea*) in Jammu region. *Indian Journal of Agronomy* **67**(3): 287-293.
- Bhuiyan, M.S., Mondol, M.R.I., Rahaman, M.A., Alam, M.S. and Faisal, A.H.M.A. (2008). Yield and yield attributes of rapeseed as influenced by date of planting. *International Journal of Sustainable Crop Production* **3**(3): 25-29.
- Chand, P., Govardhan, M. and Sujatha M. 2000. Effect of dates of sowing on performance of mustard varieties. *Research on Crops***1**:153-155.
- Donald, C.M. and Hamblin, J. 1976. The biological yield and harvest index of cereals as agronomic and plant breeding criteria. *Advances in agronomy* **28**: 361-405.
- Gare, B.N., Dudbade, D.D., Khade, K.K. 1996. Effects of sowing dates on yield and quality of mustard cultivars. *Journal-Maharashtra Agricultural Universities* **2**: 238-239.
- Iraddi, V.S. 2008. Response of Mustard (*Brassica juncea* L.) Varieties to Date of Sowing and Row Spacing in Northern Transition Zone of Karnataka. Thesis University of Agricultural Sciences, Dharwad. *Karnataka Journal Agricultural Science***21**(4): 44-46.

- Jadhav, S.N. and Singh, N.P. 1992. Water use and soil moisture-extraction pattern of mustard (*Brassica juncea*) under varying sowing dates, insect pest-control measures and irrigation. *Indian Journal of Agronomy* **3**: 198-198.
- Kumar, R. 2003. Screening of promising varieties of oilseed *Brassica* at different dates. Ph.D. Thesis, G.B. Pant Univ. of Agriculture and Technology, Pantnagar.
- Kumar, Rajesh, R.P., Singh and Yeshpal 2008. Yield and quality of *Brassica* species as influenced by different dates of sowing and varieties. *Pantnagar Journal Research***6**(1): 6-11.
- Kumari, A. and Singh, R.P. 2012. Productivity, nutrient uptake and economics of mustard hybrid (*Brassica juncea*) under different planting time and row spacing. *Indian journal of Agronomy* **57**(1): 61-67.
- Kumari, R.C., Koteswararao. and Obulamma U. 2004. Impact of sowing dates and land treatments on Indian mustard (*Brassica juncea*) in non-traditional areas of Andhra Pradesh. *Madras Agricultural Journal***91**(7-12): 374-377.
- Lallu, R. S., V.S. Baghel and S.B.L., Srivastava. 2010. Assessment of mustard genotypes for thermos-tolerance at seed development stage. *Indian Journal Plant Physiology* **15**(10): 36-43.
- Mondal, S.S., Acharya, D., Sarkar, S. and Ghosh, A. 2004. Effect of pre-sowing seed treatment on growth, yield, quality and nutrient uptake of Indian mustard (*Brassica juncea*) under rainfed condition. *Indian Journal of Agronomy* **49**(4): 262-263.
- Panda, B. B., Y. S. Shivay and Bandyopadhyay S. K. 2004b. Growth and development of Indian mustard (*Brassica juncea*) under different levels of irrigation and date of sowing. *Indian Journal Plant Physiology* **9**(4): 419-425.
- Panda, B.B., Bandyopadhyay, S.K., and Shivay, Y.S. 2004. Effect of irrigation level, sowing dates and varieties on yield attributes, yield, consumptive water use and water-use efficiency of Indian mustard (*Brassica juncea*).
- Pandey, B.P., S.K. Sirvastava and Lal. R.S. 1981. Genotype and environment interaction in mustard (*Brassica juncea* L.) **8**(3): 14-17.

- Panwar, K. S., Sharma S.K. and Nanwal R.K. 2000. Influence of sowing times on the yield of different mustard cultivars (*Brassica* spp.) under conserved soil moisture conditions. *Indian Journal Agricultural Science* **70**(6): 398-399.
- Patel, K.G., Rathore, S.S., Shiyani, G.L. and Raman, S. 1994. Response of different mustard varieties to dates of sowing and levels of nitrogen in South Gujarat. *Madras Agricultural Journal* **81**(6): 348-349.
- Radford, P.J. 1967. Growth analysis formula their use and abuse. *Crop Science* **5**(7): 171-175.
- Singh, R., V.U.M., Rao, Diwan S. and Suryakant 2002a. Effect of sowing date and plant density on phonological behavior, yield and its attributes in oilseed *Brassica*. *Journal Oilseeds Research* **19**(1): 119-121.
- Singh, S.K. and Singh G. 2002. Response of Indian mustard (*Brassica juncea*) varieties to nitrogen under varying sowing dates in eastern Uttar Pradesh. *Indian Journal Agronomy* **47**(2): 242-248.
- Sonani, V.V., P. T. Patel and G.G. Patel 2002. Performance of mustard under different dates of sowing in Bhal and Coastal Argo-climatic zone of Gujarat. *Journal Oilseeds Research* **19**(1): 122.
- Tyagi, P.K., 1994. Quantification of variability in growth, development and yield of Indian mustard (*Brassica juncea*L.) under different environments. M.Sc. Thesis, CCS HAU, Hisar (Haryana). University **14**(2): 155-160.
- Yadav, R.N., Bhan, S. and Uttam, S.K. 1994. Effect of sowing date, variety and plant population on growth and yield of rainfed Indian mustard (*Brassica juncea*). *Indian Journal of Agronomy* **39**(4): 682-684.