

Effect of Liquid Biofertilizer and Variable Source of Nutrients on Growth and Yield of Indian Mustard (*Brassica juncea L.*) in Western U.P., India

Abstract

A field experiment was conducted during *Rabi* season of 2021-22 at Crop Research Centre of Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut (U.P.). The experiment consisted of twenty treatment combinations with variable source of nutrients and biofertilizer. The treatment consisted of four sources of nutrients and five Biofertilizer levels were tested in RCBD (Factorial) with three replications. Results revealed that the application of 75% RDF (NPKS 90:45:45:30 kg/ha) +2-ton Vermicompost ha⁻¹ gave higher plant height (203.55 cm), LAI (2.929), branches plant⁻¹, dry matter accumulation (66.65 g plant⁻¹), and seed (1854), stover(7193)& biological (9047) yield (kg ha⁻¹) of mustard crop. Similarly, application of Azotobacter + PSB + KMB + ZnSB each @20 mlkg⁻¹ seed treatment gave higher plant height (201.46 cm), LAI (2.702), branches plant⁻¹, dry matter accumulation (64.98 g plant⁻¹), and seed (1735), stover(6888) & biological (8623) yield (kg ha⁻¹) of mustard crop. Thus, it may be concluded that the application of 75%RDF (NPKS 90:45:45:30kg/ha) + 2-ton Vermicompost/ha +Biofertilizers (Azotobacter + PSB + KMB + ZnSB each @ 20 ml/kg Seed treatment)found economical in obtaining higher growth and yield of mustard.

Keywords: Liquid Biofertilizer, NPKS, PSB, KMB, Seed Treatment

INTRODUCTION

Rapeseed and Mustard are the major *rabi* oilseed crops of India. Indian mustard [*Brassica juncea (L.) Czern&Coss*] is commonly known as *rai*. Crop is grown under a wide range of agro-climatic conditions. During the growing season, it requires somewhat cool temperatures with a good supply of soil moisture and a dry harvest period. Mustard is typically grown in temperate areas. It is also grown as a cold season crop in select tropical and subtropical climates. Indian mustard is an important oilseed crop in the world. Rapeseed and mustard are members of the genus *Brassica* and the family *Brassicaceae*. This plant is grown for vegetable, oil, fodder, condiment, and green manure. With the passage of time, this crop has evolved into one of the most important suppliers of vegetable oil. Vegetable oils are a high-value agricultural commodity. Globally, the total area under Rapeseed-Mustard crop is 35.95 Mha with 71.49 MT production and 1990 kg/ha productivity. However, India continues to be rank 2nd after Canada in acreage 6.86 Mha (19.81%) and rank 4th after Canada, European Union and China in production 9.12 MT (10.37%) with productivity 1331 kg/ha. Rapeseed-Mustard crop in India are grown in diverse agro climatic conditions. In India, Rajasthan have rank 1st with 2.84 Mha (41.44%) area and 4.10 MT (45.03%) production. Uttar Pradesh contributed 0.77 Mha (11.29 %) to the total area and 0.98 MT (10.79 %) to the

production with 1090 kg/ha productivity. (Anonymous, 2021). “It is the most important edible oil seed crops of India next to groundnut and soybean. India has 12-15% of the world’s area under oilseed but account for less than 6-7 % of world’s production to meet the need of about 16% of world population” (Anonymous, 2011).

“The oil content of Indian mustard varies between 30 to 49%. These seed and oil are used as a condiment and for flavouring various food items such as curries and pickles. The oil is utilized for human consumption throughout the Northern India, in cooking and frying purposes. The oil cake is used as feed and manure. It is also used as vegetable where the fresh leaves of young plants serve as a good source of Sulphur and minerals in the human diet. Population of India is increasing rapidly and consequently edible oil demand is also going up day by day. Hence, it has become necessary to enhance the present production by developing superior varieties of Indian mustard. Indian mustard requires optimum weather conditions for its good growth and development. Since, it is mostly grown after the harvest of long duration varieties of rice and late recession of moisture from rice fields in Uttar Pradesh. The sowing of mustard crop gets delayed and the growth and vigour of mustard is not good as timely sown crop. It is important to increase the productivity of Indian mustard, which still has a greater scope to exploit the yield potential of existing cultivars with agronomic manipulations. Among the various agronomic factors that are known to augment crop production, fertilizer and nutrient management play a significant role. The efficiency of fertilizer nitrogen is only 40-50%, phosphorus 15-20% and Sulphur 10-12% in Indian soils and this could be enhanced by efficient use of inputs” (Hegde and Sudhakar Basa, 2004).

For sustainable crop production, integrative effect of organic, inorganic and bio-fertilizers is important. Bio-fertilizers and organic manures play a significant role in sustainable agriculture. Bio-fertilizer contains selective strains of nitrogen fixing bacteria, PSB, and potash mobilizing Bacteria which helps to improve availability of NPK to crops. It mobilizes & converts insoluble plant nutrients to soluble and makes it available to plants. Farmyard manure with rich organic matter can be supplemented with N, P & K fertilizers. FYM improves soil structure through binding effect on soil aggregates, cation exchange capacity and water holding capacity, fertilizer use

efficiency, microbial activity and nutrient availability in soil besides supplying the essential plant nutrients. There is a great scope for increasing the production of Indian mustard by bringing more area under cultivation and increasing its productivity by applying organic manures (FYM) & liquid biofertilizers (Azotobacter, PSB, KMB, ZnSB) with balanced fertilization (N:P:K:S) and maintaining soil fertility status. Though some information about mustard nutrition is available but the role of nutrient use efficiency on affecting the productivity of crop under the influence of different organic, inorganic nutrients and biofertilizers needs to be worked out. The suitable treatment of different nutrient with appropriate dosages is to be worked out to understand nutrient uptake, availability and achieve maximum yield.

Material and methods

The experiment was carried out at Crop Research Centre, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut (U.P.) located in Indo – Gangetic plains of Western Uttar Pradesh at 29° 13' 96" N latitude and University is situated at 77° 68' 43" E longitude with an elevation of 228 metres above the mean sea level. Meerut lies on national highway 58 and is at a distance of 70 km from Delhi. The climate of this region is subtropical and semi-arid climate characterized with hot summers and extremely cold winters. The normal period for onset of the monsoon in this region is third week of June and, it lasts up to end of September or sometimes extends to the first week of October. The average annual rainfall is about 1073 mm and out of which about 80 percent is received by south-west monsoon while annual potential evapo-transpiration is about 1667 mm. Thus, the area falls under soil moisture deficit. The rainfall during the experimental period was recorded from the Meteorological observatory of the IIFSR, Modipuram, Meerut. The temperature begins to rise from the month of February and reaches its maximum in March month, minimum and maximum temperature ranged between (4.7 °C to 38.70 °C), respectively, whereas minimum and maximum relative humidity ranged between 28.4 % and 92.6 % percent during crop period. Mean weekly temperature, relative humidity, sunshine hours, evaporation and rainfall as recorded at nearby located meteorological observatory of Indian Institute of Farming Systems Research, Meerut.

The experiment consisted of twenty treatment combinations with variable source of nutrients and biofertilizer. The treatment consisted of four sources of nutrients i.e., Control, 100% RDF (NPKS 120:60:60:40 kg/ha), 75% RDF (NPKS 90:45:45:30 kg/ha) + 6-ton FYM/ha and 75% RDF (NPKS 90:45:45:30 kg/ha) + 2-ton Vermicompost/ha and five Biofertilizer levels i.e., No biofertilizers, Azotobacter @20 ml/kg seed treatment, Azotobacter + PSB each @20 ml/kg seed treatment), Azotobacter + PSB + KMB each @20 ml/kg seed treatment, Azotobacter + PSB + KMB + ZnSB each @20 ml/kg seed treatment were tested in RCBD (Factorial) with three replications. The mustard variety Pusa Vijay (NPJ-93) was grown and growth & yield, nutrient uptake, soil properties as influenced by different treatments were assessed. The height of the mustard plant was measured from the base to the top of the plant at harvesting, with the help of meter scale. The mean of plant height was worked out on the basis of total height of five randomly selected plants in each plot which was divided by the number of plants. Number of primary and secondary branches per plant were counted from five plants selected randomly at harvest and their mean value was taken. Five plants were randomly selected from border rows at harvest stage and after sun drying, materials were dried in oven (65 °C) till the constant weight was attained. The average value obtained was recorded as the dry matter of the plant (g/plant). The samples collected for dry matter estimation, leaves of five plants was packed at 90 DAS and leaf area was measured with the help of leaf area meter.

$$LAI = \frac{\text{Leaf area (cm)}}{\text{Land area (cm)}}$$

Mean crop growth rate of a plant for a time "t" is defined as the increase in dry weight of plant material from a unit area per unit of time.

$$CGR (g/m^2/day) = \frac{W_2 - W_1}{T_2 - T_1} \times \frac{1}{A}$$

W_1 = Total dry weight of plant at time t_1 W_2 = Total dry weight of plant at time t_2 t_1 = Time at first observation t_2 = Time at second observation A = ground area (m^2)

From the individual plot, the net plot area was harvested separately and produce was sun dried. After sun drying, the crop was threshed and produce cleaned. The final weight was recorded in $kg \text{ plot}^{-1}$ and finally converted into $kg \text{ ha}^{-1}$. Stover yield of mustard was calculated with subtraction of seed yield from biological yield and reported in $kg \text{ ha}^{-1}$. After 3-4 days sun drying, all above the ground plant parts of the net plot were dried and weighed in $kg \text{ plot}^{-1}$ to represent the biological yield and finally converted into $kg \text{ ha}^{-1}$. The harvest index will be computed by calculate the harvest index.

$$H.I (\%) = \frac{\text{Economic yield (kg/ha)}}{\text{Biological yield (kg/ha)}} \times 100$$

Result and discussion

Growth parameters:

Perusal of data given in Table 1 Showed that Plant height was significantly influence by the application of liquid biofertilizer and variable source of nutrients at harvest. In case of variable source of nutrients, treatment N₄ (75% NPKS + 2-ton VC ha⁻¹) recorded highest plant height (203.55 cm) as compared to other source of nutrients. However, the lowest plant height was recorded under N₁ (Control) at harvest. These results are in conformity with the findings of **Shivendu et al. (2019)**. Similarly, it was indicated from the data in the table that plant height also differs with application of different biofertilizers. Amongst, all the treatments, application of B₅ treatment (Azotobacter + PSB + KMB + Zn SB each @ 20 ml/kg ST) recorded superior plant height (201.40 cm) at harvest. The lowest plant height was recorded under the treatment B₁ (No biofertilizers) at harvest. However, the non-significant effect was found in respect to biofertilizer treatments at harvest. These results are in conformity with the findings of **Krishna et al. (2016) in mustard and Dhaka et al. (2001) in raya**. Number of branches increased with advancement of crop age and reached maximum at harvest. The number of primary and secondary branches at different stages was influenced by the application of liquid biofertilizer and variable source of nutrients. The treatment, N₄ (75% NPKS + 2-ton VC ha⁻¹) recorded significantly higher number of primary branches plant⁻¹ (5.87) at harvest. The lowest primary branches plant⁻¹ were observed in N₁ (Control). Similar findings were also reported by **Kansotia et al. (2013)**. Data indicated that number of primary branches also significantly influenced by different treatments of biofertilizers. Amongst the treatments, the B₅ treatment (Azotobacter + PSB + KMB + Zn SB each @ 20 ml/kg ST) recorded higher number of primary branches plant⁻¹ (5.82), which was significantly superior then remaining treatments of biofertilizers. However, the significantly lowest number of primary branches were recorded under B₁ treatment (No biofertilizer). Similar findings were also reported by **Singh et al. (2014)**. The treatment, N₄ (75% NPKS + 2-ton VC ha⁻¹) recorded significantly higher number of secondary branches plant⁻¹ (13.20) at harvest stage, respectively. which was statistically at par with N₂ (100% NPKS @ (120:60:60:40) kg ha⁻¹) at harvest. However, The lowest number of secondary branches plant⁻¹ recorded with N₁ (Control). Similar findings were also reported by **Thakur et al., (2009)**. Data indicated that number of secondary branches also influenced by different treatments of biofertilizers. Amongst the treatments, the B₅ treatment (Azotobacter + PSB + KMB + Zn SB each @ 20

ml/kg ST) recorded higher number of secondary branches plant⁻¹ (12.96), at harvest stage and the lowest number of secondary branches were recorded under B₁ (No biofertilizer). However, the non-significant effect in respect to number of secondary branches under the different treatment of biofertilizer. These findings are conformed with the findings of **Sahoo et al., (2010)**. plant dry weight was significantly influenced by the application of liquid biofertilizer and variable source of nutrients. Amongst variable source of nutrients, the treatment N₄ (75% NPKS + 2-ton VC ha⁻¹) recorded the significantly higher plant dry weight (66.65 g plant⁻¹) at harvest, as compared to other source of nutrients. However, the significantly lowest dry matter accumulation was noticed in N₁ (Control) treatment. Similar results were also reported by **Meena et al., (2014)** and **Thakur et al., (2009)**. Plant dry weight significantly varied with the treatments of different biofertilizer. Amongst all the treatments, the B₅ treatment (Azotobacter + PSB + KMB + Zn SB each @ 20 ml/kg ST) recorded significantly higher dry matter accumulation (64.98 g plant⁻¹) at harvest. which was statistically at par with B₄ (Azotobacter + PSB + KMB each @20 ml/kg ST) at 30 DAS and at harvest. However, the significantly lowest plant dry weight was found under treatment B₁ (No biofertilizer). These results were in agreement with the findings of **Singh and Singh (2014)**. Maximum LAI (2.929) was recorded with the application of N₄ (75% NPKS + 2-ton VC ha⁻¹) at 90 DAS, which was significantly superior than rest of the source of nutrients. However, minimum LAI was recorded under N₁ (Control) at harvest. Similar findings were also reported by **Kashvedet al., (2010)** and **Krishna et al., (2016)**. Similarly, application of B₅ (Azotobacter + PSB + KMB + Zn SB each @ 20 ml/kg ST) recorded significantly higher (2.702) leaf area index than B₁. The lowest leaf area index was recorded under B₁ (No biofertilizers). Similar findings were also reported by **Krishna et al., (2016)**. The maximum Crop growth rate (4.903 g m⁻² day) was recorded in the application of N₄ (75% NPKS + 2-ton VC ha⁻¹) at harvest. However, the minimum crop growth rate was recorded under N₁ (Control). Similar findings were also reported by **(Bhat et al., 2007 and Tripathi et al., 2010)**. Similarly, in case of biofertilizers, the application of B₅ (Azotobacter + PSB + KMB + Zn SB each @ 20 ml/kg ST) recorded significantly higher Crop growth rate (4.563 g m⁻² day) at harvest as compare to B₁. While the lower Crop growth rate was recorded under B₁ (No biofertilizers). Similar findings were also reported by **Saikia et al., (2013)**.

Table-1 Effect of liquid biofertilizers and variables source of nutrients on the growth parameters of mustard at harvest stage.

Treatments		Plant height (cm)	Primary branches plant ⁻¹	Secondary branches plant ⁻¹	Dry matter accumulation (g plant ⁻¹)	Leaf area index	Crop Growth Rate (g m ⁻² day ⁻¹)
Factor A (Source of nutrients)							
N ₁	Control	186.32	5.20	12.26	56.40	2.383	3.858
N ₂	100% NPKS @ (120:60:60:40) Kg ha ⁻¹	197.43	5.54	12.90	62.22	2.466	4.783
N ₃	75% NPKS + 6 ton FYM ha ⁻¹	202.46	5.72	12.78	64.36	2.689	4.593
N ₄	75% NPKS + 2 ton VC ha ⁻¹	203.55	5.87	13.20	66.65	2.929	4.903
<i>SEm ±</i>		2.23	0.07	0.09	0.70	0.077	0.273
<i>CD (P= 0.05)</i>		6.41	0.20	0.25	2.02	0.221	0.784
Factor B (Biofertilizers)							
B ₁	No biofertilizers	193.91	5.50	12.56	59.76	2.587	4.416
B ₂	Azotobacter @ 20 ml/kg ST	195.37	5.54	12.80	61.32	2.622	4.498
B ₃	Azotobacter + PSB each @ 20 ml/kg ST	197.26	5.46	12.70	62.32	2.574	4.518
B ₄	Azotobacter + PSB + KMB each @20 ml/kg ST	199.25	5.60	12.89	63.65	2.598	4.676
B ₅	Azotobacter + PSB + KMB + Zn SB each @ 20 ml/kg ST	201.40	5.82	12.96	64.98	2.702	4.563
<i>SEm ±</i>		2.39	0.07	0.10	0.78	0.036	0.04
<i>CD (P= 0.05)</i>		6.90	0.22	0.30	2.26	0.091	0.12

ST: Seed treatment, **PSB:** Phosphorus solubilizing bacteria, **VC:** Vermi-compost, **KMB:** Potassium mobilizing bacteria, **ZnSB:** Zinc solubilizing bacteria

Table-2 Effect of liquid biofertilizers and variables source of nutrients on seed, stover, biological yield (kg ha⁻¹) and Harvest index of mustard.

Treatment		Yield (kg ha ⁻¹)			
		Seed	Stover	Biological	H.I (%)
Factor A (Source of nutrients)					
N ₁	Control	632	4242	4875	12.97
N ₂	100% NPKS @ (120:60:60:40) Kg ha ⁻¹	1363	5901	7264	18.71
N ₃	75% NPKS + 6 ton FYM ha ⁻¹	1743	6740	8483	20.48
N ₄	75% NPKS + 2 ton VC ha ⁻¹	1854	7193	9047	20.39
<i>SEm ±</i>		8.13	47.10	45.84	0.17
<i>CD (P= 0.05)</i>		23.36	135.35	131.74	0.50
Factor B (Biofertilizers)					
B ₁	No biofertilizers	1123	5136	6260	17.43
B ₂	Azotobacter @ 20 ml/kg ST	1250	5613	6863	17.61
B ₃	Azotobacter + PSB each @ 20 ml/kg ST	1388	6072	7460	17.90
B ₄	Azotobacter + PSB + KMB each @20 ml/kg ST	1496	6387	7883	18.28
B ₅	Azotobacter + PSB + KMB + Zn SB each @ 20 ml/kg ST	1735	6888	8623	19.45
<i>SEm ±</i>		9.08	52.66	51.25	0.19
<i>CD (P= 0.05)</i>		26.12	151.35	147.29	0.56

ST: Seed treatment, **PSB:** Phosphorus solubilizing bacteria, **VC:** Vermi-compost, **KMB:** Potassium mobilizing bacteria, **ZnSB:** Zinc solubilizing bacteria

Perusal of data given in Table 2 Showed that seed yield of mustard influenced significantly with the application of variable source of nutrients. The significant highest value of seed yield (1854 kg ha^{-1}) was recorded under the treatment N_4 (75% NPKS + 2-ton VC ha^{-1}) followed by N_3 (75% NPKS + 6-ton FYM ha^{-1}) and lowest (632 kg ha^{-1}) in control treatment. Similar results were also reported by **Kansotia et al. (2013) and Singh et al. (2014)**. Similarly, with the application of different biofertilizer influenced significantly in the seed yield of mustard. The highest seed yield (1735 kg ha^{-1}) recorded under the treatment B_5 (Azotobacter + PSB + KMB + Zn SB each @ 20 ml/kg ST) which was followed by B_4 (Azotobacter + PSB + KMB each @ 20 ml/kg ST) and the lowest seed yield (1123 kg ha^{-1}) was recorded under the treatment B_1 (No biofertilizer). These results are confirmed with the findings of **Kumar et al., (2006) and Singh and Singh (2014)**. The significant highest value of stover yield (7193 kg ha^{-1}) was recorded under the treatment N_4 (75% NPKS + 2-ton VC ha^{-1}) followed by N_3 (75% NPKS + 6-ton FYM ha^{-1}) and lowest (4242 kg ha^{-1}) in control treatment. (**Bhat et al., 2007 and Tripathi et al., 2010**). Similarly, with the application of different biofertilizer influenced significantly in the stover yield of mustard. The highest stover yield (6888 kg ha^{-1}) under the treatment B_5 (Azotobacter + PSB + KMB + Zn SB each @ 20 ml/kg ST) followed by B_4 (Azotobacter + PSB + KMB each @ 20 ml/kg ST) and the lowest stover yield (5136 kg ha^{-1}) was recorded under the treatment B_1 (No biofertilizer). The results are in close conformity with the findings of **Krishna et al., (2016) and Kumar et al., (2019)**. The biological yield of mustard significantly influenced with the application of variable source of nutrients. The significant highest value of biological yield (9047 kg ha^{-1}) was recorded under the treatment N_4 (75% NPKS + 2-ton VC ha^{-1}) followed by N_3 (75% NPKS + 6-ton FYM ha^{-1}) and lowest (4875 kg ha^{-1}) in control treatment. Similar results were also reported by **Kansotia et al. (2013) and Singh et al. (2014)**. Similarly, with the application of different biofertilizer influenced significantly in the biological yield of mustard. The highest biological yield (8623 kg ha^{-1}) under the treatment B_5 (Azotobacter + PSB + KMB + Zn SB each @ 20 ml/kg ST) followed by B_4 (Azotobacter + PSB + KMB each @ 20 ml/kg ST) and the lowest biological yield (6260 kg ha^{-1}) was recorded under the treatment B_1 (No biofertilizer). The results are in close conformity with the findings of **Krishna et al., (2016) and Kumar et al., (2019)**. From the table it is clear that in comparison to N_1 control, the significantly higher value of harvest index (20.48) was observed by the application of N_4 (75% NPKS + 2-ton VC ha^{-1}), which is statistically at par with the treatment N_3 (75% NPKS + 6-ton FYM ha^{-1}). The lowest value of harvest index (12.97) was noticed in N_1 (Control). Similar results were also reported by **Tripathi et al., (2010)**. Application of different biofertilizer influenced in harvest index of mustard. The

significantly higher value of harvest index (19.45) recorded under the treatment B₅ (Azotobacter + PSB + KMB + Zn SB each @ 20 ml/kg ST) followed by B₄ (Azotobacter + PSB + KMB each @20 ml/kg ST) and the lowest value of harvest index (17.43) was recorded under the treatment B₁ (No biofertilizer). The results are in close conformity with the findings of Kumar *et al.*, (2019).

Conclusion

Thus, the application of 75%NPKS (90:45:45:30) + 2-ton VC ha⁻¹) + B₅ (Azotobacter + PSB + KMB + Zn SB each @ 20 ml/kgSeed treatment) seems to be best option for maximum plant growth and higher yield under western UP. The study is to be continuing for few more years to draw definite conclusion for application of liquid biofertilizers and variable source of nutrients in mustard.

REFERENCES

- Anonymous, 2011. A comprehensive global collection of statistics on agriculture. Food and Agriculture Organization of the United Nations.
- Anonymous, 2021. Response of macro and micro nutrient bio-fertilizers in enhancing rapeseed-mustard productivity and soil health, Directorate of Rapeseed-Mustard Research, Bharatpur-321303, Rajasthan. *Annual report 2021*: CP18-21.
- Bhat, M.A., Singh, R. and Kahli, A. 2007. Effect of integrated use of farm yard manure and fertilizernitrogenwithandwithoutSulphuronyieldandqualityofIndianmustard(*Brassica juncea*). *JournaloftheIndianSocietyofSoilScience* **55**:224-226.
- Dhaka, A.K., Satish, K. and Kadia, V.S.(2001). Effect of inorganicand in organic manures on growth indices of late planted Raya. *Haryana Journal Agronomy* **17**(1&2): 170-172.
- Hegde, D.M. and Sudhakara Basu, S.N. 2004. Balanced fertilization for nutritional quality inoilseeds. *FertilizerNews* **49**(4):57-62,65-66.
- Kansotia, B., Meena, R. and Meena, V. (2013). Effect of vermicompost and inorganicintegrated nutrient management in mustard [*Brassica juncea*(L.) Czern&Coss]. *International Journal ofAgricultural Sciences* **10**(2):667-670.
- Kashved,S.M.,Raskar,B.S.andTamboli,B.D.2010.Effectofintegratednitrogenmanagement and irrigation regimes on productivity of mustard (*Brassica juncea*L.). *Journalof Maharashtra Agricultural universities* **35**: 349-353.
- Krishna, M., Sanjay, C. and Hemanth, K. 2016. Effect of integrated nutrient management on growth and yield in mustard (*Brassica juncea*L.) *International Journal of*

Chemical Studies **6**(2): 3571-3573.

- Kumar, N. 2006. Integrated nutrient management practices in mustard (*Brassica juncea* L.) and its effect on the productivity of succeeding rice crop. Ph.D. thesis submitted to Chaudhary Charan Singh University, Meerut.
- Kumar, V., Singh, S. 2019 Effect of fertilizers, biofertilizers and farmyard manure on sustainable production of Indian mustard (*Brassica juncea*) *Annals of Plant and Soil Research* **21**(1):25-29.
- Meena, B. P., Kumar, A., Dotaniya, M. L., Jat, N.K. and Lal, B. 2014. Effect of organic sources of nutrients on tuber bulking rate, grades and specific gravity of potato tubers. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences*. DOI 10.1007/s40011-014-0398-4.
- Sahoo, S. K., Dwivedi, S. K. and Pradhan, L. 2010. Effect of Biofertilizers and Levels of Nitrogen on Yield and Nutrient Uptake of Indian Mustard (*Brassica juncea*). *Environment and Ecology* **28**(1): 129-131.
- Saikia, R., Bora, P.C. and Sarma, A. 2013. Effect of integrated nutrient management (INM) on productivity, nutrient uptake and economics of rape seed (*Brassica campestris* var. Toria) in Assam. *Advances in Plant Sciences* **26**(2):491-493.
- Shivendu, K. C., Sanjay, K.S. and Anil, P. 2019. Influence of integrated nutrient management on yield and quality of Indian mustard (*Brassica juncea* L.) in calcareous soil of Bihar. *Annals of Plant and Soil Research* **21**(1):76-81.
- Singh, A. and Meena, N.L. 2004. Effect of nitrogen and Sulphur on growth yield attributes and seed yield of mustard (*Brassica juncea*) in eastern plain of Rajasthan. *Indian Journal of Agronomy* **49**(3):186-188.
- Singh, R. K. and Singh, A. K. 2014. Production potential, nutrient uptake and economics of Indian mustard (*Brassica juncea*) under integrated nutrient management practices. *Indian Journal of Agricultural Sciences* **84**: 142-148.
- Singh, R., Anil, K. S. and Praveesh, K. 2014. Performance of Indian mustard (*Brassica juncea* L.) in response to integrated nutrient management. *Journal of Agricultural Research* **7**(2):104-107.
- Thakur, A.K., Choudhari, S.K., Singh, R. and Kumar, A. 2009. Performance of rice varieties at different spacing grown by the system of rice intensification in eastern India. *Indian Journal of Agricultural Sciences* **79**(6): 443-447.
- Tripathi, M. K., Chaturvedi, S., Shukla, D. K. and Mahapatra, B. S. 2010. Yield performance

and quality in Indian mustard (*Brassica juncea*) as affected by integrated nutrient management. *Indian Journal of Agronomy* **55**(2): 138-142.

UNDER PEER REVIEW