

Original Research Article

Efficacy of Combined Eco-Friendly Evaluation of biocontrol-based Formulation formulations for Managing against Late Leaf Spot and Rust Diseases and Enhancing Yield in Groundnut

Formatted: Left

ABSTRACT

This study investigated the efficacy of various ~~eco-friendly biocontrol-based components formulations~~ against late leaf spot (LLS) (*Phaeoisariopsis personata*) and rust (*Puccinia arachidis*) in groundnut. Pot culture and field experiments were conducted during the Kharif season (June-September) of 2022. Disease incidence of both late leaf spot and rust was significantly lower in groundnut plants treated with a combination of ~~biocontrol-based formulation~~ ~~eco-friendly formulations~~ compared to control. The most effective eco-friendly treatment included a combination of *Trichoderma asperellum*, *Pseudomonas fluorescens*, fortified lignite fly ash and Annamalai mixture. This combination significantly reduced lesion frequency (~~0.38, 0.72 and 1.08~~) and Percent Disease Index (PDI) (~~8.0, 9.4 and 12.5~~) for both diseases at (50, 70 and 90 DAS) ~~respectively~~, compared to other treatments. The ~~combination of combined eco-friendly treatment~~ ~~biocontrol-based formulations~~ also significantly increased the pod yield (~~67g/plant~~). Furthermore, the same treatment increased the induction of ~~defence~~ ~~defense~~ enzymes peroxidase (2.22 fold), polyphenol oxidase (1.28 fold) and phenylalanine ammonia-lyase (6.25 fold). While Carbendazim, ~~a chemical fungicide~~, effectively reduced disease incidence, it was surpassed by the ~~biocontrol-based module~~ ~~combined eco-friendly treatment~~ in terms of both disease control and yield. These findings suggest that ~~biocontrol-based formulations~~ ~~a combination of eco-friendly formulations can be~~ holds promise as a sustainable approach for managing LLS and rust diseases in groundnut cultivation. ~~This method offers potential benefits over chemical fungicides, including reduced environmental impact and potentially higher crop yields.~~

Keywords: Leaf spot, rust, Defence ~~defense~~ enzyme, survey, biocontrol-based module fortified lignite fly ash, Annamalai mixture, groundnut

Formatted: Font: Not Bold

1. INTRODUCTION

Groundnut (*Arachis hypogaea* L.), ~~belonging to the Fabaceae family, is a globally important legume prized for its nutritional and agricultural contributions. Ois~~ often hailed as the "king of oilseeds," ~~." Globally, India holds a dominant position, holding~~ has the largest cultivation area ~~and~~ ranking second in global production. In India, groundnut production in the kharif season is 68.57 lakh tonnes with an average productivity of 1,562 kg/ha (IOPEPC, 2023). ~~Groundnut offers a compelling nutritional profile, rich in edible oil and vegetable protein [1]. Beyond its direct benefits, this legume fosters soil health through symbiotic nitrogen fixation with Rhizobia bacteria, promoting sustainable agricultural practices.~~ ~~Despite its economic and nutritional significance, g~~ Groundnut cultivation faces a formidable challenge from fungal diseases like late leaf spot (LLS), rust, ~~alternaria~~ *Alternaria* blight, stem rot, dry root rot and collar rot [2,3]. ~~Among these, LLS disease caused by~~ *Phaeoisariopsis personata* and rust disease caused by *Puccinia arachidis* pose the most significant threats. These widespread fungal pathogens inflict severe damage on groundnut crops. In most cases, these two pathogens occur together and hamper groundnut production [4]. In India, these diseases have been documented to cause yield losses exceeding

Formatted: Indent: First line: 0"

Comment [GN1]: Please change the serial numbers as ref.1 is deleted.

70%, significantly impacting agricultural productivity[5]. Leaf rust alone could cause yield reduction reaching up to 65%, especially in areas with high rainfall [6]. The detrimental effects extend beyond yield reduction, as the quality of groundnut seeds is also compromised by these fungal infections.

Comment [GN2]: Groundnut? If so, please change accordingly.

Formatted: Highlight

In groundnut cultivation, conventional disease management practices for controlling LLS and rust often rely on chemical fungicides. ~~T~~ Nevertheless, their indiscriminate use ~~raises~~ has led to concerns about environmental and human health risks[2]. Environmental contamination, disruption of ecological balance due to harm to beneficial soil microbes, and potential health risks to humans and non-target organisms are some of the concerning consequences associated with chemical fungicide application[7,8]. The emergence of eco-friendly alternatives presents a promising solution for sustainable groundnut production[9]. Antagonistic microorganisms offer a particularly effective approach for managing these fungal diseases. These beneficial microbes can act against pathogens through multiple mechanisms, including competition for resources, production of antibiosis compounds, predation and parasitism [10]. Additionally, some antagonists have the potential to induce systemic resistance within the plant itself[11]. These mechanisms offer a multifaceted approach to disease control, potentially mitigating the need for harmful chemical fungicides.

~~This research delves into the potential of eco-friendly components, particularly work explores use of~~ antagonist microorganisms, as a ~~sustainable and effective disease management strategy against~~ strategy to enhance groundnut's defense mechanisms against the detrimental fungal pathogens, *P.personata* and *P.arachidis* in groundnut. ~~By exploring this approach, we aimed to the development of sustainable and effective disease management practices for groundnut cultivation.~~

2. MATERIALS AND METHOD

2.1 Survey of late leaf spot and rust disease incidence in groundnut

~~This study employed a~~ Field survey ~~approach~~ was employed to assess the prevalence and severity of groundnut LLS and rust diseases in Cuddalore district. Villages with a history of groundnut cultivation were targeted to capture representative data. Ten locations were chosen for fixed-plot surveys. Within each plot, the number of groundnut plants exhibiting symptoms of LLS and rust disease were recorded alongside the total number of plants observed.

Formatted: Normal (Web), Indent: First line: 0.5", Right: 0", No bullets or numbering, Pattern: Clear (White)

2.2 Inoculum culture of *P. personata* and *P. arachidis*

Naturally infected leaves with LLS and rust disease symptoms were collected from fields within the Cuddalore district. These diseased leaves served as the inoculum source for *P. personata* and *P. arachidis*. The inoculum was maintained through pot culture containing groundnut plants.

Comment [GN3]: How was the spore suspension made? What was the spore strength applied as inoculum, please?

Formatted: Highlight

Formatted: Font: Italic

2.3 Artificial Inoculation Method ~~Assessment of disease severity~~

Groundnut seeds (cv. VRI 2) were grown in pots (20 cm dia, 5 plants/pot) for 30 days within a polyhouse environment. The potting medium consisted of a 2:1:1 ratio of field soil, farmyard manure (FYM) and sand. The plants were then inoculated with a spore suspension containing both mixture ~~P. of~~ *P. personata* and *P. arachidis*. To achieve optimal infection conditions, an alternating wet and dry period was implemented following inoculation. The inoculated plants were maintained at $24 \pm 2^\circ\text{C}$. Each treatment group within the glasshouse experiments comprised 12 plants with three replicates. Disease severity of both *P. personata* and *P. arachidis* were assessed using disease incidence and lesion frequency following the methods established by Subrahmanyam *et al.* (12).

Comment [GN4]: Please mention how many plants per pot were present for each group. Are there 4 plants per pot and three such pots were three replicates? Please explain.

2.3.1 Assessment of disease severity

Quantification of lesion incidence in response to various treatments for combined LLS and rust disease [4].

Lesion frequency (LF) = Number of lesion or pustules/cm² leaf area
Disease scoring was done with modified 9-point (1-9) scale [12].

$$\text{Percent diseases index (PDI)\%} = \frac{\text{Sum of individual rating}}{\text{Total no. of leaves observed}} \times \frac{100}{\text{Maximum grade}}$$

2. 4 Annamalai mixture preparatioPreparation of biocontrol based formulations.

Aliquots of ~~bovine urine~~, bovine dung, ovine dung, poultry litter, and neem cake were obtained at a conc. of 100% (w/w) and homogenously mixed in a 1:1:1:1 ratio ~~to prepare Annamalai mixture~~. The resulting organic amendment was subsequently diluted to 20%, 40%, 60%, and 80% (v/v) using distilled water at the time of application[13].

2. 5 Lignite fly ash fortification

~~The antagonists~~ *T.asperellum* and *P. fluorescens*, ~~the bio-inoculants~~ were cultivated in a Potato Dextrose broth and King's B broth respectively. Class F lignite fly ash (LFA), obtained from Neyveli Lignite Corporation, Neyveli, India, was used as the carrier material. One kilogram of LFA was mixed thoroughly with 10 grams of carboxymethyl cellulose (CMC). The LFA-CMC mixture was then autoclaved for two consecutive days to ensure proper sterilization. ~~To 1000 g of sterilized LFA-CMC carrier~~, Followed by 400 ml of each ~~bacterial suspension~~, containing approximately 8×10^9 CFU/mL, were separately added ~~to 1000 g of sterilized LFA-CMC carrier~~. 20 ml of molasses were then incorporated into the mixture, followed by thorough mixing. The fortified LFA was shade-dried for 2 hours under sterile conditions to remove excess moisture. The final product, designated as fortified lignite fly ash, was packaged in polyethylene bags and stored at $28^\circ\text{C} \pm 2^\circ\text{C}$. To monitor the viability, the population of antagonistic strains within the fortified LFA was periodically evaluated using the serial dilution technique[14].

2. 6 Efficacy of Evaluation of eco-friendly productsantagonists based formulations against LLS and rust in groundnut under pot culture condition

~~A study was carried out under pot culture conditions to assess the effect of various eco-friendly components against LLS and rust in groundnut~~. The experiment included 10 treatments with three replications, conducted from June to September 2022 at the Department of Plant Pathology, Annamalai University. Previously isolated and available biocontrol agents (BCAs), specifically *T.asperellum*(PP348038) and *P. fluorescens*(PP348041) were evaluated for their ability to suppress the targeted pathogens. Fifteen kilograms of sterilized topsoil from a groundnut field were filled in 45 x 30 cm earthen pots. Groundnut seeds of a susceptible cultivar (VRI-2) were surface sterilized using 3% NaOCl solution for 30 sec, followed by two washes with sterile water. Five seeds were sown per pot. ~~Eco-friendly components~~ ~~Biocontrol-based components~~ including these BCAs, fortified lignite fly ash (FL-ash) and Annamalai mixture were tested against LLS and rust. The treatments include talc formulation of *T.asperellum* and *P. fluorescens* (2×10^8 CFU g⁻¹) as seed treatments (10 g/kg) and soil drench after sowing, FL-ash as seed treatment (10 g/kg) and soil drench (40 kg/ha equivalent), Annamalai mixture as a seed treatment (10 ml/kg) and foliar spray (20 L/ha equivalent). Spore suspensions of *P. personata* and *P. arachidis* (10^8 spores/ml) were sprayed on plants in the evening at two-time points with a four-day interval, starting 30 days after sowing (DAS). Agar plate technique was employed to confirm the absence of seed-borne pathogens before sowing. Disease incidence and severity were likely assessed throughout the experiment.

2. 7 Efficacy of eco-friendly amendmentsField evaluation of Biocontrol based strategy against *P.personata* and *P.arachidis* in field-grown groundnut

A field experiment was carried out in an endemic field from June to September, 2022 in Sivapuri village, Cuddalore, Tamil Nadu. Groundnut cv. VRI-2 was sown in 33 x 13 feet

Comment [GN5]: Please check: PDI= Sum of individual rating * 100/ (Total no. of leaves observed * Maximum grade)

Formatted: Font: Highlight

Formatted: Font: Highlight

Formatted: Font: Highlight

Formatted: Font: Highlight

Formatted: Font: Highlight

Comment [GN6]: This was taken by weight? Or litre?

Formatted: Font: (Default) Times New Roman, 12 pt, English (India), Highlight

Formatted: Font: (Default) Times New Roman, 12 pt, English (India), Highlight

Comment [GN7]: Was this mixture used as such or filtered?

Comment [GN8]: This mixture is too watery to be shade-dried only for two hours. Please check

Comment [GN9]: How many bacteria are used, please? The culture is only one *P. fluorescens* or many? The other antagonist is fungus.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, English (India), Highlight

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, English (India), Highlight

Formatted: Font: (Default) Arial, Bold, Not Italic

Formatted: Font: Bold, Not Italic

Formatted: Font: (Default) Arial

plots. The ~~same eco-friendly biocontrol-based~~ components tested in a previous greenhouse experiment were applied here except for pathogen control, following the same schedule. Standard agricultural practices were maintained throughout the season. PDI and lesion frequency along with growth plant parameters and yield attributes were likely measured.

2.87 Effect of eco-friendly amendments on activity of defence defense enzyme induction in groundnut against *P. personata* and *P. arachidis*

2.8.1 Sample Collection

A ~~In~~ polyhouse experiment ~~on~~ was employed to evaluate the efficacy of various ~~eco-friendly component formulations in inducing defense enzyme activity within~~ VRI-2 variety of groundnut (as mentioned in 2.3) against ~~*P. personata* and *P. arachidis*~~. Subsequent ~~subsequent~~ to inoculation with the ~~respective two target~~ pathogens at 35 days post-sowing, plant samples were obtained at designated time points (1-, 3-, 5- and 7- days post inoculation) from each treatment.

2.8.2 Enzyme extraction

Groundnut plant tissues were immediately frozen in liquid N₂ and subsequently homogenized. A ~~sample of~~ ~~1 g~~ aliquot of the powdered sample was homogenized ~~mixed~~ with 2 mL of 0.1 M sodium phosphate buffer (pH 7.0) at 4°C. The homogenate was subsequently centrifuged at 10,000 rpm for 20 min. ~~and used for enzyme assays.~~ The activity of peroxidase (PO), polyphenol oxidase (PPO) and phenylalanine ammonia lyase (PAL) were assayed in the resultant supernatant.

2.8.3 Enzymatic assay

2.8.3.1 Peroxidase (PO)

Peroxidase (PO) activity was determined by following the method of Srivastava [15]. The reaction mixture contained 1.5 mL of 0.05 M pyrogallol, 0.5 mL of enzyme extract, and 0.5 mL of 1% hydrogen peroxide (H₂O₂) in a final volume of 2.5 mL. The mixture was incubated at 28±1°C. Absorbance change at 420 nm was monitored every 30 sec for a total duration of 3 min. ~~Boiled enzyme preparations served as a negative control. And~~ the enzyme activity was expressed as the change in absorbance/min/g of fresh weight tissue.

2.8.3.2 Polyphenol Oxidase (PPO)

~~This~~ The polyphenol oxidase (assay measures PPO) activity through the oxidation of catechol at 495 nm. The reaction mixture comprises 1.5 mL of 0.1 M sodium phosphate buffer (pH 6.5), 200 µL of enzyme extract, and 200 µL of 0.01 M catechol, was assayed as per Mayer and Harel (16). The change in absorbance at 495 nm/min/g of fresh-weight tissue is used to express PPO activity. ~~Phenylalanine Ammonia-Lyase (PAL)~~ [16].

2.8.3.3 Phenylalanine Ammonia-Lyase (PAL)

~~In~~ a cuvette, 100 µL of enzyme solution was combined with 500 µL of 50 mM Tris HCl buffer (pH 8.8) and 600 µL of 1 mM L-phenylalanine substrate solution. This reaction mixture was incubated for 60 minutes at 32± 2°C to allow enzymatic conversion of the substrate. The reaction was then quenched by the addition of 2 N HCl, stopping further enzyme activity. To extract the product, trans cinnamic acid, 1.5 mL of toluene was added, and the mixture was vigorously vortexed for 30 seconds to facilitate the transfer of the product into the organic phase. The mixture was then centrifuged at 1000 rpm for 5 minutes to achieve phase separation. The upper toluene layer, containing the extracted trans cinnamic acid, was carefully transferred and its absorbance was measured at 290 nm using a spectrophotometer. A blank cuvette containing only toluene served as the reference for background subtraction. A standard curve for trans cinnamic acid was previously established using known concentrations of the product in toluene. The enzyme activity was then ~~was~~ assayed following the procedure of Srivastava (15) ~~calculated~~ and expressed as micromoles (µmol) of trans-cinnamic acid formed per minute per gram of fresh tissue weight (µmol min⁻¹ g⁻¹ fresh weight) [15].

Comment [GN10]: Same method of inoculation?

Formatted: Font: (Default) Times New Roman, Not Italic

Formatted: Font: (Default) Times New Roman

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: (Default) Times New Roman, Not Italic

Formatted: Font: (Default) Times New Roman

Formatted: Font: Times New Roman, Not Italic

Formatted: F2-Body Text, Left

Formatted: Font: Not Italic

Formatted: Font: Times New Roman, Not Italic

Formatted: Font: Not Italic

Formatted: Font: Times New Roman, Not Italic

Formatted: Font: Times New Roman, Bold

Formatted: Font: (Default) Times New Roman, Not Italic

Formatted: Font: (Default) Times New Roman

Formatted: Font: Times New Roman, Not Italic

Formatted: F2-Body Text, Left, Right: 0", Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Formatted: Font: Not Italic

Formatted: Font: Times New Roman, Not Italic

Formatted: Font: Not Italic

Formatted: Font: (Default) Times New Roman, Not Italic

Formatted: Font: Not Italic

Formatted: Font: (Default) Times New Roman, Not Italic

Formatted: Font: Times New Roman

2. 9-8 STATISTICAL DATA ANALYSIS

A randomized block design (RBD) with three replicates was used for the pot culture and field experiments. Statistical analysis was performed using SAS software (version 9.4, SAS Institute, Cary, NC, USA) to assess treatment effects. Following a significant analysis of variance (ANOVA), Dunnett's multiple comparison test (DMRT) was employed to identify statistically significant differences ($p < 0.05$) between treatment means and the control. All the treatments were replicated thrice.

3. RESULTS AND DISCUSSION

3. 1 Survey on the disease incidence

A spatial survey was conducted in various locations within the Cuddalore district, Tamil Nadu, to assess the prevalence of LLS and rust diseases in groundnut crops. The investigation revealed a substantial variation in disease incidence across the surveyed locations. LLS exhibited a prevalence ranging from 8.24% to 32.84%, with the highest incidence observed in Mathur. Similarly, rust incidence varied between 7.21% and 30.78%, with Mathur again recording the most severe infection. These findings suggest that geographical factors, cultivar type, and pathogen genetics within the region may significantly influencing the severity and distribution of the pathogens [17-19]. The co-infection of both *P. personata* and *P. arachidis* are causes severe foliar disease in all major groundnut growing areas of Tamil Nadu.

3.2 Evaluation of various Efficacy eco-friendly biocontrol-based components against *P. personata* and *P. arachidis* isolates under pot culture conditions

The study investigated the effectiveness of eco-friendly formulations compared to a control group. Results demonstrated a significant reduction in disease incidence of LLS and rust for in plants treated with several eco-friendly biocontrol formulations based formulations was found compared to the control. Among the treatments, the combination (seed and soil application) of *T. asperellum*, *P. fluorescens*, fortified lignite fly ash, and Annamalai mixture (denoted as T7) exhibited the most pronounced disease reduction. T7 had most significantly lowered the lesion frequency of LLS (0.38, 0.72 and 1.08) and disease severity index (PDI) (8.0, 9.4 and 12.5) at various observation points (50, 75 and 90 DAS) respectively. The same treatment also significantly reduced the disease incidence of rust which recorded 0.24, 0.51, 0.83 lesion frequency and 5.9, 7.6, 9.8 and PDI of rust at 50, 70 and 90 DAS respectively. Additionally, T7 resulted in a considerable increase in pod yield (67 g/plant). The performance of T7 was followed by the application of carbendazim (chemical fungicide) which also displayed a significant reduction in LLS (0.47, 0.94 and 1.12 lesion frequency and 9.2, 10.8 and 13.7 PDI) and rust (0.49, 0.73, 0.99 lesion frequency and 6.9, 8.1, 10.7 PDI) at 50, 70 and 90 DAS respectively, compared to the control. However, the combined eco-friendly formulation (T7) demonstrated superior disease control and yield improvement compared to the chemical treatment (carbendazim). Overall, the findings also suggested that strategically combined eco-friendly biocontrol-based formulations offer a promising approach for managing LLS and rust diseases, potentially surpassing the efficacy of individual eco-friendly components and even a standard chemical fungicide (Table 1 and Fig. 1). Being a combination of biocontrol agents and eco-friendly amendments is an added advantage in that the microbial consortia could potentially proliferate in the soil and survive throughout the cropping season and protect the crop from harmful pathogenic infection [20,21]. The Earlier workers have proven that the native isolate of bioagents performs well ultimately reducing could reduce the disease intensity of LLS and rust [22,23]. During the tripartite interaction, the biocontrol agents like *Trichoderma* and *Pseudomonas* produce certain kind of antibiotics, iron-chelating agents, cell wall degrading enzymes and secondary metabolites that acts on pathogens leads to suppressing the pathogens' growth effectively [24,25]. Annamalai mixture has been proven as an effective disease suppressor and plant

Comment [GN11]: Pot culture expts. are done usually as CRBD.

growth promoter. This mixture contains animal excretes that improve the soil health and protect the plants against various diseases by promoting plant growth [14,13].

Comment [GN12]: Reference?

Table 1. Evaluation of various eco-friendly biocontrol based formulations components against *Phaeoisariopsis personata* and *Puccinia arachidis* Leaf spot and rust under pot culture condition

Formatted: Font: Not Italic

Treatment	Disease incidence						PDI						Yield g/plant
	Late leaf spot (No. of lesion)			Rust (No. of pustule)			Late leaf spot			Rust			
	50 DAS	70 DAS	90 DAS	50 DAS	70 DAS	90 DAS	50 DAS	70 DAS	90 DAS	50 DAS	70 DAS	90 DAS	
T ₁	0.98 ^d	1.14 ^d	1.32 ^e	0.64 ^e	0.89 ^e	1.27 ^e	12.9 (21.04) ^e	13.6 (21.64) ^e	14.1 (22.05) ^e	10.8 (19.18) ^e	11.2 (19.55) ^e	13.8 (21.80) ^e	53 ^e
T ₂	1.34 ^g	1.52 ^g	2.02 ^h	1.10 ^g	1.48 ^h	1.63 ^h	15.9 (23.49) ^h	16.1 (73.65) ^h	16.9 (24.27) ^h	15.2 (22.94) ^h	16.4 (23.88) ^h	17.7 (24.87) ^h	50 ^h
T ₃	0.63 ^c	1.02 ^c	1.24 ^d	0.49 ^d	0.71 ^d	1.09 ^d	10.3 (18.71) ^d	11.9 (20.17) ^d	13.6 (21.64) ^d	7.0 (15.34) ^d	8.3 (16.74) ^d	11.7 (20.00) ^d	57 ^d
T ₄	1.19 ^e	1.26 ^e	1.57 ^f	0.87 ^f	1.16 ^f	1.32 ^f	14.1 (22.05) ^f	15.3 (23.02) ^f	15.9 (24.27) ^f	12.5 (20.70) ^f	13.0 (21.13) ^f	13.9 (21.89) ^f	53 ^f
T ₅	1.27 ^f	1.44 ^f	1.63 ^g	1.05 ^g	1.39 ^g	1.52 ^g	15.0 (22.78) ^g	15.9 (23.49) ^g	16.3 (23.81) ^g	14.5 (22.38) ^g	15.3 (23.02) ^g	16.1 (23.65) ^g	51 ^g
T ₆	0.42 ^b	0.97 ^c	1.13 ^c	0.37 ^c	0.67 ^c	1.03 ^c	9.7 (18.14) ^c	10.2 (18.62) ^c	12.9 (21.04) ^c	6.2 (14.41) ^c	7.9 (16.32) ^c	10.8 (19.18) ^c	59 ^c
T ₇	0.31 ^a	0.62 ^a	0.98 ^a	0.18 ^a	0.31 ^a	0.60 ^a	7.3 (15.67) ^a	8.9 (17.35) ^a	11.3 (19.64) ^a	5.2 (13.18) ^a	7.1 (15.45) ^a	9.0 (17.45) ^a	69 ^a
T ₈	0.37 ^b	0.84 ^b	1.02 ^b	0.29 ^b	0.53 ^b	0.89 ^b	8.9 (17.35) ^b	9.8 (18.24) ^b	12.6 (20.79) ^b	5.9 (14.05) ^b	7.6 (16.00) ^b	9.9 (18.33) ^b	63 ^b
T ₉	3.11 ⁱ	4.56 ⁱ	6.05 ^j	2.17 ⁱ	3.63 ^j	5.22 ^j	38.3 (38.23) ^j	41.8 (40.28) ^j	42.1 (40.45) ^j	35.5 (36.57) ^j	38.7 (38.46) ^j	41.2 (39.93) ^j	45 ^j
T ₁₀	2.45 ^h	3.26 ^h	3.40 ⁱ	2.03 ^h	3.25 ⁱ	3.47 ⁱ	23.8 (29.19) ⁱ	28.3 (32.13) ⁱ	34.9 (36.21) ⁱ	19.6 (26.27) ⁱ	23.7 (29.13) ⁱ	30.8 (33.70) ⁱ	51 ⁱ

* Means of three replications, Values in the column followed by same letters not differ significantly by DMRT ($p=0.05$)



Comment [GN13]: Plant population seem to vary among the pots.

Formatted: Font: Not Italic

Fig. 1. Pot culture study of eco-friendly components against *Phaeoisariopsis personata* and *Puccinia arachidis*

3. 3 Evaluation of various eco-friendly biocontrol-based formulations components against *P. personata* and *P. arachidis* isolates under field conditions

The results of the potculture experiments were confirmed in the field. The experiment investigated the impact of these eco-friendly formulations on disease incidence and yield compared to a control. The aforesaid treatment (T7) demonstrated the most significant reductions in disease incidence for both LLS and rust. Treatment T7 displayed consistently lower lesion frequency and percent disease index (PDI) of LLS (0.22, 0.38, 0.51 and 7.20, 8.14, 9.50) and rust (0.12, 0.24, 0.41 and 3.92, 5.67, 6.81) diseases at 50, 70, and 90 DAS, respectively. Additionally, T7 resulted in the highest yield (2060 kg/ha). Carbendazim application (T8) served as a positive control and exhibited a lower disease incidence of late leaf spot (0.27, 0.44, 0.56 lesion frequency and 8.32, 8.89, 10.07 PDI) and rust (0.19, 0.27, 0.33 lesion frequency and 4.19, 6.05, 7.37 PDI) at 50, 70 and 90 DAS respectively, compared to the control but was surpassed by the combined eco-friendly formulation treatment (T7). Treatment T8 displayed a moderate reduction of LLS and rust along with a yield of 1984 kg/ha. The study suggests that the combined application of these eco-friendly biocontrol based formulations offers a more effective and potentially more sustainable approach for managing LLS and rust in groundnut crops compared to individual applications or a chemical fungicide control (Table 2 & Fig 2). Overall, the combined application of bioproducts is known to control diseases effectively by employing more than one mechanism of direct or indirect pathogen suppression [26,27]. This approach has been evaluated against various diseases with a combination of a variety of bioproducts including a combination of biocontrol agents [27]. Such approaches include, a combination of organic products with biocontrol agents [14], a combination of plant products with biocontrol agents [28,29] and a combination of fungicides with biocontrol agents [30]. These bioproducts also promoted plant growth significantly and increased the yield of the crop [31-33]. The major reason for the increased plant growth and yield parameters might be due to the significant induction of plant growth-promoting substances like auxin, gibberellic acid and abscisic acid induced by the bioproducts [34,35]. The lignite fly ash used as carrier material contains a trace amount of minor nutrients like Ca, K, and Si [36-38]. These minor nutrients trigger the plant defense response and protect the plants [39-41].

Table 2. Evaluation of various eco-friendly components against *Phaeoisariopsis personata* and *Puccinia arachidis* under field condition

Treatment	Disease incidence						PDI						Yield kg/ha
	Late leaf spot (No. of lesion)			Rust (No. of pustule)			Late leaf spot			Rust			
	50 DAS	70 DAS	90 DAS	50 DAS	70 DAS	90 DAS	50 DAS	70 DAS	90 DAS	50 DAS	70 DAS	90 DAS	
T ₁	0.38 ^c	0.54 ^c	0.72 ^c	0.34 ^c	0.38 ^c	0.43 ^c	10.09 (18.52) ^e	10.79 (19.17) ^e	10.97 (19.34) ^e	6.80 (15.11) ^e	7.14 (15.49) ^e	8.83 (17.28) ^e	1893 ^e
T ₂	0.48 ^d	0.61 ^d	0.79 ^d	0.42 ^d	0.48 ^d	0.53 ^d	11.76 (20.05) ^h	12.13 (20.38) ^h	12.69 (20.86) ^h	8.87 (17.32) ^h	9.24 (17.67) ^h	10.19 (18.61) ^h	1828 ^h
T ₃	0.35 ^b	0.50 ^c	0.69 ^b	0.29 ^c	0.31 ^b	0.39 ^b	9.14 (17.59) ^d	9.39 (17.84) ^d	10.51 (18.91) ^d	5.93 (14.09) ^d	6.94 (15.27) ^d	8.02 (16.45) ^d	1897 ^d
T ₄	0.41 ^c	0.56 ^d	0.75 ^c	0.37 ^d	0.42 ^c	0.47 ^c	10.51 (18.91) ^f	10.93 (19.30) ^f	11.39 (19.72) ^f	7.46 (15.85) ^f	7.91 (16.33) ^f	9.09 (17.54) ^f	1868 ^f
T ₅	0.46 ^d	0.59 ^d	0.77 ^c	0.40 ^d	0.45 ^d	0.49 ^d	11.19 (19.54) ^g	11.78 (20.07) ^g	12.04 (20.30) ^g	8.51 (16.96) ^g	8.89 (17.34) ^g	9.71 (18.15) ^g	1843 ^g

T ₆	0.32 ^b	0.47 ^b	0.66 ^b	0.23 ^b	0.30 ^b	0.37 ^b	8.42 (16.86) ^c	8.96 (17.41) ^c	10.39 (18.80) ^c	5.02 (12.94) ^c	6.27 (14.50) ^c	7.98 (16.40) ^c	1938 ^c
T ₇	0.22 ^a	0.38 ^a	0.51 ^a	0.12 ^a	0.24 ^a	0.31 ^a	7.20 (15.56) ^a	8.14 (16.57) ^a	9.50 (17.95) ^a	3.92 (11.41) ^a	5.67 (13.77) ^a	6.81 (15.12) ^a	2060 ^a
T ₈	0.27 ^a	0.44 ^b	0.56 ^a	0.19 ^b	0.27 ^a	0.33 ^a	8.32 (16.76) ^b	8.89 (17.34) ^b	10.07 (18.50) ^b	4.19 (11.81) ^b	6.05 (14.23) ^b	7.37 (15.75) ^b	1984 ^b
T ₉	1.36 ^e	1.51 ^e	1.68 ^e	1.31 ^e	1.33 ^e	1.38 ^e	14.46 (22.34) ⁱ	16.89 (24.26) ⁱ	18.45 (25.43) ⁱ	10.67 (19.06) ⁱ	12.02 (20.28) ⁱ	18.48 (25.46) ⁱ	1739 ⁱ

* Means of three replications, Values in the column followed by same letters not differ significantly by DMRT ($p=0.05$).

(T₁—Tale formulation of *T. asperellum*, as seed treatment @ 10g/kg of seed, soil application @ 10kg/ha at 25 and 45 DAS, T₂—Tale formulation of *P. fluorescens* as seed treatment @ 10g/kg of seed, soil application @ 10 kg/ha at 25 and 45 DAS, T₃—T₁+T₂, T₄—Application of fortified lignite fly ash as seed treatment @ 10g/kg of seed, soil application @ 40 kg/ha at 25 DAS, T₅—Application of Annamalai mixture as seed @ 10 ml/kg of seed, foliar spray @ 20 lit/ha at 25 and 45 DAS, T₆—T₃+T₄, T₇—T₆+T₅, T₈—Carbendazim seed treatment @ 2g/kg, foliar application @ 0.1 g/lit at 25 and 45 DAS and T₉—control.




Fig. 2. Field study of eco-friendly components against *Phacoisariopsis personata* and *Puccinia arachidis*

3. 4 DefenceDefense enzyme activity in groundnut plants to eco-friendly amendments against *P. personata* and rust *P. arachidis*

Plants rely on a robust enzymatic defense system to combat various threats, including pathogens, herbivores, and abiotic stresses. Pot culture experiments investigated the influence of various eco friendly components on defense enzyme activity in groundnut plants challenged with *P. personata* and *P. arachidis*. The eco friendly bio products included *T. asperellum*, *P. fluorescens*, fortified lignite fly ash and Annamalai mixture applied as mentioned previously. Combined application of bioproducts (treatment T7) resulted in the

Formatted: Font: Not Italic

most significant induction of ~~defense enzyme~~ activities ~~of the defense enzymes~~, PO (2.22-fold), PPO (1.28-fold) and PAL (6.25-fold). This peak induction occurred 5 days post-challenge inoculation with the respective pathogens (Fig. 2). ~~Treatment T8 also displayed substantial enzyme activity increases, although to a lesser extent than T7. Overall, enzyme activity declined following the fifth day of pathogen inoculation across all treatments.~~ Interestingly, plants solely inoculated with the pathogens also exhibited elevated levels of defense enzymes compared to the healthy control group up to the fifth day (Fig. 3). This stimulation of defense enzymes suggests that the combined application of these ~~eco-friendly biocontrol~~ components ~~possibly are involved in triggers~~ systemic resistance in groundnut plants, potentially improving their tolerance against these fungal diseases [42]. ~~These results are supported by previous studies demonstrating the induction of defense enzymes in plants treated with biocontrol agents and other eco-friendly products [43,44]. Peroxidases are a diverse group of enzymes that detoxify reactive oxygen species (ROS) produced by pathogens [45,46]. These ROS, like hydrogen peroxide (H₂O₂) and superoxide anion (O₂⁻), are highly reactive and can damage cellular components [47]. By detoxifying ROS, peroxidases prevent oxidative damage to proteins, lipids, and nucleic acids, maintaining cellular integrity and promoting plant survival[48]. In addition to ROS detoxification, plants employ Polyphenol Oxidase (PPO) to fortify their defenses. PPO catalyzes the oxidation of phenolic compounds, leading to the formation of quinones. These quinones play a multifaceted role in contributing to cell wall stiffening, triggering phytoalexin synthesis [49]. This disrupts pathogen membranes and inactivates their enzymes, directly inhibiting infection(49). Additionally, these bioproducts trigger the induced systemic resistance (ISR) in plants against pathogenic organisms[50]. Overall, these findings are consistent with previous reports suggesting that various modes of action of these eco-friendly components, including rhizosphere colonization [51], antibiotic production[52], and induction of systemic resistance[50], contribute to their potential as biocontrol agents for managing pests and diseases in crop plants. The enhanced defense enzyme activity observed in plants treated with the combination of eco-friendly components suggests their role in strengthening the plant's defense mechanisms against pathogen attack.~~



Comment [GN14]: 1,3,5,7 days. Please mention "days".

Fig. 31. Induction of defence enzymes by eco-friendly components in groundnut plants against *Phaeoisariopsis personata* and *Puccinia arachidis*. (a) Peroxidase, (b) Polyphenol oxidase, (c) Phenylalanine ammonia-lyase. T₁ — Tale formulation of *T. asperellum* seed treatment @ 10g/kg of seed, soil application @ 10 kg/ha at 25 and 45 DAS, T₂ — Tale

4. CONCLUSION

This study investigated the prevalence of Late Leaf Spot (LLS) and rust diseases in groundnut crops across the Cuddalore district, Tamil Nadu, India. The findings revealed significant geographical variation in disease incidence, with Mathur exhibiting the highest infection rates for both diseases. This suggests that location-specific factors likely play a role in disease distribution and severity. Furthermore, the research explored the efficacy of various eco-friendly components in managing these fungal diseases. A combination of *T. richoderma asperellum*, *Pseudomonas fluorescens*, fortified lignite fly ash and Annamalai mixture demonstrated the most significant reduction in disease incidence for both LLS and rust and increased groundnut yield under both pot culture and field conditions. This eco-friendly approach also resulted in increased groundnut yield compared to control groups. Additionally, the study observed induction of defense enzyme activity in groundnut plants treated with these biocontrol-based components, suggesting a potential mechanism for disease resistance. These findings highlight the potential of eco-friendly formulations for managing LLS and rust in groundnut crops, offering a sustainable alternative to chemical fungicides.

8. CONSENT

All the authors approved the manuscript at the time of submission

9. REFERENCE

1. Arya SS, Salve AR, Chauhan S. Peanuts as functional food: a review. *J. Food Sci. Technol.* 2016;53(1):31–41. doi: 10.1007/s13197-015-2007-9.
2. Pal KK, Dey R, Tilak KV. Fungal diseases of groundnut: control and future challenges. *Future challenges in crop protection against fungal pathogens.* 2014:1–29. DOI 10.1007/978-1-4939-1188-2.
3. Joshi E, Sasode DS, Singh N, Chouhan N. Diseases of groundnut and their control measures. *Biotica Res. Today.* 2020;2(5):232–237.
4. Kishore GK, Pande S. Integrated management of late leaf spot and rust diseases of groundnut (*Arachis hypogaea* L.) with *Prosopis juliflora* leaf extract and chlorothalonil. *Int. J. Pest Manag.* 2005;51(4):325–332. DOI: 10.1080/09670870500404567.
5. Subrahmanyam P, Rao VR, McDonald D, Moss JP, Gibbons RW. Origins of resistances to rust and late leaf spot in peanut (*Arachis hypogaea*, Fabaceae). *Econ. Bot.* 1989;43(4):444–455.
6. Cole DL. Rust disease of groundnut in southern Africa: present situation and possible interactions with other groundnut foliar diseases. In Citation: ICRISAT (International Crops Research Institute for the Semi-Arid Tropics). 1987. Groundnut rust disease. Proceedings of a Discussion Group Meeting, 24–28 Sep 1984, ICRISAT Center, India. Patancheru, AP 502324, India: ICRISAT;1987.
7. Goswami SK, Singh V, Chakdar H, Choudhary P. Harmful effects of fungicides- Current status. *Int. J. Agric. Environ. Biotechnol.* 2018;11:1011–1019. ISSN: 1974–1712.
8. Boudh S, Singh JS. Pesticide Contamination: Environmental Problems and Remediation Strategies. In: Bharagava R, Chowdhary P. (eds) *Emerging and Eco-Friendly Approaches for Waste Management.* Springer, Singapore. 2019:245–269. https://doi.org/10.1007/978-981-10-8669-4_12.
9. Verma DK, Pandey AK, Mohapatra B, Srivastava S, Kumar V, Talukdar D, Yulianto R, Zuan AT, Jobanputra AH, Asthir B. Plant growth-promoting rhizobacteria: An eco-friendly approach for sustainable agriculture and improved crop production. In *Microbiology for Sustainable Agriculture, Soil Health, and Environmental Protection* 2019:3–80. Apple Academic Press.
10. Raaijmakers JM, Mazzola M. Diversity and natural functions of antibiotics produced by beneficial and plant pathogenic bacteria. *Annu. Rev. Phytopathol.* 2012;50:403–424. <https://doi.org/10.1146/annurev-phyto-081211-172908>.
11. Adam M, Heuer H, Hallmann J. Bacterial antagonists of fungal pathogens also control root-knot nematodes by induced systemic resistance of tomato plants. *PloS one.* 2014;9(2):e90402. <https://doi.org/10.1371/journal.pone.0090402>.
12. Subrahmanyam P, McDonald D, Waliyar F, Reddy LJ, Nigam SN, Gibbons RW, Rao VR, Singh AK, Pande S, Reddy PM, Rao PS. Screening methods and sources of resistance to rust and late leaf spot of groundnut. *Information Bulletin no. 47.* International Crops Research Institute for the Semi-Arid Tropics; 1995: 1–20.
13. ThamamaraiSelvi M, Kurucheve V, Darwin Christdhas Henry I, Sutha Raja Kumar R and Jaiganesh V. Effect of animal excreta on mycelial growth of *Pyriculariaoryzae*. *J PharmacognPhytochem* 2019;8(3):3373–3375.
14. Newton DK, Christopher DJ. Evaluation of Bio Inoculants Fortified Lignite Fly Ash

Comment [GN15]: Please remove the references deleted from text.

Comment [GN16]: Some journals are presented in abbreviated form and some aren't. Please follow uniformity.

Comment [GN17]: The references may be kept below 30 .

- (LFA) against Sheath Blight (*Rhizoctonia solani* Kuhn) and Sheath Rot (*Sarocladiumoryzae* (Sawada) Games and Haksworth) Diseases of Rice in Cauvery Delta Region of Tamil Nadu. *Journal of Biological Control*. 2017;31(3):172–183. DOI: <https://doi.org/10.18311/jbc/2017/15848>.
15. Srivastava SK. Peroxidase and poly- phenol oxidase in *Brassica juncea* plants infected with *Macrophominaphaseolina* (Tassai) Goid. and their implication in disease resistance. *Journal of Phytopathology*. 1987;120(3):249–254. <https://doi.org/10.1111/j.1439-0434.1987.tb04439.x>.
 16. Mayer AM, Harel E. Polyphenol oxidases in plants. *Phytochemistry*. 1979;18(2):193–215. [https://doi.org/10.1016/0031-9422\(79\)80057-6](https://doi.org/10.1016/0031-9422(79)80057-6).
 17. Shaw MW, Osborne TM. Geographic distribution of plant pathogens in response to climate change. *Plant Pathology*. 2011;60(1):31–43. <https://doi.org/10.1111/j.1365-3059.2010.02407.x>.
 18. West JS, Townsend JA, Stevens M, Fitt BD. Comparative biology of different plant pathogens to estimate effects of climate change on crop diseases in Europe. *European Journal of Plant Pathology*. 2012;133:315–331. <https://doi.org/10.1007/s10658-011-9932-x>.
 19. Jain A, Sarsaiya S, Wu Q, Lu Y, Shi J. A review of plant leaf fungal diseases and its environment speciation. *Bioengineered*. 2019;10(1):409–424. <https://doi.org/10.1080/21655979.2019.1649520>.
 20. Vishwakarma K, Kumar N, Shandilya C, Mohapatra S, Bhayana S, Varma A. Revisiting plant–microbe interactions and microbial consortia application for enhancing sustainable agriculture: a review. *Frontiers in Microbiology*. 2020;11:560406. <https://doi.org/10.3389/fmicb.2020.560406>.
 21. Mondal S, Halder SK, Yadav AN, Mondal KC. Microbial Consortium with Multifunctional Plant Growth-Promoting Attributes: Future Perspective in Agriculture. In: Yadav A, Rastegari A, Yadav N, Kour D. (eds) *Advances in Plant Microbiome and Sustainable Agriculture. Microorganisms for Sustainability*, vol 20. Springer, Singapore. 2020:219–258. https://doi.org/10.1007/978-981-15-3204-7_10.
 22. Dey U, Dhutraj DN, Suryawanshi AP, DAS A, Das SP. Comparative economics of spray scheduling for the management of late leaf spot of groundnut incited by *Phaeoisariopsispersonata*. *Indian Phytopathology*. 2016;69(2):149–154.
 23. Kumhar DR, Meena AK, Meena PN. Efficacy of bio agents and fungicides against collar rot and early leaf spot of groundnut. *Indian Phytopathology*. 2018;71:549–554. <https://doi.org/10.1007/s42360-018-0080-5>.
 24. Prashar P, Kapoor N, Sachdeva S. Biocontrol of plant pathogens using plant growth promoting bacteria. *Sustainable Agriculture Reviews*: 2013;12:319–360. https://doi.org/10.1007/978-94-007-5961-9_10.
 25. Lahlali R, Ezrari S, Radouane N, Kenfaoui J, Esmael Q, El Hamss H, Belabess Z, Barka EA. Biological control of plant pathogens: A global perspective. *Microorganisms*. 2022;10(3):596. <https://doi.org/10.3390/microorganisms10030596>.
 26. Lastochkina O, Seifikalhor M, Aliniaiefard S, Baymiev A, Pusenkova L, Garipova S, Kulabuhova D, Maksimov I. *Bacillus* spp.: efficient biotic strategy to control postharvest diseases of fruits and vegetables. *Plants*. 2019;8(4):97. <https://doi.org/10.3390/plants8040097>.
 27. Nguyen SD, Trinh TH, Tran TD, Nguyen TV, Chuyen HV, Ngo VA, Nguyen AD. Combined application of rhizosphere bacteria with endophytic bacteria suppresses root diseases and increases productivity of black pepper (*Piper nigrum* L.). *Agriculture*. 2020;11(1):15. <https://doi.org/10.3390/agriculture11010015>.
 28. Muthukumar A, Eswaran A, Nakkeeran S, Sangeetha G. Efficacy of plant extracts and

- biocontrol agents against *Pythium aphanidermatum* inciting chilli damping-off. Crop protection. 2010;29(12):1483–1488. <https://doi.org/10.1016/j.cropro.2010.08.009>.
29. Akila R, Rajendran L, Harish S, Saveetha K, Raguchander T, Samiyappan R. Combined application of botanical formulations and biocontrol agents for the management of *Fusarium oxysporum* f. sp. *cubense* (Foc) causing Fusarium wilt in banana. Biological control. 2011;57(3):175–183. <https://doi.org/10.1016/j.biocontrol.2011.02.010>.
 30. Santos FM, Viera LS, Camargo DP, Muniz MF, Costa IF, Guedes JV, Santos JR, Silva JC. Integrating a Bacillus- based product with fungicides by foliar application to protect soybean: a sustainable approach to avoid exclusive use of chemicals. Pest Management Science. 2022;78(11):4832–4840. <https://doi.org/10.1002/ps.7104>.
 31. Radwan MA, Farrag SA, Abu-Elamayem MM, Ahmed NS. Biological control of the root-knot nematode, *Meloidogyne incognita* on tomato using bioproducts of microbial origin. Applied Soil Ecology. 2012;56:58–62. <https://doi.org/10.1016/j.apsoil.2012.02.008>.
 32. Pylak M, Oszust K, Fraç M. Review report on the role of bioproducts, biopreparations, biostimulants and microbial inoculants in organic production of fruit. Reviews in Environmental Science and Bio/Technology. 2019;18(3):597–616. <https://doi.org/10.1007/s11157-019-09500-5>.
 33. Ali O, Ramsubhag A, Jayaraman J. Biostimulant properties of seaweed extracts in plants: Implications towards sustainable crop production. Plants. 2021;10(3):531. <https://doi.org/10.3390/plants10030531>.
 34. Khalid A, Arshad M, Shaharoon B, Mahmood T. Plant Growth Promoting Rhizobacteria and Sustainable Agriculture. In: Khan M, Zaidi A, Musarrat J. (eds) Microbial Strategies for Crop Improvement. Springer, Berlin, Heidelberg. 2009:133–160. https://doi.org/10.1007/978-3-642-01979-1_7.
 35. Maheshwari DK, Dheeman S, Agarwal M. Phytohormone-Producing PGPR for Sustainable Agriculture. In: Maheshwari D. (eds) Bacterial Metabolites in Sustainable Agroecosystem. Sustainable Development and Biodiversity, 12. Springer, Cham. 2015:159–182. https://doi.org/10.1007/978-3-319-24654-3_7.
 36. Ram LC, Srivastava NK, Tripathi RC, Jha SK, Sinha AK, Singh G, Manoharan V. Management of mine spoil for crop productivity with lignite fly ash and biological amendments. Journal of environmental management. 2006;79(2):173–187. <https://doi.org/10.1016/j.jenvman.2005.06.008>.
 37. Kishor P, Ghosh AK, Kumar D. Use of fly ash in agriculture: A way to improve soil fertility and its productivity. Asian Journal of Agricultural Research. 2010;4(1):1–4. ISSN: 1819–1894.
 38. Alterary SS, Marei NH. Fly ash properties, characterization, and applications: A review. Journal of King Saud University-Science. 2021;33(6):101536. <https://doi.org/10.1016/j.jksus.2021.101536>.
 39. Hajiboland R. Effect of Micronutrient Deficiencies on Plants Stress Responses. In: Ahmad P, Prasad M. (eds) Abiotic Stress Responses in Plants. Springer, New York, NY. 2012:283–329. https://doi.org/10.1007/978-1-4614-0634-1_16.
 40. Ali N, Réthoré E, Yvin JC, Hosseini SA. The regulatory role of silicon in mitigating plant nutritional stresses. Plants. 2020;9(12):1779. <https://doi.org/10.3390/plants9121779>.
 41. Khan I, Awan SA, Rizwan M, Brestic M, Xie W. Silicon: an essential element for plant nutrition and phytohormones signaling mechanism under stressful conditions. Plant Growth Regulation. 2023;100(2):301–319. <https://doi.org/10.1007/s10725-022-00872-3>.

42. Andrés JA, Pastor NA, Ganuza M, Rovera M, Reynoso MM, Torres A. Biopesticides: An Eco-Friendly Approach for the Control of Soilborne Pathogens in Peanut. In: Singh D, Singh H, Prabha R. (eds) *Microbial Inoculants in Sustainable Agricultural Productivity*. Springer, New Delhi. 2016;1:161–179. https://doi.org/10.1007/978-81-322-2647-5_9.
43. Rais A, Jabeen Z, Shair F, Hafeez FY, Hassan MN. *Bacillus* spp., a bio-control agent enhances the activity of antioxidant defense enzymes in rice against *Pyriculariaoryzae*. *PloS one*. 2017;12(11):e0187412. <https://doi.org/10.1371/journal.pone.0187412>.
44. Omidvari M, Abbaszadeh-Dahaji P, Hatami M, Kariman K. Biocontrol: a novel eco-friendly mitigation strategy to manage plant diseases. *Plant Stress Mitigators*. 2023;27–56. <https://doi.org/10.1016/B978-0-323-89871-3.00020-3>.
45. Kawano T. Roles of the reactive oxygen species-generating peroxidase reactions in plant defense and growth induction. *Plant Cell Rep. Plant cell reports*. 2003;21:829–837. <https://doi.org/10.1007/s00299-003-0591-z>.
46. Almagro L, Gómez Ros LV, Belchi-Navarro S, Bru R, Ros Barceló A, Pedreño MA. Class III peroxidases in plant ~~defence~~defense reactions. *Journal of experimental botany*. 2009;60(2):377–390. <https://doi.org/10.1093/jxb/ern277>.
47. Madkour LH. Function of reactive oxygen species (ROS) inside the living organisms and sources of oxidants. *Pharm. Sci. Anal. Res. J*. 2019;2:180023. ISSN: 2640–6659.
48. Sharma P, Sharma P, Arora P, Verma V, Khanna K, Saini P, Bhardwaj R. Role and regulation of ROS and antioxidants as signaling molecules in response to abiotic stresses. In *Plant signaling molecules*, Woodhead Publishing; 2019.
49. Jamiolkowska A. Natural compounds as elicitors of plant resistance against diseases and new biocontrol strategies. *Agronomy*. 2020;10(2):173. <https://doi.org/10.3390/agronomy10020173>.
50. Nguvo KJ, Gao X. 2019. Weapons hidden underneath: bio-control agents and their potentials to activate plant induced systemic resistance in controlling crop *Fusarium* diseases. *J. Plant Dis. Prot*. 2019;126(3):177–190.
51. Yang X, Chen L, Yong X, Shen Q. Formulations can affect rhizosphere colonization and biocontrol efficiency of *Trichoderma harzianum* SQR-T037 against *Fusarium* wilt of cucumbers. *Biol Fertil Soils*. 2011;47:239–248. <https://doi.org/10.1007/s00374-010-0527-z>.
52. Ulloa-Ogaz AL, Muñoz-Castellanos LN, Nevárez-Moorillón GV. Biocontrol of phytopathogens: antibiotic production as mechanism of control. *The battle against microbial pathogens: basic science, technological advances and educational programs*. 2015;1:305–309.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

Formatted: Normal, Left, Indent: Left: 0.25", Right: 0", No bullets or numbering, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Formatted: Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto

Formatted: Font: (Default) Times New Roman, 12 pt