

TRAUMA-RELATED DEATHS IN THE ACCIDENT AND EMERGENCY DEPARTMENT OF A NIGERIAN TEACHING HOSPITAL: A retrospective study

Abstract:

Background: Trauma is a significant contributor to morbidity, disabilities, loss of man hours and mortality in both the developed and developing world with 90% of global trauma deaths occurring in low and middle income countries. The extent of motor vehicular use, degree of adherence to use of safety gadgets, strength of enforcement of industrial safety regulations, level of civil unrest as well as other socio-demographic indices predict the contribution of various etiologic factors. Deaths from the trauma may perhaps have far more psychologic and physical impact on families that from acute/chronic medical cases

Method:: A retrospective study of the admission records and in-patient records of all trauma related deaths in the accident and emergency unit of the University of Port Harcourt Teaching Hospital from 1st January 2018 to 31st December 2022 was done. Relevant information was extracted and analyzed. Data obtained was analyzed descriptively. Results were presented in tables and charts where necessary. Means and standard deviations were used to represent certain variables.

Results: Total emergencies (medical and surgical) seen within the study period were 42,443, total injuries seen were 10,787 giving a crude injury prevalence rate of 25.4%. There were a total of 766 deaths among injured patients giving a crude mortality rate of 7.1%. Road traffic accidents (RTAs) (46.3%) and Firearm injuries (23.5%) were the most common injury mechanisms seen. Traumatic brain injuries are by far the leading cause of trauma-related deaths in the ER (82.5%) with most deaths (96.1%) occurring within the first 72hours of Admission.

Conclusion: Trauma-related deaths occur in 7.1% of all injuries presenting at the study sight. The most common mechanism is RTA while the most affected victim is the young adult male. Traumatic brain injury accounts for more deaths than all others combined with most of the deaths occurring within the first 72hours of presentation. All efforts at reducing these injuries should be encouraged.

Key word: Trauma, disabilities, public health crisis, domestic injuries

INTRODUCTION

Trauma is a significant contributor to morbidity, disabilities, loss of man hours and mortality in both the developed and developing world. Ninety percent of global trauma deaths occur in low and middle income countries¹. Trauma in general contribute to 12% of the global disease burden². It has been regarded as a neglected public health crisis in low and middle income countries³

Common causes include, Road traffic accidents (RTAs), fall from height, industrial injuries, occupational injuries, domestic injuries, assault and firearm injuries. The extent of motor vehicular use, degree of adherence to use of safety gadgets, strength of enforcement of industrial safety regulations, level of civil unrest as well as other socio-demographic indices predict the contribution of various etiologic factors.

Road traffic accidents are by far the most common cause of both trauma- related injuries and trauma-related deaths⁴⁻⁷. The National Bureau of Statistics in Nigeria reported an increase in road traffic crashes (RTC) by approximately 14% from 2013 to 2019⁸. These injuries typically affects the young adult male who in most part of the developing world is the most dynamic, most enterprising and most daring⁹⁻¹⁰. Falls from heights are particularly a major concern in the children and in the elderly population. Firearm injuries have also been reported by some authors^{6,10} as growing contributors to injuries. The degree to which firearm injuries contribute to trauma depend on the gun control measures and the presence or otherwise of conflicts within the region served by the care facility⁴.

In low and middle income countries, trauma systems- a structured framework that provides optimal pre-hospital, intra-hospital and post treatment rehabilitation for the injured patients through appropriate co-ordination of health resources, health care personnel and healthcare facilities- is still at the rudimentary stage^{11,12}. This creates a serious intervention gap and could contribute to the poorer outcome of care seen in these regions compared to the more developed world¹³.

In addition, available trauma data are not centralized, prehospital care is almost non-existent and prompt/ optimal intra-hospital care is still hindered by poor health insurance coverage and lack of needed resources¹⁴⁻¹⁶. These may further worsen outcome indices¹⁵.

Disability-adjusted life years (DALYs) which combine the number of years of life lost from premature death with the loss of health from disability among persons with nonfatal injuries is an attempt to measure the non-fatal impact of trauma on a given population¹⁷.

Deaths from the trauma may perhaps have far more psychologic and physical impact on families than from acute/chronic medical cases. The demography of those affected explains the enormity of its impact on Gross domestic product of any nation and the social and mental health implications on family members left behind.

A study of the mortality pattern of such injuries will among other things help detect any change in epidemiologic trend, understand the common injuries that account for trauma deaths and appreciate the demographic distributions with the view of improving the management of trauma within the study area and beyond.

AIM: The aim of this study was to profile trauma-related deaths at the accident and emergency department of the University of Port Harcourt Teaching Hospital.

MATERIALS AND METHODS

STUDY DESIGN: this was a retrospective study of deaths from trauma patients that presented at the accident and emergency department of the University of Port Harcourt Teaching Hospital over a five year period 1st January 2018- 31st December 2022.

METHOD: The admission records and in-patient records were retrospectively reviewed to extract relevant information on patient's demographics, presenting complaint at admission, treatment spectrum, Length of stay in the ward, emergency treatment received and the specialists for definitive treatment. For the purpose of analysis, patient comorbidity was defined as the presence of pre-existing medical conditions that could alter the outcome of treatment of the traumatic condition. Deaths recorded in this study were those that occurred while the patients were receiving care at the Accident and emergency department. Deaths among trauma patients transferred to other units of the hospital were not included.

Trauma patients brought in dead and those who died on arrival were also excluded from the study. For the purpose of analysis, dead-on-arrival (DOA) refers to patients with severe traumatic conditions who died at the emergency room before commencement of any form of intervention while brought in dead patients were regarded as those who were already dead before arrival at the study site.

STUDY LOCATION

The University of Port Harcourt teaching Hospital (UPTH) is in Rivers State along East-West Road with coordinates of 4.45305800N and 6.5504300E. UPTH serves as a tertiary

referral center and receives referrals from neighboring states such as Bayelsa, Abia, Imo, Akwa-Ibom, Delta, Cross River, and other states in Nigeria. The accident and emergency department is the first point of call for all adult and pediatric trauma emergencies.

The department has medical officers of several categories and varying level of training who provide emergency care for trauma patients before transfer to the more appropriate specialty for definitive treatment. More than 50% of the medical and nursing staff do not have requisite certification in Basic life support (BLS) Advanced cardiovascular life support (ACLS) and Advanced trauma life support (ATLS)

Payment for elective interventions are on out-of-pocket basis for most patients since health insurance coverage is quite low⁶. The emergency care policy of the hospital provides some cover for emergency procedures but drugs and advanced surgical interventions are not included in this policy.

STUDY POPULATION

Patients who presented at the accident and emergency department of the study center within the study period. Patients with incomplete medical records that will not aid in analysis were excluded.

DATA ANALYSIS

Data obtained were analyzed descriptively. Results were presented in tables and charts where necessary. Means and standard deviations were used to represent certain variables. Chi-square test and students' T-test for continuous variables were used where appropriate to test observed differences, and p-value of <0.05 was deemed statistically significant

RESULTS

A total of 10, 939 patients presented to the accident and emergency department with traumatic injuries. Records of 152 patients were either not completely reported or not found in the archives. Available records from 10,787 were recruited into the study giving 98.6% recruitment rate. Total emergencies (medical and surgical) seen within the study period were 42,443 giving a crude injury prevalence rate of 25.4%

The age range for all trauma patients within the study period was from 0-96 years with a mean age of 33.17+/-16.1years. The most common age group was the 21-30 year age group. There were 8,342 males and 2,445 females making a male female ratio of 3.4:1

Table 1: Demographics

Age (years)	n (%)
<1	23 (0.2)
1-10	302 (2.7)
11-20	924 (8.6)
21-30	3941 (36.6)
31-40	2523 (23.4)
41-50	2086 (19.3)
51-60	719 (6.7)
61-70	117 (1.1)
71-80	82 (0.7)
80-90	53 (0.5)
>90	17 (0.2)
Sex	
Male	8342 (77.3)
Female	2445 (22.7)
Occupation	
preschool Children/students	1323 (12.2)
Driver	508 (4.7)
Civil servant/ self -employed	2409 (22.3)
Unemployed	5638 (52.3)
Businessman	3981 (36.9)
Professionals	349 (3.2)
Others	60 (0.6)
Marital status	
Single	4434 (41.1)

Married	5969 (55.3)
Widow	306 (2.8)
Divorced	78 (0.7)
Level of education	
Un-educated	228 (2.1)
Primary	714 (6.6)
Secondary	5717 (53.0)
Tertiary	4128 (38.3)

Road traffic accidents (RTAs) accounted for 46.3% of all injury mechanisms seen. Firearm injuries (23.5%) burns (8.7%) and assault (6.0%) also made good contributions as stated in table 2.

Table 2: Distribution of etiologies for all trauma patients

Etiology	n (%)
Road Traffic Accidents (RTA)	4993 (46.3)
Firearm injuries	2531 (23.5)
Burns	934 (8.7)
Falls	643(6.0)
Sports injuries	244 (2.3)
Industrial injuries	786 (7.3)
Assaults	637 (6.0)
Birth trauma	19 (0.1)
Total	10,787 (100)

Table 3: Distribution of injury mechanisms across age groups for all trauma patients.

Age group (years)	Etiology								Total
	RTA	Falls	Firearms	Burns	Sports injuries	Industrial injuries	Assaults	Birth trauma	
<1	3	1	-	-	-	-	-	19	23

Age group (years)	Etiology								Total
	RTA	Falls	Firearms	Burns	Sports injuries	Industrial injuries	Assaults	Birth trauma	
1-10	29	108	27	34	60	-	44	-	302
11-20	641	89	5	54	48	-	87	-	924
21-30	1501	21	1418	329	83	283	306	-	3941
31-40	1412	38	425	117	53	352	126	-	2523
41-50	1047	81	434	358	-	130	36	-	2086
51-60	327	99	217	31	-	21	24	-	719
61-70	18	79	3	9	-	-	8	-	117
71-80	12	63	1	2	-	-	4	-	82
80-90	1	50	-	-	-	-	2	-	53
>90	2	14	1	-	-	-	-	-	17
Total	4993	643	2531	934	244	786	637	19	10,787
(%)	(46.8)	(6.0)	(23.5)	(8.7)	(2.3)	(7.3)	(6.0)	(0.1)	(100)

Skin and subcutaneous injuries (39.5%) were the most commonly injured body region, followed by bone fractures (31.6%).

Table 4: Distribution of Body regions affected

Injury pattern	<i>n</i> (%)
Bone fractures	3,405 (31.6)
Skin and subcutaneous Injuries	4,261 (39.5)
Traumatic brain injuries	1894 (17.6)
Spinal cord injuries	910 (8.4)
Maxillo-facial injuries	493(4.6)
ENT injuries	676 (6.3)
Ophthalmic injuries	390 (3.6)
Urologic injuries	341 (3.2)
Chest trauma	681 (6.3)

Injury pattern	<i>n</i> (%)
Abdomino-pelvic injuries	108(1.0)
Multiple injuries	1,994 (18.5)

Most patients were stabilized and transferred to the ward/ theater/ICU (63.7%) while 21.7% of patients were discharged home after initial stabilization as shown in table 5.

Table 5: outcome of ER treatment

ER treatment outcome	<i>n</i> (%)
Stabilized and discharged home	2342 (21.7)
Stabilized and transferred to the ward/theatre/ICU	6867 (63.7)
Signed against Medical advice	249 (2.3)
Absconded	115(1.1)
Referred out by specialist team	445 (4.1)
Died in the ER	766 (7.1)
Total	10,787 (100)

Road traffic accidents (68.3%) and firearm injuries (15.3) made the highest contribution to mortality.

Table 6: Distribution of mortality pattern across injury mechanism

Etiology	<i>n</i> (%)
Road Traffic Accidents (RTA)	523 (68.3)
Firearm injuries	117 (15.3)
Burns	22 (2.8)
Falls	14(1.8)

Etiology	<i>n</i> (%)
Industrial injuries	19 (2.5)
Assaults	11 (1.4)
Total	766 (100)




Figure1: distribution of mortality pattern across five years




Figure 2: distribution of mechanisms of mortality.

Table 7: Distribution of mortality against duration of stay in the ER

	Duration of hospital stay		
	<24hours	24-72hours	>72hours
Severe TBI	458 (59.8%)	156 (20.4%)	20 (2.6%)
Hemorrhagic shock	88 (11.5%)	3 (0.4%)	-
Spinal cord injury	12 (1.6%)	16 (2.1%)	2 (0.2%)
Septic shock	-	-	8 (1.0%)
Others	1 (0.1%)	2 (0.2%)	-
Total	559 (73%)	177 (23.1%)	30(3.9%)

DISCUSSION

Results from this study showed that 42,443 patients presented to the Accident and emergency department (emergency room) within the five year period under review. Of this number 10,787 were injuries giving a crude injury prevalence rate of 25.4%. Both Onyemachi et al¹⁸ in Enugu (24.3%) and Prekker *et al*¹⁰(24.4%) in Minnesota, USA reported similar findings.

The crude mortality rate was 7.1%. This is lower than findings from Elachi et al⁴ in Markurdi North central Nigeria (15.2%) and higher than findings from other authors within the region including Thanni and Kehinde⁵ (2%) for south-west Nigeria, Solegberu et al¹⁶(4.4%) from Lagos, south-west Nigeria, Onyemaechi et al¹⁸ (4.5%) from Enugu, South-East Nigeria, as well as Egol et al¹⁹(4.7%) from USA. The rate disparities may be a reflection of the geographic disparity in the spread of competent emergency and trauma care specialist within the region, the distribution of injury mechanisms as well as the quality of care available at various centers. This disparity also highlights the need for a central national trauma database which among other benefits have the potential of revealing the existing health infrastructure and health man-power gaps across different tertiary health facilities in the country.

The most common victim of trauma and trauma-related deaths is the adult male of the 21-40 years age group. The young adult male have been shown by several studies^{2,4,5} as the most vulnerable to injuries and deaths from injuries. This group of patients constitute the large chunk of the demographics of young populations typical of low and middle-income countries and are the most dynamic and most productive fraction of the population in these countries²⁰⁻²². Their loss therefore is not merely a subtraction from the population but a huge bite on the nation's productive capacity and potentially a reduction in its gross earnings.

Road traffic accidents accounted for far more deaths (68.3%) than all other injury mechanism combined. Most authors²³⁻²⁵ have reported the leading role of road traffic accidents in trauma-related injuries and deaths. The fraction of its contribution being a mirror image of the level of urbanization, the extent of motor vehicular use and the degree to which safe driving policies and helmet use is adhered to²⁶. Though existent within cities served by the study center, safe driving laws are flagrantly disobeyed by drivers and other road-users, helmet use is poorly implemented and available road networks are in terribly poor conditions.

This study particularly showed a high incidence of deaths from firearm injuries (15.3%) in the region much higher than those from falls, burns, Industrial accidents and assaults put together. This is a clear reflection of the high rate of gun violence in the oil-rich

regions of the Niger delta and perhaps the unguided use of weapons by security personnel in the bid to maintain law and order. This also underscores the need to curb the proliferation of illegal weapons and train security personnel on proper use of weapons within their reach.

Distribution of mortality across the five years of the study showed that the number of deaths were lower in 2020 and in 2022 compared to the other three years. The former may have been due to the covid-19 restrictions which limited vehicular movements and crashed trauma incidences and trauma deaths while the latter may be explained by the establishment of a trauma team in the study center earlier that year. The trauma team, though still at its evolving stage was a collection of specialized trauma residents from surgery and related departments pulled to the Emergency room to provide competent first-hand treatment to trauma patients with the aim of achieving prompt resuscitation and stabilization before definitive treatment is carried out by the specialist teams. Its dramatic impact on the death rates for trauma-related injuries clearly shows the role of deliberate and intervention-focused management policies in reducing mortalities and improving the quality of care.

Traumatic brain injuries are by far the leading cause of trauma related deaths in our ER (82.5%). Amaefule et al²⁷ also showed that traumatic brain injuries caused more deaths (66.2%) within the first 72hours in the ER than all other causes combined. Etebu and Ekere²⁸ in a similar study 2 decades ago in the same institution showed that traumatic brain injury was the leading cause of trauma-related deaths in children. The persistently high impact of head injuries to trauma-related deaths may result from poor adherence to helmet use, terrible condition of roads within the country, the abysmally low ratio of neurosurgeon per population (worsened by health workforce exports) as well as inability of a large fraction of the population to afford brain CT scan^{29,30,31}.

Hemorrhagic shock accounted for 12% of mortalities within the study period. These deaths were probably from unrecognized concealed bleed coupled with poor assessment of hemodynamic status with inadequate intervention in the ER. Reducing mortalities from hemorrhagic shock will require more attention to clinical details by the ER personnel and a deliberate management policy aimed at prompt and unrestricted access to blood for trauma victims without payment irrespective of health insurance status. Reduction of existing bottlenecks to early surgical intervention will also go a long way to reducing mortality from hemorrhagic shock.

More than 95% of trauma-related deaths took place within the first 72hours of admission. Both Osime et al³¹ in a similar tertiary center in Nigeria and Elachi et al⁴ have reported the

same pattern, clearly reflecting the severity of such injuries at presentation and the need for emergency room personnel to optimally utilize this golden period for prompt and adequate resuscitation.

Reducing trauma-related mortality clearly need an ambivalence of coordinated efforts from vehicle manufacturers, drivers/road users, prehospital personnel, the ER physicians, health system managers and the government at all levels. The enforcement of safe driving laws such as helmet use, use of seat belts, government-backed vehicle inspection, speed restrictions, no drink-driving policies, safer vehicle manufacturing as well as provision of good roads are some primordial and primary preventive measures³³.

Secondary preventive measures will include, prompt evacuation and transport of injured victims to appropriate facilities within an organized trauma system, adequate prehospital resuscitation as well as prompt and optimal in-hospital care by Advance trauma life support trained ER personnel.

As tertiary measures, early and adequate rehabilitation of injured victims will go a long way to incorporating these victims back to work and social life as early as possible

Though most of these are stills non-existent in the country, this study will hopefully stimulate the needed change.

CONCLUSION:

Trauma-related deaths occur in 7.1% of all injuries presenting at the study sight. The most common mechanism is RTA while the most affected victim is the young adult male. Traumatic brain injury accounts for more deaths than all others combined with most of the deaths occurring within the first 72hours of presentation. Since these deaths are preventable, all effort should be on deck to prevent or at least reduce them.

Ethical Approval:

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

REFERENCES

1. Gosselin RA, Spiegel DA, Coughlin R, Zirkle LG. Injuries: The neglected burden in developing countries. *Bull World Health Organ* 2009;87:246–246a.
2. Peden M, McGee K, Sharma G. *The Injury Chart Book: A Graphical Overview of the Global Burden of Injuries*. Geneva, Switzerland: World Health Organization; 2002. Available from: <http://www.who.int/publications/924156220x.pdf>
3. Hyder AA. Injuries in low- and middle-income countries: a neglected disease in global public health. *Injury*, 44 (5) (2013), pp. 579-580.
4. Elachi IC, Yongu WT, Odoyoh OO, Mue DD, Ogwuche EI, Ahachi CN, et al. An epidemiological study of the burden of trauma in Makurdi, Nigeria. *Int J Crit Illn Inj Sci* 2015;5:99–102. [
5. Thanni LO, Kehinde OA. Trauma at a Nigerian teaching hospital: Pattern and documentation of presentation. *Afr Health Sci* 2006;6:104–7.
6. Chalya PL, Dass RM, Mchembe MD, Mbelenge N, Ngayomela IH, Chandika AB, et al. Citywide trauma experience in Mwanza, Tanzania: A need for urgent intervention. *J Trauma Manag Outcomes* 2013;7:9
7. Tadesse B, Tekilu S, Nega B, Seyoum N. Pattern of injury and associated variables as seen in the emergency department at tikuranbessa specialized referral hospital; Adis Ababa Ethiopia. *East Cent Afr J Surg* 2014;19:73–82.
8. Reports | National Bureau of Statistics [Internet]. [cited 2021 Oct 13]. Available from: <https://nigerianstat.gov.ng/elibrary/read/1164>
9. Lagarde E Road traffic injury is an escalating burden in Africa and deserves proportionate research efforts. *PLoS Med* 2007;4:e170.
10. Prekker ME, Miner JR, Rockswold EG, Biros MH. The prevalence of injury of any type in an urban emergency department population. *J Trauma* 2009;66:1688–95.
11. Howard JM. Historical background to accidental death and disability: the neglected disease of modern society. *Prehosp Emerg Care*, 4 (4) (2000), pp. 285-289
12. GJ Bazzoli, EJ. MacKenzie. Trauma centers in the United States: identification and examination of key characteristics. *J Trauma Acute Care Surg*, 38 (1) (1995), pp. 103-110
13. JC He, LA Kreiner, N Sajankila, DL Allen, JA. Claridge. Performance of a regional trauma network: a state-wide analysis. *J Trauma Acute Care Surg*, 81 (1) (2016), pp. 190-195
14. A Gwaram, OG Okoye, OO Olaomi. Observed benefits of a major trauma centre in a tertiary hospital in Nigeria. *Afr J Emerg Med*, 11 (2) (2021), pp. 311-314
15. D. Adeloje. Prehospital trauma care systems: potential role toward reducing morbidities and mortalities from road traffic injuries in Nigeria. *Prehospital Disaster Med*, 27 (6) (2012), pp. 536-

16. BA Solagberu, AO Adekanye, CPK Ofoegbu, US Udoffa. Epidemiology of trauma deaths. *West Afr J Med*, 22 (2) (2003), pp. 177-185
17. Krug EG, Sharma GK, Lozano R. The global burden of injuries. *Am J Public Health*. 2000;90:523–6.
18. Onyeamachi NO, Nwankwo OE, Ezeadawi RA. Epidemiology of Injuries Seen in a Nigerian Tertiary Hospital. *Niger J ClinPract*. 2018 Jun; 21(6): 752–757.
19. Egol KA, Tolisano AM, Spratt KF, Koval KJ. Mortality rates following trauma: The difference is night and day. *J Emerg Trauma Shock* 2011;4:178–83.
20. Swarnkar M, Singh P, Dwivedi S. Pattern of trauma in central India: An epidemiological study with special reference to mode of injury. *Internet J Epidemiol*. 2009;9.
21. Banthia P, Koirala B, Rauniyar A, Chaudhary D, Kharel T, Khadka SB. An epidemiological study of road traffic accident cases attending emergency department of teaching hospital. *JNMA J Nepal Med Assoc*. 2006;45:238
22. Mohammed AZ, Edino ST, Ochicha O, Umar AB. Epidemiology of gunshot injuries in Kano, Nigeria. *Niger J Surg Res*. 2005;7:296–9
23. Abhilash KP, Chakraborty N, Pandian GR, Dhanawade VS, Bhanu TK, Priya K. Profile of trauma patients in the emergency department of a tertiary care hospital in South India. *J Family Med Prim Care*. 2016 Jul-Sep;5(3):558-563.
24. Rastogi D, Meena S, Sharma V, Singh GK. Epidemiology of patients admitted to a major trauma centre in northern India. *Chin J Traumatol*. 2014;17:103–7.
25. Shameem AM, Shabbir KM, Agrawal D, Sharma BS. Outcome in head injured patients: Experience at a level 1 trauma centre. *Indian J Neurotrauma*. 2009;6:119–22.
26. Ernstberger A, Joeris A, Daigl M, Kiss M, Angerpointner K, Nerlich M, Schmucker U. Decrease of morbidity in road traffic accidents in a high income country - an analysis of 24 ,405 accidents in a 21 year period. *Injury*. 2015 Oct;46 (Suppl 4):S135-143.
27. Amaefule KE, Dahiru IL, Ejagwulu FS, Maitama MI. Trauma Mortality in The Emergency Department of a Tertiary Hospital in a Low-Income Country: It's Time To Walk The Talks. *West African Journal of Medicine*. 2020 Apr-Jun;37(2):131-137. PMID: 32150631
28. Etebu EN, Ekere AU. Paediatric accidental deaths in Port Harcourt, Nigeria: a 10-year retrospective study. *Niger J Med*. 2004 Apr-Jun;13(2):140-143.
29. Liu BC, Ivers R, Norton R, Boufous S, Blows S, Lo SK. Helmets for preventing injury in motorcycle riders. *Cochrane Database Syst Rev*. 2008 Jan 23;(1):CD004333.
30. Siddiqui AA, Zafar H, Bashir SH. An audit of head trauma care and mortality. *J Coll Physicians Surg Pak*. 2004 Mar;14(3):173-177

31. Shekhar C, Gupta LN, Premsagar IC, Sinha M, Kishore J. An epidemiological study of traumatic brain injury cases in a trauma centre of New Delhi (India). *J Emerg Trauma Shock*. 2015 Jul-Sep;8(3):131-139
32. Osime OC, Ighedosa SU, Oludiran OO, Iribhogbe PE, Ehikhamenor E, Elusoji SO. Patterns of trauma deaths in an accident and emergency unit. *Prehosp Disaster Med*. 2007 Jan-Feb;22(1):75-78.
33. Kleber C, Giesecke MT, Tsokos M, Haas NP, Buschmann CT. Trauma-related preventable deaths in Berlin 2010: need to change prehospital management strategies and trauma management education. *World J Surg*. 2013 May;37(5):1154-1161.
34. Nzegwu MA, Aligbe JU, Banjo AA, Akhiwui W, Nzegwu CO. Patterns of morbidity and mortality amongst motorcycle riders and their passengers in Benin-city Nigeria: One-year review. *Ann Afr Med*. 2008;7:82-5
35. Murad MK, Larsen S, Husum H. Prehospital trauma care reduces mortality. Ten-year results from a time-cohort and trauma audit study in Iraq. *Scand J Trauma, Resusc and Emerg Med*. 2012, 20:13

UNDER PEER REVIEW