

Case report

Intestinal occlusion on flange taking Meckel's diverticulum: about a case report

Abstract

Meckel's diverticulum is a congenital anomaly of the digestive tract resulting from the persistence of the omphalomesenteric duct, and is often asymptomatic and discovered by chance. However, it can sometimes lead to complications. We report the case of a 30-year-old oncology patient with sigmoidal adenocarcinoma and liver metastasis who presented with intestinal obstruction due to a gregolepipoic bridle taking Meckel's diverticulum.

Key words: meckel's diverticulum, intestinal obstruction

Introduction

Meckel's diverticulum (MD) is a persistent defect of the omphalomesenteric duct, first described in 1598 [1]. It is the most common congenital malformation of the digestive tract. Its incidence is between 1% and 4% of the general population [2]. Generally benign and asymptomatic, DM is a pathology of childhood, but can manifest itself in adulthood. Complicated forms, such as digestive haemorrhage, intestinal obstruction and intestinal intussusception, account for 4 to 16% of DM cases, and often constitute the circumstances of their discovery [1-3]. The aim of this article was to describe the clinical case and management of an intestinal obstruction secondary to a Meckel diverticulum.

Case Presentation

This is a 30-year-old oncology patient with a sigmoidal adenocarcinoma for which he had undergone a sigmoidal colostomy and three sessions of chemotherapy. His current history dates back to 24 hours before admission, with

the onset of an occlusive syndrome involving cessation of bowel movements and gas, bilious vomiting and no other associated signs. All this evolved in a context of apyrexia and altered general condition. Clinically, the patient was conscious (Glasgow score 15/15), hemodynamically stable (BP=12/08cmhg) and respiratory (FR=18cycles/min). Abdomen was distended, tympanic, stoma non-functional. Stomal and rectal examination were without abnormalities. An abdominal CT scan (Fig. 1) showed distension of some of the bowel loops, measuring 38 mm in maximum diameter, with local hydroaerosal levels, contrasting with a collapsed appearance of the colonic frame and ileal loops upstream of a transitional level opposite L5.




Figure 1 Abdominopelvic CT scan, showing distension and hydroaeric level

The patient underwent surgery for section of a gremoepiploic flange taking up Meckel's diverticulum, with wedge-shaped bowel resection of Meckel's diverticulum and exploration= Presence of epiplo-parietal, gremo-epiploic and gremo-grelic adhesions (adhesiolysis performed). Presence of a gremo-epiploic flange 50cm from the ICJ responsible for 4cm of distension upstream, taking up Meckel's diverticulum located 50cm from the ICJ, which is in pain.



Figure 2 Meckel's diverticulum (intraoperative photo)

Discussion

Meckel's diverticulum is the partial persistence of the omphalomesenteric duct. It is the most common congenital anomaly of the gastrointestinal tract, with a slight male predominance [4,5]. It is rare, occurring in between 2% and 4% of the population [6,4]. Meckel's diverticulum is usually asymptomatic, and is only diagnosed incidentally or when complications arise. The diagnosis of intestinal obstruction due to Meckel's diverticulum can be evoked preoperatively, either on abdominal ultrasound, technetium-99m scintigraphy, abdominal computed tomography or magnetic resonance imaging (MRI) [7]. Distinguishing between an intestinal loop and Meckel's diverticulum on CT remains difficult [8]. Mechanical obstruction is the most frequent complication in adults, accounting for 24 to 53%. Most often, this is an occlusion with a variable mechanism [9,10,11]: volvulus, invagination, fixation of the diverticulum at the umbilicus or at any other point in the abdomen. The frequency of complications is slightly higher in men [4,5]. In this case, the Meckel diverticulum was located 50cm from Bauhin's valve, with a diameter of 2cm and a length of 6cm, and a flange was involved in the occlusion. The location of Meckel's diverticulum varies between 10 and 100 cm from Bauhin's valve in 50% of cases, and its dimensions average 2cm in diameter and 5cm in length [5]. When a Meckel's diverticulum is discovered, surgical treatment is often required. Two methods are commonly reported in the literature: wedge-shaped resection (diverticulectomy) and segmental bowel resection involving Meckel's diverticulum and terminal-graft anastomosis [12], although the latter is preferred to the former. In our case, we opted for a wedge-shaped bowel resection of the meckel diverticulum, with section of an epiploic bridle taking up the diverticulum. However, in the event of a complication, segmental resection of the diverticulum is the rule. This is preferred to wedge-shaped resection, which risks leaving ectopic tissue in place [13]. Laparoscopy is playing an increasingly important role in these procedures. Regardless of the approach used, digestive sutures must be performed on a perfectly healthy, non-inflammatory and well-vascularized area [14].

Conclusion:

Meckel's diverticulum is the partial persistence of the omphalomesenteric duct. It is usually asymptomatic. Complicated Meckel's diverticulitis is mainly diagnosed intraoperatively. It is a surgical emergency requiring early and appropriate management. The outcome depends on how early the diagnosis is made.

References

1.Park JJ, Wolff BG, Tollefson MK, Walsh EE, Larson DR. Meckel Diverticulum: the Mayo Clinic Experience with 1476 Patients (1950-2002). *Ann Surg.* 2005 Mar;241(3):529-33. PubMed | Google Scholar

2.Bani-Hani KE, Shatnawi NJ. Meckel's Diverticulum: comparison of incidental and symptomatic cases. *World J Surg.* 2004 Sep;28(9):917-20. PubMed | Google Scholar

3.Mendelson KG, Bailey MB, Balint TD, Pofahl WE. Meckel's Diverticulum: review and surgical management. *Current Surgery.* 2001;58:455-7. PubMed | Google Scholar

4.Barbary C, Tissier S, Floquet M, Régent D. Imaging complications of complications of Meckel's diverticulum. *J Radiol.* March 1 2004;85(3):273-9. PubMed | Google Scholar

5.Ouangré E, Zida M, Bazongo M, Sanou A, Patindé Bonkoungou G, Doamba ND et al. Complications of Meckel's diverticulum (DM) in adults. in adults: about 11 cases at CHU-Yalgado Ouédraogo in Burkina Faso. *Pan Afr Med J.* 2015;22:74. PubMed | Google Scholar

6.Hamma K, Iouknane D. Acute intestinal occlusions in l'adulte: étude rétrospective à propos de 75 cas de Janvier 2012 à

December 2013 at the department of general surgery. University of béjaia. 2014. Google Scholar

7. Ajmi S, Sahnoun L, Boujelbene N, Sriha B, Essabbah H. Hemorrhage from a child's Meckel diverticulum discovered by Technétium 99m scintigraphy. *J Chir (Paris)*. July 1 2005;142(4):244-6. PubMed | Google Scholar

8. Malhotra S, Roth AD, Gouge TH, et al. Gangrene of Meckel's diverticulum secondary to axial torsion: a rare complication. *Am J Gastroenterol*. 1998 Aug;93(8):1373-5. PubMed | Google Scholar

9. Barbary C, Floquet M, Riebel N, Tissier S, Laurent V, Régent D. Enterolith occlusion and Meckel's diverticulitis associated with a hernia in an adult. *J Radiol*. 1 Jan. 2004;85(1):49-50. PubMed | Google Scholar

10. Franchi S, Martelli H, Paye-Jaouen A, Goldszmidt D, Pariente D. Acute intestinal invagination in infants and children. *EMC- Pédiatrie*. Feb 1, 2005;2(1):45-57. Google Scholar

11. Khemekhem R, Ben Ahmed Y, Rahay H, Soufiane G, Said J, Douira W et al. Pathological aspects of Meckel's diverticulum in children. *J Pédiatrie Puériculture*. June 1, 2013;26(3):146-50. Google Scholar

12.Sagar J, Kumar V, Shah DK. Meckel's diverticulum: a systematic review. J R Soc Med. Oct 2006;99(10):501-5.

13.Cissé M, Konaté I, Dieng M, Fall MB, Ka O, Ngom G, Touré CT.

Meckel diverticula complicated by intestinal occlusions:

prise en charge à Dakar, à propos de 10 cas. J Afr Chir Digest.

2008;8(2):782-787

14.Grabin C, Bonnard A, Helardot P-G. Meckel's diverticulum surgery.

diverticulum. EMC-Chirurgie. 2005;(2):613-620. Google Scholar

UNDER PEER REVIEW