

SEQUENTIALLY GENERATED MEASURABLY BOUND PRODUCTS

Abstract

Many studies have been done on products of measurable sets. The most recent results highlight the properties of tensor products expressed as matrix products. In this study, we investigate the conditions under which sequentially generated products of functions are measurably bound using $(\epsilon - \delta)$ criterion for uniform continuity. The study sheds light on the application of r -neighbourhood topological properties of refinement of measurable sets in determining the boundedness of sequentially generated products of measurable functions. Concepts such as monotonicity of functions, continuity from above of set functions, almost everywhere properties and r -neighbourhood partition of measurable sets are applied in the context of p -integrable functions. The results of this research can be applied to develop the r -neighbourhood business models where r represents the physical distance around a fixed business focal point that geographically creates a fruitful business environment for achievement of the optimal industrial and commercial profit margins determined by the boundedness of product functions. For a fixed product of functions i.e. the target of achievement, one can sequentially and by monotonicity of measurable functions determine the quantitative (or measurable) convergence of the product of functions which represents the interactive operational activities towards the defined business goals. Further, the results of this study can be applied in developing geometrical models

in engineering by quantitative approximation to desired standards.

Keywords : Refinement, Measurably bound, Monotonically decreasing, r -neighbourhood .

1 Introduction

This study makes the sequence $(f_i \otimes \chi_A)_{i=1}^{\infty}$ of products of f_i and χ_A where $f_i \otimes \chi_A \in L_p(\mu) \otimes \Sigma$ move quantitatively closer to $f \otimes x'$ for each i . With an appropriate choice of a real number $r > 0$, the r -neighbourhood $N_r(x_i)$ of a point $x_i \in X$ as discussed in [2, 14] partitions an open set G_i for $i \in I$ [1, 7] such that the set

$$(L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^n : \| (\langle \mu_{|f_j|^p}(N_r(x_i)), x' \rangle)^{1/p} - (\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p} \| \| x' \|^{1/p'} \geq \epsilon$$

is made progressively smaller for large values of j .

Concepts on uniform continuity of functions (see[5]) and geometrical estimation of distance from a point to a given set (see[14]) are utilized so that the quantity

$(\langle \mu_{|f_j|^p}(N_r(x_i)), x' \rangle)^{1/p} \| x' \|^{1/p'}$ is kept within the ϵ -distance of the quantity $(\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p} \| x' \|^{1/p'}$ as we restrict a point $y \in G_i$ for each $i \in I$ to smaller intervals $N_r(x_i)$ of $x_i \in X$

While the continuity of functions is discussed, we study the behaviour of sequentially generated tensor products within the ϵ -distance parameter where the $(\epsilon - \delta)$ criterion for uniform continuity is applied.

2 Preliminaries

In this study, we consider $1 \leq p < \infty$ and the conjugate real number p' such that $1/p + 1/p' = 1$ (see[1, 3, 10, 12]). Throughout this paper, (Ω, Σ, μ) denotes a measure space where Σ is a sigma ring of subsets of Ω , $\mu : \Sigma \rightarrow X$ is a countably

additive vector measure, X a Banach space, $L_p(\mu)$ the space of p -integrable functions with respect to μ . The function $\langle x, x' \rangle$ denotes the duality between the Banach space X with its topological dual X' . For each $x' \in X'$, we have $\langle \mu(A), X' \rangle = \langle x, x' \rangle$ for every $A \in \Sigma$ (see[1]). If a sequence $(f_n)_{i=1}^\infty \in L_p(\mu)$ and χ_A is the characteristic function of a measurable set A of finite measure, then $f_n \otimes \chi_A$ denotes the product of f_n and χ_A such that $f_n \otimes \chi_A \in L_p(\mu) \otimes \Sigma$ for each n

Let $\langle \mu, x' \rangle = \mu_{x'}$ for every $x' \in X'$ such that $\mu_{x'} = \mu_{x' \div \|x'\|}(\|x'\|)$

Therefore,

$$\begin{aligned} (\int |f_j|^p \cdot \chi_A d\langle \mu, x' \rangle)^{1/p} \|x'\|^{1/p'} &= (\langle \mu_{|f_j|^p}(A), x' \div \|x'\| \rangle)^{1/p} (\|x'\|) \\ &= (\langle \mu_{|f_j|^p}(A), x' \rangle)^{1/p} (\|x'\|) \|x'\|^{-1/p} \\ &= (\langle \mu_{|f_j|^p}(A), x' \rangle)^{1/p} \|x'\|^{1/p'} \end{aligned}$$

Therefore, $(\langle \mu_{|f_j|^p}(A), x' \rangle)^{1/p} \|x'\|^{1/p'}$ is well defined as demonstrated in [1, 3, 7, 8, 9, 10, 11], where $\mu_{|f_j|^p}(A) \in X$ for every $A \in \Sigma$.

Definition 1(Refinement)

A family $(A_j : j \in \alpha)$ of subsets of X is called a refinement of a set G if for every r_i -neighbourhood $N_{r_i}(x_i)$ of a point x_i in A_i , there is a subset G_i in G such that

$$\langle \mu_{|f_j|^p}(N_{r_i}(x_i)), x' \rangle^{1/p} \leq \langle \mu_{|f_j|^p}(G_i), x' \rangle^{1/p} \text{ for each } i \text{ and } j$$

Therefore,

$$\langle \mu_{|f_j|^p}(\bigcup_{i=1}^\infty N_{r_i}(x_i)), x' \rangle^{1/p} = \langle \mu_{|f_j|^p}(G_i), x' \rangle^{1/p}$$

$$\sum_{i=1}^\infty \langle \mu_{|f_j|^p}(N_{r_i}(x_i)), x' \rangle^{1/p} = \langle \mu_{|f_j|^p}(G_i), x' \rangle^{1/p}$$

for each j

Definition 2(Measurably bound Products)

Let (Ω, Σ, μ) be a measure space and $E_{n_j+1} \subset E_{n_j}$ for each n_j be set of measurable sets each of which is a refinement of G such that $\mu(E_{n_1}) < \infty$ for all n_j . There

exists a neighbourhood $N_r(x_i)$ such that $N_r(x_i) \subset G_i \in G$.

$$\text{Define } E_{n_j} = (L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^n : \| [(\langle \mu_{|f_j|^p}(N_r(x_i)), x' \rangle)^{1/p} - (\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p}] \| x' \|^{1/p'} \geq \epsilon$$

Such that

$$(\langle \mu_{|f_j|^p}(E_n), x' \rangle)^{1/p} = (\langle \mu_{|f_j|^p}(\bigcap_{j=1}^{\infty} E_{n_j}), x' \rangle)^{1/p} \text{ where } E_n \downarrow \emptyset \text{ for each } n.$$

Therefore,

$$\begin{aligned} (\langle \mu_{|f_j|^p}(E_n), x' \rangle)^{1/p} &= (\langle \mu_{|f_j|^p}(\bigcap_{j=1}^{\infty} E_{n_j}), x' \rangle)^{1/p} \\ &\leq (\langle \mu_{|f_j|^p}(E_{n_j}), x' \rangle)^{1/p} \leq (\langle \mu_{|f_j|^p}(E_{n_1}), x' \rangle)^{1/p} \end{aligned}$$

for all n_j

As noted in [6, 7] regarding monotonically increasing sets, it follows that

$$(\langle \mu_{|f_j|^p}((E_{n_j})^c), x' \rangle)^{1/p} \uparrow (\langle \mu_{|f_j|^p}((E_n)^c), x' \rangle)^{1/p} \text{ for each } n$$

where $(E_{n_j})^c$ and $(E_n)^c$ are the complements of E_{n_j} and E_n respectively (for examples on complements of sets, see[2, 13]).

The results in [4, 8] on monotonically decreasing sets show that ,

$$(\langle \mu_{|f_j|^p}((E_{n_j})), x' \rangle)^{1/p} \downarrow (\langle \mu_{|f_j|^p}((E_n)), x' \rangle)^{1/p} \text{ for each } n_j$$

From the results in [4] on the convergence of measurable sets to zero with respect to c^* -algebra valued measures, in (see[1]) on integral mappings and order continuous Banach spaces of integrable functions with respect to vector measure and in [8, 12] on projective tensor products, the sequence $(f_i \otimes \chi_A)_{i=1}^{\infty}$ of products of measurable functions, where $f_i \otimes \chi_A \in L_p(\mu) \otimes \Sigma$ is therefore said to be measurably bound to $f \otimes x'$ at a point x if

$$\begin{aligned} (L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^n : \| [(\langle \mu_{|f_j|^p}(N_r(x_i)), x' \rangle)^{1/p} - (\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p}] \| x' \|^{1/p'} \geq \epsilon \end{aligned}$$

is monotonically decreasing to a null set as $n \rightarrow \infty$ for $\epsilon > 0$

Definition 3(r-Neighbourhood Partition)

Let $(G_i : i \in I)$ be a family of measurable open subsets of the normed space X and $(x_i)_{i=1}^{\infty}$ be a sequence of elements in G_i for each $i \in I$ where I is an index set. The sequence $(N_{r_1}(x_1), N_{r_2}(x_2), \dots, N_{r_{i-1}}(x_{i-1}), N_{r_i}(x_i), N_{r_{i+1}}(x_{i+1}) \dots)$ of r_i - neighbourhoods of x_i such that $N_{r_i}(x_i) \cap N_{r_j}(x_j) = \emptyset$ for $i \neq j$ is said to partition G_i into disjoint sets for each $i \in I$ if

$$N_{r_i}(x_i) \uparrow G_i \text{ for each } i \in I \text{ (see[6, 7]).}$$

Therefore

$$\bigcup_{i=1}^{\infty} N_{r_i}(x_i) = G_i \text{ for each } i \in I.$$

If $r = \min (r_1, r_2, \dots, r_{i-1}, r_i, r_{i+1}, \dots)$, then

$$N_r(x_i) \subset G_i \text{ for each } i \in I \text{ (see[2, 14])}$$

Definition 4(Directed Set of Vector Measure Duality)(see [8, 9])

A set $(\langle \mu_i, x' \rangle)_{i=1}^n$ of non-negative vector measure duality is said to be increasingly directed if for $\langle \mu_i, x' \rangle \leq \langle \mu_{k_i}, x' \rangle$ where $1 \leq i < k_i \leq n$ we have

$$\langle \mu_i(A), x' \rangle = LUB_k \langle \mu_{k_i}(A), x' \rangle \text{ for every } A \in \Sigma$$

Definition 4(Almost Everywhere Property)(see [1, 3, 10])

Let x be an element in X . A proposition $P(x)$ is true almost everywhere if there exists a null set E such that $x \in X \sim E$

3 Main Results

Proposition 1

Let $(f_n)_{i=1}^{\infty}$ and f be p -integrable functions such that f_n converges to f

uniformly. The set $(f_i \otimes \chi_A)_{i=1}^n$ of measurable functions where $f_i \otimes \chi_A \in L_p(\mu) \otimes \Sigma$ for each i is said to be measurably bound to $f \otimes x'$ at a point x if

$$(L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=m}^n : \| [(\sum_{j=m}^n < \mu_{|f_j|^p}(N_r(x_i)), x' >)^{1/p} - (< \mu_{|f|^p}(G_i), x' >)^{1/p}] \| x' \|^{1/p'} \| \geq \epsilon/2$$

is monotonically decreasing to a null set as $n \rightarrow \infty$ for $\epsilon > 0$

Proof

Since f_n converges to f uniformly and $f_i \otimes \chi_A \in L_p(\mu) \otimes \Sigma$ for $i = 1, \dots, n$ is measurably bound to $f \otimes x'$ at a point x (by hypothesis), we need to sequentially show that set $(f_1 \otimes \chi_A, \dots, f_n \otimes \chi_A)$ and $f \otimes x'$ which are at a distance greater than an arbitrary real number $\epsilon > 0$, can be made progressively smaller for values that are in some r -neighbourhood $N_r(x_i)$ of $x_i \in X$ (see [2, 14]). Further, the $(\epsilon - \delta)$ criterion on uniform continuity of functions (see [5]) is applied to obtain the desired results. Therefore, given $\epsilon > 0$ there exists a $\delta > 0$ such that

$$\nabla f_n = (L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^n : \| [(\sum_{j=1}^n < \mu_{|f_j|^p}(N_r(x_i)), x' >)^{1/p} - (< \mu_{|f|^p}(G_i), x' >)^{1/p}] \| x' \|^{1/p'} \| \geq \epsilon$$

is monotonically decreasing to a null set as $n \rightarrow \infty$

provided $y \in N_\delta(x_i) \forall y \in G_i$.

Suppose we choose $n > m$ such that

$$\nabla f_{m-1} = (L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^{m-1} : \| [(\sum_{j=1}^{m-1} < \mu_{|f_j|^p}(N_r(x_i)), x' >)^{1/p} - (< \mu_{|f|^p}(G_i), x' >)^{1/p}] \| x' \|^{1/p'} \| \geq \epsilon/2$$

The choice of $\epsilon > 0$ and $\epsilon/2 > 0$ Implies that

$$\nabla f_{m-1} \subseteq \nabla f_n$$

Therefore,

$$(L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^n : \| [(\sum_{j=1}^n < \mu_{|f_j|^p}(N_r(x_i)), x' >)^{1/p} - (\sum_{j=1}^m < \mu_{|f_j|^p}(N_r(x_i)), x' >)^{1/p}] \| x' \|^{1/p'} \geq \epsilon/2)$$

Fix n , taking limits as $m \rightarrow \infty$ and applying Cauchy criterion as discussed in [1, 5], we obtain

$$(L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=m}^n : \| [(\sum_{j=m}^n < \mu_{|f_j|^p}(N_r(x_i)), x' >)^{1/p} - (< \mu_{|f|^p}(G_i), x' >)^{1/p}] \| x' \|^{1/p'} \geq \epsilon/2) = \nabla^{f_{m-1}} \cap \nabla^{f_n}$$

whenever $y \in N_\delta(x_i) \forall y \in G_i$

Therefore,

$$(L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=m}^n : \| (\sum_{j=m}^n < \mu_{|f_j|^p}(N_r(x_i)), x' >)^{1/p} \| x' \|^{1/p'} - (< \mu_{|f|^p}(G_i), x' >)^{1/p} \| x' \|^{1/p'} \geq \epsilon/2) \subseteq \nabla^{f_n} \downarrow \emptyset$$

as $n \rightarrow \infty$

Hence,

$$(L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=m}^n : \| [(\sum_{j=m}^n < \mu_{|f_j|^p}(N_r(x_i)), x' >)^{1/p} \| x' \|^{1/p'} - (< \mu_{|f|^p}(G_i), x' >)^{1/p} \| x' \|^{1/p'} \geq \epsilon/2) \downarrow \emptyset$$

as $n \rightarrow \infty$ provided $y \in N_\delta(x_i) \forall y \in G_i$

Proposition 2

Let $(f_n)_{i=1}^\infty$, f and g be p -integrable functions such that the set $(f_1 \otimes \chi_A \dots f_n \otimes \chi_A)$ of products where $f_i \otimes \chi_A \in L_p(\mu) \otimes \Sigma$ for each i is measurably bound to $f \otimes x'$ and $g \otimes x'$ at a point x . If for $\epsilon > 0$, we have

$$\| (< \mu_{|f|^p}(G_i), x' >)^{1/p} - (< \mu_{|g|^p}(G_i), x' >)^{1/p} \| < \epsilon$$

where $(G_i : i \in I)$ is a family of measurable open subsets of the normed space X ,

then

$$(\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p} = (\langle \mu_{|g|^p}(G_i), x' \rangle)^{1/p} \text{ a.e.}$$

Proof

$$\text{Let } (\nabla)_f^g = ((L_p(\mu)_{f \otimes x'}^{g \otimes x'}) : \| [(\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p} \\ - (\langle \mu_{|g|^p}(G_i), x' \rangle)^{1/p}] \| \| x' \|^{1/p'} \| < \epsilon)$$

for each $i \in I$

Let the collection $(A_j : j \in \alpha)$ be a refinement of a set G . For every r_i -neighbourhood $N_{r_i}(x_i)$ of a point x_i in A_j , there exists a subset G_i in G such that $N_{r_i}(x_i) \subseteq G_i$ for each i and j

Take $r = 1/2 \min (r_1, r_2, \dots, r_n)$. It follows that $r > 0$ and

$$N_r(x_i) \subseteq N_{r_i}(x_i) \subseteq G_i \text{ for each } i \in I$$

On application of $(\epsilon - \delta)$ criterion on uniform continuity as discussed in [5] and the duality function $\langle \mu, x' \rangle$ (see[1, 3, 9, 10, 11]), we take y closer to x_i such that for $\delta > 0$, we have

$$y \in N_\delta(x_i) \forall y \in G_i$$

Consequently,

$$(L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^n : \| [(\sum_{j=1}^n \langle \mu_{|f_j|^p}(N_r(x_i)), x' \rangle)^{1/p} \\ - (\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p}] \| \| x' \|^{1/p'} \| \geq \epsilon/2)$$

$$\text{Let } \nabla_{f_n}^f = (L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^n : \| [(\sum_{j=1}^n \langle \mu_{|f_j|^p}(N_r(x_i)), x' \rangle)^{1/p} \\ - (\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p}] \| \| x' \|^{1/p'} \| \geq \epsilon/2)$$

and

$$\nabla_{f_n}^g = (L_p(\mu)_{f_j \otimes \chi_A}^{g \otimes x'})_{j=1}^n : \| [(\sum_{j=1}^n \langle \mu_{|f_j|^p}(N_r(x_i)), x' \rangle)^{1/p}$$

$$- (\langle \mu_{|g|^p}(G_i), x' \rangle)^{1/p} \|x'\|^{1/p'} \geq \epsilon/2)$$

Since $\|[(\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p}$

$$- (\langle \mu_{|g|^p}(G_i), x' \rangle)^{1/p} \|x'\|^{1/p'} \| < \epsilon \text{ on } (\nabla)_f^g$$

for each $i \in I$, it follows that

$$\begin{aligned} ((\nabla)_f^g)^c &= ((L_p(\mu)_{f \otimes x'}^{g \otimes x'}) : \|[(\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p} \\ &- (\langle \mu_{|g|^p}(G_i), x' \rangle)^{1/p} \|x'\|^{1/p'} \geq \epsilon) \end{aligned}$$

Therefore,

$$((\nabla)_f^g)^c \subseteq \nabla_{f_n}^f \cup \nabla_{g_n}^f$$

Since the set

$(f_1 \otimes \chi_A \dots f_n \otimes \chi_A)$ is measurably bound to $f \otimes x'$ and $g \otimes x'$ at a point x (by hypothesis), it follows that

$\nabla_{f_n}^f$ and $\nabla_{g_n}^f$ are both monotonically $\downarrow \emptyset$ as $n \rightarrow \infty$

Subsequently,

The $\langle \mu, x' \rangle$ - measure of $((\nabla)_f^g)^c$ is zero,

provided $y \in N_\delta(x_i) \forall y \in G_i$

Suppose,

$$\begin{aligned} ((L_p(\mu)_{f \otimes x'}^{g \otimes x'}) : \|[(\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p} \\ - (\langle \mu_{|g|^p}(G_i), x' \rangle)^{1/p} \|x'\|^{1/p'} \neq 0) \\ = \bigcup_{k=1}^{\infty} ((L_p(\mu)_{f \otimes x'}^{g \otimes x'}) : \|(\langle \mu_{|f|^p}(G_i), x' \rangle \\ - (\langle \mu_{|g|^p}(G_i), x' \rangle)^{1/p} \|x'\|^{1/p'} \geq 1/k) = 0 \end{aligned}$$

It follows that

$$\begin{aligned} ((L_p(\mu)_{f \otimes x'}^{g \otimes x'}) : \| [(\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p} \\ - (\langle \mu_{|g|^p}(G_i), x' \rangle)^{1/p}] \| \| x' \|^{1/p'} \| \neq 0) \end{aligned}$$

is a null set since it is equal to the countable union of null sets (see [1, 7]).

The results on almost everywhere property in [1, 3, 10] demonstrate that

$$\langle \mu_{|f|^p}(G_i), x' \rangle^{1/p} = \langle \mu_{|g|^p}(G_i), x' \rangle^{1/p} \text{ a.e.}$$

Proposition 3

Let $(f_n)_{i=1}^\infty$, f and g be p -integrable functions such that the set $(f_1 \otimes \chi_A, \dots, f_n \otimes \chi_A)$ of products where $f_i \otimes \chi_A \in L_p(\mu) \otimes \Sigma$ for $i = 1, \dots, n$ is measurably bound to $f \otimes x'$ at a point x . If

$$((L_p(\mu)_{f \otimes x'}^{g \otimes x'}) : \langle \mu_{|f|^p}(G_i), x' \rangle^{1/p} \neq \langle \mu_{|g|^p}(G_i), x' \rangle^{1/p})$$

where G_i for each $i \in I$ is null set, then $(f_1 \otimes \chi_A, \dots, f_n \otimes \chi_A)$ is measurably bound to $g \otimes x'$ at a point x

Proof

Let E denote the set

$$((L_p(\mu)_{f \otimes x'}^{g \otimes x'}) : \langle \mu_{|f|^p}(G_i), x' \rangle^{1/p} \neq \langle \mu_{|g|^p}(G_i), x' \rangle^{1/p})$$

Then E is a null set (by hypothesis). From the results on almost everywhere

property in [1, 3, 10], it follows that

$$\langle \mu_{|f|^p}(G_i), x' \rangle^{1/p} = \langle \mu_{|g|^p}(G_i), x' \rangle^{1/p} \text{ a.e.}$$

$$\begin{aligned} \text{Let } F_n = ((L_p(\mu)_{f_j \otimes \chi_A}^{g \otimes x'})_{j=1}^n : \| (\sum_{j=1}^n [\langle \mu_{|f_j|^p}(N_r(x_i)), x' \rangle]^{1/p} \\ - \langle \mu_{|g|^p}(G_i), x' \rangle)^{1/p} \| \| x' \|^{1/p'} \| \geq \epsilon) \end{aligned}$$

$$\begin{aligned} &\subseteq E \cup (L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^n : \| (\sum_{j=1}^n < \mu_{|f_j|^p}(N_r(x_i)), x' >)^{1/p} \| x' \|^{1/p'} \\ &- (< \mu_{|f|^p}(G_i), x' >)^{1/p} \| x' \|^{1/p'} \geq \epsilon) \end{aligned}$$

Since E is null set and $(f_1 \otimes \chi, \dots, f_n \otimes \chi_A)$ is measurably bound to $f \otimes x'$ at a point x by hypothesis), then

$$\begin{aligned} &((L_p(\mu)_{f_j \otimes \chi_A}^{g \otimes x'})_{j=1}^n : \| ((\sum_{j=1}^n [(< \mu_{|f_j|^p}(N_r(x_i)), x' >)^{1/p} \\ &- (< \mu_{|g|^p}(G_i), x' >)^{1/p}] \| x' \|^{1/p'} \geq \epsilon) \end{aligned}$$

From which the results follows (see Proposition 1) that

$$F_n \downarrow \emptyset \text{ as } n \rightarrow \infty$$

Proposition 4

Let $(f_n)_{i=1}^\infty, (g_n)_{i=1}^\infty$ and f be p -integrable functions such that the set $(f_1 \otimes \chi_A, \dots, f_n \otimes \chi_A)$ is measurably bound to $f \otimes x'$ at a point x . If

$$\begin{aligned} &((L_p(\mu)_{f_j \otimes \chi_A}^{g_i \otimes \chi_A}) : \| [(< \mu_{|f_j|^p}(N_r(x_i)), x' >)^{1/p} \\ &- (< \mu_{|g_i|^p}(N_r(x_i)), x' >)^{1/p}] \| x' \|^{1/p'} = 0) \end{aligned}$$

is a non empty set for each $j = 1, \dots, n$, then given a real $\epsilon > 0$, the set

$$\begin{aligned} &((L_p(\mu)_{g_j \otimes \chi_A}^{f \otimes x'}) : \| [(\sum_{j=1}^n < \mu_{|g_j|^p}(N_r(x_i)), x' >)^{1/p} \\ &- (< \mu_{|f|^p}(G_i), x' >)^{1/p}] \| x' \|^{1/p'} \geq \epsilon) \end{aligned}$$

Proof

$$\begin{aligned} \text{Let } E_n &= ((L_p(\mu)_{f_j \otimes \chi_A}^{g_i \otimes \chi_A}) : \| [(< \mu_{|f_j|^p}(N_r(x_i)), x' >)^{1/p} \\ &- (< \mu_{|g_i|^p}(N_r(x_i)), x' >)^{1/p}] \| x' \|^{1/p'} = 0) \end{aligned}$$

Since $E_n \neq \emptyset$ (by hypothesis), then by almost everywhere property for pairwise distinct sets as discussed in [1, 3, 10] , it follows that

$$(E_n)^c = ((L_p(\mu)_{f_j^{\otimes \chi_A}})^{g_i} : (\langle \mu_{|f_j|^p}(N_r(x_i)), x' \rangle)^{1/p} \\ \neq (\langle \mu_{|g_i|^p}(N_r(x_i)), x' \rangle)^{1/p})$$

is a null set.

Suppose

$$((L_p(\mu)_{g_j^{\otimes \chi_A}})^{f \otimes x'})_{j=1}^n : \| [(\sum_{j=1}^n \langle \mu_{|g_j|^p}(N_r(x_i)), x' \rangle)^{1/p} \\ - (\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p}] \| \| x' \|^{1/p'} \geq \epsilon \\ \subseteq (E_n)^c \cup ((L_p(\mu)_{f_j^{\otimes \chi_A}})^{f \otimes x'})_{j=1}^n : \| (\sum_{j=1}^n \langle \mu_{|f_j|^p}(N_r(x_i)), x' \rangle)^{1/p} \| \| x' \|^{1/p'} \\ - (\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p} \| \| x' \|^{1/p'} \geq \epsilon)$$

Since $(E_n)^c$ is a null set and $(f_1 \otimes \chi_A, \dots, f_n \otimes \chi_A)$ is measurably bound to $f \otimes x'$ at x it follows from Proposition 3 that

$$((L_p(\mu)_{g_j^{\otimes \chi_A}})^{f \otimes x'})_{j=1}^n : \| [(\sum_{j=1}^n \langle \mu_{|g_j|^p}(N_r(x_i)), x' \rangle)^{1/p} \\ - (\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p}] \| \| x' \|^{1/p'} \geq \epsilon)$$

is monotonically $\downarrow \emptyset$ as $n \rightarrow \infty$ as required

Proposition 5

Let the family $(A_j : j \in \alpha)$ of subsets of X be a refinement of a set G_i for each i and $(f_n)_{i=1}^\infty, f$ be p -integrable functions such that the set $(f_1 \otimes \chi_A, \dots, f_n \otimes \chi_A)$ is measurably bound to $f \otimes x'$ at $x \in X$. If for given real numbers $\beta > 0$ and $\epsilon > 0$, the set

$$((L_p(\mu)_{f_k^{\otimes \chi_A}})^{f \otimes x'})_{k \geq n} : \| [(\sum_{k \geq n} \langle \mu_{|f_k|^p}(N_r(x_i)), x' \rangle)^{1/p} \\ - (\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p}] \| \| x' \|^{1/p'} \geq 1/\beta)$$

then for $\epsilon(\beta) > 0$,

$$\| [(\sum_{k \geq n} \langle \mu_{|f_k|^p}(N_r(x_i)), x' \rangle)^{1/p}$$

$$- (\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p} \|x'\|^{1/p'} < \epsilon$$

Proof

Since $(A_j : j \in \alpha)$ is a refinement of a set G_i for each i (by hypothesis), then for every r -neighbourhood $N_r(x_i)$ of a point x_i in A_i , there is a subset G_i in G such that $N_r(x_i) \subseteq G_i$ for each i and j

For each natural number i , each $N_r(x_i)$ has non empty lower bound B in G_i such that

$$(\langle \mu_{|f_j|^p}(B), x' \rangle)^{1/p} \|x'\|^{1/p'} < \epsilon$$

$$(\langle \mu_{|f_j|^p}(N_r(x_i)), x' \rangle)^{1/p} \|x'\|^{1/p'} < \epsilon$$

Therefore,

$$\lim_{j \rightarrow \infty} (\langle \mu_{|f_j|^p}(N_r(x_i)), x' \rangle)^{1/p} \|x'\|^{1/p'} < \epsilon$$

Define $E_n^\beta = \bigcup_k ((L_p(\mu)_{f_k \otimes \chi_A}^{f \otimes x'})_{k \geq n} : \|[(\sum_{k \geq n} \langle \mu_{|f_k|^p}(N_r(x_i)), x' \rangle)^{1/p}$

$$- (\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p} \|x'\|^{1/p'} \| \geq 1/\beta)$$

Since $(f_1 \otimes \chi_A, \dots, f_n \otimes \chi_A)$ is measurably bound to $f \otimes x'$ at $x \in X$, as noted in proposition 1, there exists a real number $\beta > 0$ satisfying

$$((L_p(\mu)_{f_k \otimes \chi_A}^{f \otimes x'})_{k \geq n} : \|[(\sum_{k \geq n} \langle \mu_{|f_k|^p}(N_r(x_i)), x' \rangle)^{1/p}$$

$$- (\langle \mu_{|f|^p}(G_i), x' \rangle)^{1/p} \|x'\|^{1/p'} \| \geq 1/\beta) \downarrow \emptyset \text{ as}$$

$$n \rightarrow \infty$$

Subsequently,

$$(\langle \mu_{|f|^p}(E_n^\beta), x' \rangle)^{1/p} \downarrow 0$$

Define $E = \bigcup_{\beta=1}^\infty E_n^\beta$

$E^c = \bigcap_{\beta=1}^\infty (E_n^\beta)^c$ where E^c is the complement of E in $L_p(\mu) \otimes X'$

Therefore,

$$\begin{aligned} E^c &= \bigcap_{\beta=1}^{\infty} \bigcap_k ((L_p(\mu)_{f_k \otimes \chi_A}^{f \otimes x'})_{k \geq n} : \| [(\sum_{k \geq n} < \mu_{|f_k|^p}(N_r(x_i)), x' >)^{1/p} \\ &\quad - (< \mu_{|f|^p}(G_i), x' >)^{1/p}] \| x' \|^{1/p'} \geq 1/\beta)^c \\ &= \bigcap_{\beta=1}^{\infty} \bigcap_k ((L_p(\mu)_{f_k \otimes \chi_A}^{f \otimes x'})_{k \geq n} : \| [(\sum_{k \geq n} < \mu_{|f_k|^p}(N_r(x_i)), x' >)^{1/p} \\ &\quad - (< \mu_{|f|^p}(G_i), x' >)^{1/p}] \| x' \|^{1/p'} < 1/\beta) \end{aligned}$$

Let $1/\beta < \epsilon$ for $\beta > 0$

Therefore,

$$\begin{aligned} \bigcap_{\beta=1}^{\infty} \bigcap_k ((L_p(\mu)_{f_k \otimes \chi_A}^{f \otimes x'})_{k \geq n} : \| [(\sum_{k \geq n} < \mu_{|f_k|^p}(N_r(x_i)), x' >)^{1/p} \\ - (< \mu_{|f|^p}(G_i), x' >)^{1/p}] \| x' \|^{1/p'} < \epsilon) \end{aligned}$$

On E^c .

Corollary 1

Let $(f_n)_{i=1}^{\infty}, (g_n)_{i=1}^{\infty}$ and f be p -integrable functions with respect to an increasingly directed scalar measure $< \mu_i, x' >$ for each i and $(G_i : i \in I)$ be a family of closed measurable subsets of X . If $(f_1 \otimes \chi_A, \dots, f_n \otimes \chi_A)$ is measurably bound to $f \otimes x'$ over the r -neighbourhood $N_r(x_i)$ of a point x_i in $A \in \Sigma$, then

$$\begin{aligned} ((L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^n : \| [(\sum_{j=1}^n < \mu_{k_j} |f_j|^p(N_r(x_i)), x' >)^{1/p} \\ - (< \mu_{k_j} |f|^p(X \sim G_i), x' >)^{1/p}] \| x' \|^{1/p'} \geq \epsilon) \downarrow \emptyset \end{aligned}$$

Proof

Since G_i is closed for each i , then $G^C = X \sim G_i$ is topologically open follows from the results in [2, 13, 14].

Therefore,

$$A \cap G_i^c = A \sim G_i \neq \emptyset$$

If $r = \min (r_1, r_2, \dots, r_n)$, the results in [2, 13, 14] demonstrate that for a topologically open set G_i , there exists an r -neighbourhood $N_r(x_i)$ such that

$$N_r(x_i) \subset G_i \text{ for } r > 0 \text{ and for each } i$$

By definition of measurably bound products over $N_r(x_i)$, it suffices to show that

$$\begin{aligned} & ((L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^n : \| [(\sum_{j=1}^n < \mu_j | f_j |^p(N_r(x_i)), x' >)]^{1/p} \\ & \quad - (< \mu_j | f |^p(G_i^c), x' >)^{1/p} \| \| x' \|^{1/p'} \geq \epsilon) \downarrow \emptyset \end{aligned}$$

Applying the results discussed in [8, 9] in the context of increasingly directed set of vector measure duality, it follows that

$$\begin{aligned} & ((L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^n : \| [(\sum_{j=1}^n < \mu_j | f_j |^p(N_r(x_i)), x' >)]^{1/p} \\ & \quad - (< \mu_j | f |^p(G_i^c), x' >)^{1/p} \| \| x' \|^{1/p'} \geq \epsilon) \\ & = LUB_k((L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^n : \| [(\sum_{j=1}^n < \mu_{k_j} | f_j |^p(N_r(x_i)), x' >)]^{1/p} \\ & \quad - (< \mu_{k_j} | f |^p(G_i^c), x' >)^{1/p} \| \| x' \|^{1/p'} \geq \epsilon) \\ & = LUB_k((L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^n : \| (\sum_{j=1}^n < \mu_{k_j} | f_j |^p(N_r(x_i)), x >)^{1/p} \\ & \quad - (< \mu_{k_j} | f |^p[(A \cup G_i)^c] \cup (X \sim G_i), x' >)^{1/p} \| \| x' \|^{1/p'} \geq \epsilon) \\ & \subseteq LUB_k((L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^n : \| [(\sum_{j=1}^n < \mu_{k_j} | f_j |^p(N_r(x_i)), x' >)]^{1/p} \\ & \quad - (< \mu_{k_j} | f |^p[(A \cup G_i)^c], x' >)^{1/p} \\ & \quad + (< \mu_{k_j} | f |^p[(X \sim G_i)], x' >)^{1/p} \| \| x' \|^{1/p'} \geq \epsilon) \end{aligned}$$

Therefore

$$\begin{aligned} & ((L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^n : \| [(\sum_{j=1}^n < \mu_{k_j} | f_j |^p(N_r(x_i)), x' >)]^{1/p} \\ & \quad - (< \mu_{k_j} | f |^p[(X \sim G_i)], x' >)^{1/p} \| \| x' \|^{1/p'} \geq \epsilon) \end{aligned}$$

$$\subseteq ((L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^n : \| (\sum_{j=1}^n < \mu_j | f_j |^p(N_r(x_i)), x' >)^{1/p} - (< \mu_j | f |^p(G_i^c), x' >)^{1/p} \| \| x' \|^{1/p'} \geq \epsilon)$$

The preceding discussion satisfy the monotone property in [6, 13] from which the results follows

$$((L_p(\mu)_{f_j \otimes \chi_A}^{f \otimes x'})_{j=1}^n : \| [(\sum_{j=1}^n < \mu_{k_j} | f_j |^p(N_r(x_i)), x' >)^{1/p} - (< \mu_{k_j} | f |^p(X \sim G_i), x' >)^{1/p} \| \| x' \|^{1/p'} \geq \epsilon) \downarrow \emptyset$$

3 Conclusion

The results obtained in this study highlight the boundedness of sequentially generated measurable products at a point using the neighbourhood topological properties of open sets.

REFERENCES

- [1] Agud, L., Calabuig, J. M., Juan, M. A. and Sanchez Perez, E. A. Banach Lattice Structures and Concavifications in Banach Spaces. *Mathematics*, 2020, 8(1), 127.
- [2] Almuhr, E., Miqdad, H., Al-labadi, M. and Idrisi, M. I. μ -L-Closed Subsets of Noetherian Generalized Topological Spaces. *International Journal of Neutrosophic Science*, 2024, 23(3), 148-48.
- [3] Calabuig, J. M., Rodrez, J., Shez-Pz, E. A. Strongly embedded subspaces of p-convex Banach function spaces. *Positivity*, 2013, 17(3), 775-791.
- [4] Gbemou, K. and Mensah, Y. Aspects of the Fourier-Stieltjes transform of C-algebra valued measures. *Journal of Advances in Mathematics and Computer Science*, 2019, 33(5), 1-8
- [5] Konstantogiannis, S. Relaxation of the Sequential Criteria for Continuity and

- Uniform Continuity of a Real Function. *The Teaching of Mathematics*, 2022, 25(2), 116-121
- [6] Otanga, O. L. On the finiteness and sigma finiteness of a cubic measure function. *Inter. Math. Forum, Hikari Ltd*, 2015, 10(3), 111 - 114.
- [7] Otanga, O. L., Oduor, M. O. and Aywa, S. O. Partition of measurable sets. *Journal of Advances in Mathematics*, 2015, 10 (8), 3759 - 3763.
- [8] Otanga, O. L., Oduor, M. O. and Aywa, S. O. On Generation of Measurable Covers for Measurable Sets Using Multiple Integral of Functions. *Journal of textit Mathematics and Statistical Science*, 2015, 2015, 32 -41
- [9] Otanga, O. L. and Oduor, M. O. On Pointwise product vector measure duality. *Journal of Advances in Mathematics*, 2021, 20, 8 - 18
- [10] Otanga, O. L.. On Absolute continuity of non negative functions. *Asian Research Journal of Mathematics*, 2022, 8(11), 385 - 392.
- [11] Olwamba, L. O. On Projection Properties of Monotone Integrable . Functions *Journal of Advances in Mathematics and Computer Science*, 2024, 39(3), 29-36.
- [12] Rodrez, J.and Zoca, A. R. Weak Precompactness in Projective Tensor Products. *Indagationes Mathematicae*, 2024, 35(1), 60-75.
- [13] Tyagi, B. K. and Chauhan, H. On Generalized Closed Sets in Generalized Topological Spaces. *Cubo (Temuco)*, 2016, 18(1), 27-45.
- [14] Van Ngai, H., Tron, N. H. and Tinh, P. N. Directional Hlder metric subregularity and application to tangent cones. *Journal of Convex Analysis*, 2017, 24(2), 417-457.