

# COMPARATIVE STUDY ON THE PREVALENCE OF GASTROINTESTINAL PARASITE BETWEEN WILD AND POND RAISED AFRICAN SHARPTOOTH CATFISH *Clarias gariepinus* IN AKWA IBOM STATE.

## ABSTRACT

A comparative study on the prevalence of gastrointestinal parasites of *Clarias gariepinus* in selected location in Oron and Uyo, Akwa Ibom state, Nigeria. A total of one hundred (100) samples were used, fifty (50) from the wild and fifty (50) from concrete pond. The wild samples were collected from fishermen in oron river, while pond samples were collected in Dominta farm in Uyo. Both location samples were examined for gastrointestinal parasite using standard scientific methods. The parasite obtained where nematode, cestode, and trematode. The research revealed that all the fish samples were infected with at least one of these parasites. Samples from concrete pond had 38%, 24% and 10% of nematode, cestode and trematode respectively while samples from the wild had a prevalence of 34%, 24, and 18% of nematode, cestode and trematode. Samples from the river and pond revealed that there was no significant difference ( $P>0.05$ ) in the prevalence of gastrointestinal parasite. Parasitic infection is a great hinderance to fish growth and therefore have adverse effects on fish productivity. Hence fish farmers should be sensitized about the presence and the effects of gastrointestinal parasite in the aquatic ecosystem. Communities should be enlightened about disposal of waste in the natural water bodies which increases the level of intermediate host that harbors this parasite.

**Keywords:** Comparison, Prevalence, Gastrointestinal parasites, Concrete ponds, Wild, *Clarias gariepinus* and Akwa Ibom State

## INTRODUCTION

Fish is a valuable food source that provides humans with income (Adelganet *et al.*, 2014) and high-quality protein and minerals (Soofi *et al.*, 2017). Fish can fulfil a person's nutritional needs due to their high digestible energy content (Rahei *et al.*, 2017). According to Ndupuhet *et al.* (2016), fish accounts for 40% of the average Nigerian's daily protein intake. The growing demand for animal protein has led to a notable growth in the development of fish farming cultures (Agbabiaka *et al.*, 2017). According to reports, the African catfish (*Clarias gariepinus*) is a suitable species for aquaculture in Nigeria and other African countries (Eyo *et al.*, 2015). These favourable characteristics include its ability to withstand a wide range of environmental conditions, its ability to grow quickly, its acceptance of artificial feed and its high fecundity rate. It is a resilient fish that can be raised in locations with scarce water supplies since it can be heavily stocked in low oxygen environments (Abuelhassan *et al.*, 2014). Due to its extremely high nutritional content, it is also well-liked by Nigerian customers (Udeze *et al.*, 2012). Parasites are an important group of pathogens, causing infection and diseases of fish both in fresh and marine environments (Emmy-Egbe, 2013). Economic losses have resulted from parasitic infection due to direct fish death, decreased fish development and fecundity and increased fish susceptibility to

diseases. Thus, fish health and the production of aquatic crops are increasingly threatened by parasitic infestations (Appah *et al.*, 2018). Because fish parasites coexist peacefully with their hosts as a means of survival, fish parasitic diseases are especially significant in tropical regions (Bichiet *al.*, 2018). However, parasitic disease increases quickly and causes significant mortality in situations where hosts are overcrowded, such as in fish farms or aquaria. In most cases, parasitic disease occurs in the wild as a result of anthropogenic activities and interference, such as pollution, affect the ecosystem and change the natural distribution of their parasite groups (Dabo *et al.*, 2018), hence making the environment conducive for parasite growth and development. Amidst many approaches to mitigate the fish parasite outbreak in Nigeria, data from archives suggest that a wide variety of parasitic infections may be present in both wild and homestead-cultured fish (Bichi and Ibrahim, 2009). Since fish cultivation is growing more intensive and popular, there is a need to monitor Nigerian freshwater bodies for parasitic illnesses as they pose a severe health risk to consumers. Furthermore, the intermediate host (crustaceans and aquatic insects) of these parasites are a contributing factor too (Pal and Ghosh, 1985). One reason for the rise in unskilled farmers is the rise in domestic fish farming. Improved agricultural practices for fish farmers must be taught to them (Ibrahim and Yahaya 2011). Ineffective and unprofessional methods of fish farming causes stress, disease and ultimately fish mortality. In ponds, compared to the wild, their exposure to parasites is restricted because of their confines. Fish farming has significant challenges as a result of the present economic unrest, including limited financial resources and an increase in new diseases that severely reduce fish market productivity and yield. Therefore, by identifying and contrasting the gastrointestinal parasites of fish bred in the wild and in concrete ponds, this study aims to close the knowledge gap. The fish studied are African Sharp-tooth Catfish (*Clarias gariepinus*) from Uyo, Akwa Ibom State, Nigeria.

## **MATERIALS AND METHODS**

### **The Study Area**

This study was carried out in Uyo (concrete pond) and Oron Local Government Areas (of Akwa Ibom State, Nigeria). Uyo is the capital of Akwa Ibom State which is located in the rainforest zone of the country and lies between latitude 4°32'N and 5°33'N and longitude 7°25'E and 8°25'E. Domita farm used for this study is located at no 4 Ring Road, Nsukara Offot, Uyo, Akwa Ibom State, Nigeria. Oron Local Government Area is one of the local government areas in Akwa Ibom State, it is a coastal area and the people of this area are majorly fishermen.

### **Ethics and consent**

The ethic and consent concerning the use of fish for this research were deemed unnecessary according to the University of Uyo. Meanwhile, the experimental protocol including the number of fishes used for this research was approved and regulated by the Institute. In addition, all the methods were carried out in accordance with relevant guidelines.

### **Collection of Fish Samples**

One Fifty (50) *Clarias gariepinus* samples were collected from water bodies in Oron (wild) and 50 from man-made reservoir pond (concrete pond). The concrete pond samples were bought at Domita farm (concrete), Uyo. The fish were harvested using a plastic basket. Harvested fish were put in a plastic bucket and transported alive immediately to the Animal and Environmental Biology laboratory in the Department of Animal and Environmental Biology, Faculty of Biological Science, University of Uyo. Uyo: and examined for the presence of gastrointestinal parasite within 24 hours of collection. Samples of fish from Oron River were booked and bought from local fishermen, they were collected fresh and alive from the fishermen as early as 6:00 am and transported to the laboratory alive in plastic bucket.

### Processing of Samples

At the laboratory, the fishes were rinsed with clean tap water, killed by suffocation and then parameters such as the standard lengths, total length and weights of the sample were measured using a meter rule and a weighting balance respectively. The sexes were determined using urogenital papillae and recorded. The samples were dissected to expose the alimentary canal which was then isolated and placed in petri dish containing physiological saline. The gastrointestinal tract was then slit longitudinally and examined for parasite under a light microscope (OLYMPUS.USA).

### Microscope Examination of Parasite

Gastrointestinal content was placed on a slide, mounted on the microscope stage and viewed under X10 and X40 objective for parasite observation. The parasite seen were identified using the pictorial guide manual of Eric (2010).

### Results

The results of this study reveal that, all samples were infected with at least one of either nematode, cestode or trematode. Samples from the wild recorded high prevalence of parasite than concrete pond (Table 1).

**Table 1:** Result of morphometric analysis of *Clarias gariepinus* from wild and concrete pond.

Group	Weight(g)	Total length(cm)	Standard length(cm)
Wild sample	263.9 ± 67.11	35.74 ± 5.0	30.79 ± 3.99
Concrete Sample	216.5 ± 52.43	34.66 ± 4.44	28.52 ± 3.90
Tcal	3.95	1.14	2.91
Df	49	49	49
Ttab	2.01	2.01	2.01
Decision:	No significant difference	No significant difference	No significant difference

**Table 2:** Sex base comparison of the prevalence of gastrointestinal parasite of *Clarias gariepinus* in concrete pond and wild.

Pond type	Sex	Number Examined	Number Infected with Nematode (%)	Number Infected with Cestode (%)	Number Infected with Trematode (%)	Total (%)
Samples from Concrete	Male	25	6(24.0)	7(28.0)	2(8.0)	15(60)
	Female	25	13(52.0)	5(20.0)	3(12.0)	21(85)
	Total	50	<b>19(38.0)</b>	<b>12(24.0)</b>	<b>5(10.0)</b>	<b>36(72)</b>
	Chi-Square		4.16	0.439	0.222	
	Df		1	1	1	
	p-value		0.041	0.508	0.637	
Samples from Wild	Male	26	7(26.9)	8(30.8)	7(26.9)	22(84.6)
	Female	24	10(41.7)	4(16.7)	2(8.3)	16(66.7)
	Total	50	<b>17(34.0)</b>	<b>12(24.0)</b>	<b>9(18.0)</b>	<b>38(76.0)</b>
	Chi-Square		1.209	1.361	2.922	
	Df		1	1		
	p-value		0.272ns	0.243ns	0.087ns	

**Table 3:** A weight-based comparison of the prevalence of gastrointestinal parasite of *Clarias gariepinus* between concrete pond and wild.

Pond type	Weight (g)	Number examined	Number infected with Nematode (%)	Number infected with Cestode (%)	Number infected by Trematode (%)
Concrete Pond	93.5-147.4	5	3(60.0)	1(20.0)	0(0.0)
	147.5-201.4	14	4(28.6)	1(7.1)	2(14.3)
	201.5-255.4	21	8(38.1)	9(42.9)	2(8.0)
	255.5-309.4	10	4(40.0)	1(10.0)	1(10.0)
	<b>Total</b>	<b>50</b>	<b>19(38.0)</b>	<b>12(24.0)</b>	<b>9(18.0)</b>
	Chi square		1.572	7.393	0.985
	Df		3	3	3
P		0.666ns	0.060ns	0.805ns	

	value				
<b>Sample from wild.</b>	158.5-257.4	27	9(33.3)	8(29.6)	5(18.5)
	257.5-356.4	19	6(31.6)	3(15.8)	4(21.1)
	356.5-455.4	3	1(33.3)	1(33.3)	0(0.0)
	455.5-554.4	1	1(100.0)	0(0.0)	0(0.0)
	<b>Total</b>	<b>50</b>	<b>17(34.0)</b>	<b>12(24.0)</b>	<b>9(18.0)</b>
	Chi square		1.997	1.63	1.003
	Df		3	3	3
P value		0.573ns	0.653ns	0.801ns	

**Table 4:** Total length comparison in the prevalence of gastrointestinal parasite in *Clarias gariepinus* between concrete pond and fishes from the wild.

Pond type	Total length	Number examined	Number infected with Nematode (%)	Number infected with Cestode (%)	Number infected with Trematode (%)
Concrete sample	25.5-30.4	12	4(33.3)	2(16.7)	1(8.3)
	30.5-35.4	16	6(37.5)	5(31.3)	2(12.5)
	35.5-40.4	17	7(41.2)	4(23.5)	1(5.9)
	40.5-45.4	5	2(40.0)	1(20.0)	1(20.0)
	<b>Total</b>	<b>50</b>	<b>19(38.0)</b>	<b>12(24.0)</b>	<b>5(10.0)</b>
	Chi square		0.194	0.861	1.024
	Df		3	3	3
P value		0.979ns	0.835ns	0.795ns	
Sample from wild	25.5-31.4	15	3(20.0)	6(40.0)	2(13.3)
	31.5-35.4	16	8(50.0)	4(25.0)	1(6.3)
	35.5-43.4	15	4(26.7)	1(6.7)	6(40.0)
	43.5-49.4	4	2(50.0)	1(25.0)	0(0.0)
	<b>Total</b>	<b>50</b>	<b>17(34.0)</b>	<b>12(24.0)</b>	<b>9(18.0)</b>
	Chi square		3.951	4.587	7.515
	Df		3	3	3
P value		0.267ns	0.205ns	0.057ns	

## Discussion

This study reveals that the fish were infected with nematode, cestode and trematode which is in conformity with previous researches. This infection was higher in female than in male samples and male than in female samples from concrete pond and wild respectively. Biu *et al.* (2018); Bekele *et al.* (2015) reported higher prevalence in male than female samples of *Clarias gariepinus* in the wild. Williams and Jones (1994) suggested that parasitism differs in various aquatic ecosystems and this is determined by the interaction between biotic and abiotic factors. Fish species in good environmental conditions rarely come down with diseases (Kawe *et al.*, 2016). Reports have recorded helminths to be found in freshwater fishes, at high prevalence which are dependent on factors of parasite species and their biology, host and its feeding habits, presence of intermediate hosts where necessary, hygiene and physical factors of the water body (Doreen *et al.*, 2009; Shukerova *et al.*, 2010; Hussien *et al.*, 2012). This difference in infection between male and female maybe due to differential feeding habit, either quality or quantity of food eaten or as a result of different degree of resistance to infection. The high rate of infection could also be attributed to competition for food, mate partner and territorial defense.

The lower prevalence of infection in female sample from wild could also be as a result of physiological status of female fish in which most gravid female could have increased resistance to infection by parasites. However, this report contradicts with the findings of Enyidi *et al.* (2016); Ambrose *et al.* (2018) and Adegbaaju (2010) who reported higher parasitic infection in female *Clarias gariepinus* from wild than male due to their search for food and egg laying habitat. Female *Clarias gariepinus* from wild had more prevalence of parasite than male. This still agrees with Danyaro (2018) but contradicts with Bekele *et al.* (2015).

Considering the body weight, these results show that fish from concrete pond weighing 93.5-147.4g had the highest prevalence. This conforms to the study of Ibrahim *et al.* (2013) who reported that fishes (juvenile) with low weight have high parasitic infestation than adult which may have acquired immunity which could be attributed to earlier exposure. High prevalence of parasite could also be attributed to high stocking density of pond. Fishes from wild in weight range of 455.5-554.4g had highest prevalence of parasite. This is likely attributed to high stocking density. This report conforms to Iyajiet *et al.* (2014) who reported that fishes with weight 542.5-621.4g had high prevalence of parasite due to large surface area and long duration to accumulate parasite than younger ones which provides more internal and external space for parasite residence and therefore tends to have heavier worm burdens because they eat more parasitized prey (foodchain). However, it contradicts with Ibrahim *et al.* (2013) who recorded high prevalence in fishes with low weight.

Fish with the total length range of 40.5-45.4cm and 30.5-35.4 recorded the highest prevalence (80%) with no significant difference amongst other groups. This report agrees to Adebote *et al.* (2004) and Akinsanya *et al.* (2007), who recorded high prevalence amongst fish with length ranging of 40.0-49.9cm. While fish sample from wild showed high prevalence of parasite in total length range of 31.5-37.4cm and 43.5-49.4 cm. This report shows that there was competition for

food and other resources. In both the wild and pond, it was observed that fishes of standard-length range of 30-40cm were more infected than those with either lower or higher length.

## REFERENCES

- Abuelhassan.E.Y., Atef. I.S. and Jihan, M. R. (2014). The occurrence of *Contraecaecum* sp. Luræ (Nematoda: Anisakidae) in four teleostean species from lake Nasser, Egypt: Morphological and Molecular Studies. *The Journal of Basic and Applied Zoology*. 9:21-27.
- Adegbaju, W. A., Hassan. A. A. and Akinsanya. B. (2010). Haemoparasites of *Clarias gariepinus* and synodontis clarias from Lekki Lagoon, Lagos. Nigeria, *Journal of American Science*, 3:3.
- Akinsanya, B., Otubanjo, O. A. and Hassan, A. A. (2007). Helminth parasites of *Malapterurus electricus* Malapteruridae from Lekki Lagoon, Lagos, Nigeria. *Journal of American Science*, 3(3): 1-5.
- Akinsanya. B., Otunanjo, O.A. and Hassan. A. A. (2007). Helminth parasite of *Malapterurus electricus* from Lekki Lagoon. Lagos, Nigeria. *Journal of American Science*. 3(3):5-10.
- Ambrose. M. and Beatty. V.M. (2018). Survey of gastrointestinal parasites in African Cat fish (*Clarias gariepinus*) in some fish ponds in Kaduna Metropolis, Nigeria. *Integrity Found*. 3(6):184-189.
- Bekele, J. and Daniel, H. (2015). Prevalence of intestinal parasite of *Oreochromis niloticus* and *Clarias gariepinus* fish species in lake Ziway, Ethiopia. *Journal of Aquaculture Research and Development*. 6:2.
- Biu, A. A., Diyaware. M. Y., Yakaka, W. and Jerry, R. D. (2014). Incidence of parasites of *Clarias gariepinus* caught from Lake Alan, Maiduguri, Borno State, Nigeria. *Nigerian Journal of Fisheries and Aquaculture*, 2(1):74-80.
- Doreen, Z. M., Chakanesta, C. and Phumuzile, Y. (2009). Observation on the helminth parasite of fish in Insukamini Dam, Zimbabwe. *Research Journal of Agriculture and Biological Science*, 5(5): 782-785.
- Danyaro. N., Miga, H. L. Bichi, A. H., Dabo, N. T and Maruf, N. (2018). Prevalence of intestinal helminth parasites in *Heterotis niloticus* in marma water channel along River Hadejia, Jigawa Nigeria. *Dutse Journal of Pure and Applied Sciences*, 4:2.
- Emmy-Egbe, I. O. (2013). Faecal Disposition methods and Incidence of intestinal helminth parasites among school children in Ihiala Local Government Area Anambra State, Nigeria. *International Science Resources Journal*, 4(2):81-86.

- Enyidi,U. D. and Eneje, U. I. (2015). Parasite of African catfish *Clarias gariepinus* cultured in homestead ponds. *Journal of Agriculture*, 2:12.
- Gbabiaka, L.A., Akande,T. T. and Ekeocha, C.A. (2017). Assessment of parasites associated with African catfish farmed at Owerri Federal Constituency, Imo State Nigeria. *Agriculture and Biology Journal of North America*, 8(5):168-172.
- Hassan,A. A. Akinsanya, B. and Adegbaju. W. A. (2007). Haemo parasites of *Clariasgariepinus* and *Synodontiselarias* from Lekki Lagoon, Lagos. *Nigeria Journal of American Science*, 3(3):17-19.
- Hussen, A., Tefera, M. and Asrate, S. (2012). Gastrointestinal helminth parasites of *Clarias gariepinus*(Catfish) in Lake Hawassa Ethiopia. *Scientific Journal of Animal Science*, 1(4): 131-136.
- Ibrahim H.Y. and Yahaya, H. (2011). Women participation in homestead fish farming in North central Nigeria. *Livestock Research for Rural Development*. 23: 35.
- Ibrahim,B.Anta.I. K. Badaru, A. A.and A Bdllahi S. A (2010). Occurrence of helimithy on *Clarias gariepinus* (African cat fish) caught in selected points along river Kaduna, Nigeria. *Science World Journal*.14:3.
- Iyaji.F.O.. Eyo, J. Falola, E. and Okpanachi, M. A. (2015). Parasite of *Synodontissorex* (gunther, 1866 mochokidae, siluriformes) in river Niger and Benue at the confluence area in Lokoja, Nigeria. *FutaJournal of Research in Sciences*, 1:87-94.
- Kawe, S. M., God'spower, R. O., Balarabe, M. R. and Akaniru, R. I. (2016). Prevalence of gastrointestinal helminth parasites of *Clarias gariepinus*in Abuja, Nigeria. *Sokoto Journal of Veterinary Sciences*, 14 (2):26-33.
- Kawe.S. M. and Godspower, R. O., Balarabe, M. R. and Akaniru, R.I. (2016). Prevalence of gastrointestinal helminth parasites of *Clarias gariepinus* in Abuja, Nigeria. *Sokoto Journal of Verterinary Science*. 14:2.
- Ndupuh, E.E., Okoye, U.,and Adeleye S. A. (2016). A survey on endo - parasites of *Clarias gariepinus* in some selected fish Farms in Owerri west local government area of Imo State,Nigeria. *International Journal of Fisheries and Aquaculture studies*, 4(5): 624-631.
- Shukerova. S., Kirin, D. and Hanzelova, V. (2010). Endohelminth communities of the perch, *Percfluviatilis*(Perciformes, Percidae) from Srebama Biosphere Reserve, Bulgaria. *Helminthologia*. 42(2): 99-104.
- Williams H & Jones A (1994). Parasitic worms of fish. Taylor and Francis, Bristol, UK. Pp. 593.