

Effects of Organic and Inorganic fertilizers on the growth and yield of Rice (*Oryza sativa* L.)

ABSTRACT

To evaluate the effect of different levels of organic and inorganic fertilizers on the growth and yield of Sin Thu Kha rice variety. The experiment was conducted with randomized complete block (RCB) design by three replications. The experiments were carried out in Water Utilization Research section, Department of Agricultural Research (DAR) at Yezin, Naypyitaw, Myanmar during dry and wet seasons, 2023. The organic sources used were poultry manure and cow dung manure. The treatments were T₁ (Control), T₂ (100% RRF) (Recommended Rate of Fertilizer), T₃ (6 t ha⁻¹ Cow dung manure) (CM), T₄ (6 t ha⁻¹ Poultry Manure) (PM), T₅ (25% RRF + 4.5 t ha⁻¹ CM), T₆ (25% RRF + 4.5 t ha⁻¹ PM), T₇ (50% RRF + 3 t ha⁻¹ CM), T₈ (50% RRF + 3 t ha⁻¹ PM), T₉ (75% RRF + 1.5 t ha⁻¹ CM), T₁₀ (75% RRF + 1.5 t ha⁻¹ PM). It was observed that the grain yields as well as the yield attributing parameters like number of tillers hill⁻¹, panicle length, number of panicles hill⁻¹, number of spikelets panicle⁻¹, filled grain percent were significantly influenced among different treatments except 1000 grain weight. The maximum grain yield was resulted (796.10 g m²) in dry season and (666.35 g m²) in wet season at T₈ treatment (50% RRF + 3 t ha⁻¹ PM). The higher yield of T₈ and T₁₀ were observed in both seasons followed by T₂, T₇ and T₉.

Keywords: *cow dung manure; poultry manure; organic fertilizer.*

1. INTRODUCTION

Rice (*Oryza sativa* L.) is a widely consumed staple food and one of the most important sources of nutrition all over the world human population, especially in Asia and Africa [1]. Inorganic and organic are the main sources for replenishing plant nutrients in agricultural soils [2]. The improper application of synthetic fertilizers has polluted soil and water basins. Organic matter and important plant nutrients are no longer present in the soil. Continuous use of inorganic fertilizers leads to deterioration in soil chemical, physical, and biological properties, and soil health [3]. The negative impacts of chemical fertilizers, coupled with escalating prices, have led to growing interests in the use of organic fertilizers as a source of nutrients [4], [3]. Inorganic fertilizers are also becoming very costly for farmers to apply the fully recommended rates. It uses in several negative health and environmental consequences [5]. In recent years, there has been serious concern about long-term adverse effect of continuous and indiscriminate use of inorganic fertilizers on deterioration of soil structure, soil health and environmental pollution [6], [7] and [8].

Organic manure such as farmyard manure supply a good amount of plant nutrients and contribute to crop yield. It increases the organic matter in the soil. Even the applications of

recommended NPK fertilizers, devoid of organic manure have not sustained soil quality. Soil quality has declined even with the application of recommended NPK fertilizers without organic manure. Therefore, to make the soil well supplied with all the plant nutrients in the readily available form and to maintain good soil health, it is necessary to use organic fertilizer in combination with inorganic fertilizers to obtain optimum yields [9]. Use of organic matter to meet the nutrient requirement of crops would be an inevitable practice in years to come, particularly for resource poor farmers. Therefore, the objectives of this study were to evaluate the effects of different levels of organic and inorganic fertilizers and, their combination effects on the growth and yield of Sinthukha rice variety and to develop a suitable integrated level of organic and inorganic fertilizers.

2. material and methods

The experiment was carried out from January to June (dry season) and from July to November (wet season), 2023. Both experiments were conducted in Water Utilization Research section, Department of Agricultural Research (DAR) at Yezin, Naypyitaw. The experimental design was a Randomized Completely Block Design (RCBD) with ten treatments and three replications. There were 30 experimental units. The concrete tank with the size of (1.8m length × 0.9m breath × 0.5m height) was used as an individual unit. In the second season experiment, the same treatments were applied to the same plots as in the first season. The ten treatments were assigned as Table 1. The tested rice variety was Sin Thu Kha rice variety with 135 days duration. Twenty-one days old seedlings were transplanted at the space of 20 cm × 15 cm. Before conducting the experiment, soil samples were collected for the analysis of some physiochemical properties. The results are shown in Table 2. The air-dried organic manures were applied in each treatment. According to the respective treatments, the Organic manures were well incorporated with the topsoil at the depth of 15 cm, with the date of 14 days before transplanting. Analytical results of two kinds of organic manures samples were described in Table.3. As a regular practice, the experimental units were irrigated whenever necessary. For cultural practices, irrigation, weed control, and pest and disease management practices were done when it was necessary.

2.1 Data collection

Growth parameters such as plant height and number of tillers hill⁻¹ were recorded randomly selected 4 hills from each plot in 2 - weeks interval starting from 14 days after transplanting (DAT) until heading stages. For total dry matter, two plant samples were collected at 25 DAT, 55 DAT, 84 DAT and 110 DAT and dried in the shade and then put in an oven at 65°C ± 5°C for 48 hours. Dry weights were recorded. At harvest, the grain yields were computed from a central 0.72 m² size harvested area in each plot. Four hills were selected as samples to assess the yield component parameters.

Table 1: Treatments of the experiment for both seasons

Treatments	Fertilizers Rates
T ₁	Control
T ₂	100% RRF (N:P: K- 86:18.5:31 kg ha ⁻¹)

T ₃	CM (6 t ha ⁻¹)
T ₄	PM (6 t ha ⁻¹)
T ₅	25% RRF + CM (4.5 t ha ⁻¹)
T ₆	25% RRF + PM (4.5 t ha ⁻¹)
T ₇	50% RRF + CM (3 t ha ⁻¹)
T ₈	50% RRF + PM (3 t ha ⁻¹)
T ₉	75% RRF + CM (1.5 t ha ⁻¹)
T ₁₀	75% RRF + PM (1.5 t ha ⁻¹)

RRF = Recommended Rate of Fertilizer (DAR)
CM = Cowdung Manure
PM = Poultry Manure

Table 2: Measurement of some physicochemical properties of experimental soil

Characteristics	Values	Rating	Analytical Methods
Soil Texture class		Loamy Sand	Pipette method
Soil pH	7.12	Neutral	4A1- 1:5 soil: water suspension
EC (dSm ⁻¹)	0.42	Non- saline	3A1- 1:5 soil: water suspension
Organic matter (%)	0.69 %	Low	Tyurin's method
Available N (mg.Kg ⁻¹)	25.20	Very Low	Alkaline permanganate method
Available P (mg.Kg ⁻¹)	12.87	Medium	9C-Olsen's P-Malachite green
Available K (mg kg ⁻¹)	75.21	Low	1N Ammonium acetate extraction

Table 3: Measurement of chemical composition in both manures of this experiment

Characteristics	Cow dung manure	Poultry manure
Organic Matter (%)	64.90	32.85

Total N (%)	1.47	2.02
Total P ₂ O ₅ (%)	1.0	1.87
Total K ₂ O (%)	2.40	2.83
Total S (%)	0.56	0.84
Moisture (%)	2.06	8.37
Organic Carbon (%)	37.64	19.05
C:N ratio	25.60	9.43

2.2 Statistical analysis

All experimental data were analyzed by using the Analysis of Variance (ANOVA) with Statistix 8 software. The differences in treatment means were separated by Least Significant Difference (LSD) at 5% probability level.

3.1 Growth Parameters

3.1.1 plant height (cm)

Plant height varied significantly both inorganic and organic fertilizers application. Under both seasons, plant height increased progressively in all treatments from 14 DAT to 98 DAT (Figure 1a and b). The results indicated that only the control treatment (T₁) i.e. (87.83 cm) in dry season and (95.25 cm) in wet season) showed lowest plant heights at all growth stages. The treatment T₁₀ (75% RRF + Poultry Manure 1.5 t ha⁻¹) i.e. (115.17 cm) in dry season and (122.07 cm) which combined with poultry manures and chemical fertilizers had a significantly different plant height from the other treatments. A similar result was observed by Abdul- Rahman [10], Moe et al. [11] in ascertaining the combined effect of organic manure and chemical fertilizer on rice. Major nutrient availability was thought to be the cause of the variance in plant height caused by different nutrient sources

Figure 1. Plant height (cm) as affected by organic and inorganic fertilizers in (a) dry season and (b) wet season

3.1.2 Number of tillers hill⁻¹

The number of tillers hill⁻¹ counted at different growth stages between 14 and 98 days after planting, and the count was continuously raised illustrated in Figure 2 (a and b). In the dry season, all treatments showed a significant change at the 1% level, except for 14 DAT. According to the results from T₈ (3 t ha⁻¹ Poultry manure +50% NPK) (15.66) in dry season and (16.67) in wet season gave the maximum number of tillers hill⁻¹ all different growth stages the minimum tiller numbers were recorded from control (T₁) (8.73) and (7.43). In the wet season, the maximum numbers of tillers hill⁻¹ were found from T₈ (3 t ha⁻¹ Poultry manure + 50% NPK) (16.67) which was followed by T₂ (100% NPK only) (15.83) and T₁₀ (1.5 t ha⁻¹ Poultry manure + 50% NPK) (15.83). Abdul-Rahman [10] reported that Combined application of organic and mineral fertilizers produced a higher number of tillers per hill as compared to all other treatments.

Figure 2. Number of tillers per hills as affected by organic and inorganic fertilizers in (a) dry and (b) wet season

3.1.3 Total dry matter

The total dry matter (TDM) in both seasons was recorded at active tillering, panicle initiation, heading, and harvesting stages. The progressive improvements of total dry matter from active tillering to harvesting stages were recorded Figure 3 (a and b). The application of both organic and inorganic fertilizations was significantly different on all collecting growth stages. There were some significant differences on other growth stages due to combined application of organic and inorganic fertilizers. The maximum total dry matter was produced by (T₁₀) (75%RRF + Poultry Manure 1.5 t ha⁻¹). Belay et al. [12] who demonstrated that treatment with poultry manure combined with inorganic fertilizers enhances nutrient availability and creates suitable conditions for growth by reducing nutrient loss and increasing plant dry weight. Organic manures are slow releasing N source found beneficial during subsequent stages of crop, which might have resulted in increasing the total dry matter. Similar results were reported by Moe et al. [11].

Figure 3. Total dry matter (g hill⁻¹) at different growth stages as affected by organic and inorganic fertilizers applications in (a) dry season and (b) wet season

3.2 Yield and Yield Component characters

3.2.1 Number of panicles hill⁻¹

Number of panicles hill⁻¹ were highly statistically significant difference among treatments at < 0.001 (Table 4 and 5). The maximum number of panicles per hill (14.00) in dry season and (10.43) in wet season respectively were found in combined application of poultry manure and NPK fertilizer treatment, T₈ (3 t ha⁻¹ Poultry manure + 50% NPK). The state in which no fertilizer from T₁ (control) (8.46) in dry season and (7.50) in wet season resulted in the lowest number of panicles per hill. According to Moe et al. [11] combined application of organic and mineral fertilizers produced a higher number of panicles per hill as compared to all other treatments.

3.2.2 Number of spikelets panicle⁻¹

Effect of organic and inorganic fertilizers on the number of spikelets per panicle of rice showed highly significant differences among the treatments at 1% level Table (4 and 5). In dry seasons, the highest number of spikelets panicle⁻¹ was produced by T₈, T₁₀ and T₂ treatments (175.67), (174.75) and (173.11) respectively. In wet season maximum number of spikelets per panicle gave T₈ (166.74). The minimum number of spikelets per panicle⁻¹ was recorded from the control treatment T₁ (Control) which was significantly different from other treatments in both seasons. Number of spikelets panicle⁻¹ is the most important factor to maximize yield of rice which could be increased by combined application of organic manure and chemical fertilizer compared to the sole application or control treatment [13]. According to Mahmud [14], Sarker [15] organic fertilizers combined with inorganic fertilizer have been shown to improve soil structure, nutrient

exchange and maintain soil health, thus providing higher grain yield by achieving a greater number of spikelets panicle¹.

3.2.3 Percentage of Filled grain (%)

A highly significant difference in the filled grain percentage was observed in both seasons when the SinThKha variety was treated different rates of organic and inorganic fertilizers. There was a highly significant effect on filled grain (%) among different treatments. In dry season, the greatest percentage of filled grain came from T₈ (3 t ha⁻¹ poultry manure + 50% NPK) (83.14) although T₁₀ (1.5 t ha⁻¹ poultry manure + 75% NPK) (82.79) and T₂ (NPK only) (82.43) did not differ significantly from one another. In wet season, (77.23) was produced by T₈ (3 t ha⁻¹ poultry manure + 50% NPK) treatment. The lowest filled grain (%) was obtained from T₁ (control) (62.66) and (51.57) In both seasons. Arief et al. [16] were observed that NPK fertilizers combined with poultry manure resulting in greater number of filled grain per panicle of rice.

3.2.4 1000 Grain - Weight (g)

Results showed that there was no significant difference in the effects of organic and inorganic fertilizers application on 1000-grain weight of Sin Thu Kha rice. In both seasons the highest 1000-grain weight was recorded from T₈ (3 t ha⁻¹ Poultry Manure+ 50% NPK) (18.64g) in dry season and (19.76g) in wet season. The lowest 1000-grain weight was found in both seasons from (T₁) Control treatment (17.47g) in dry season and (19.10 g) in wet season as shows in Table (4 and 5). This result was in conformity of the finding of Yang et al. [17] who recorded that 1000-grain weight was increased by the application of chemical fertilizer along with organic manure.

3.2.5 Panicle Length

The combined application of organic and inorganic fertilizers gave significant panicle length of SinThuKha rice in both seasons in Table (4 and 5). The panicle length of rice ranges from (25.90 cm) in dry season to 24.11 cm in wet season gave the maximum panicle length from T₈ (3 t ha⁻¹ Poultry Manure+ 50% NPK) whereas the minimum panicle length (20.18cm) and (21.54 cm) was observed in T₁. According to Kustera [18] panicle length increased with the application of organic and chemical fertilizers, and this could be due to the increase in the absorption of available nutrients. Alim [19] reported that the length of the panicle is one component that influences the outcome of the results of the rice plant, so the longer panicles potential results obtained tend to be larger than the shorter panicles.

3.2.6 Grain yield (g/m²)

Grain yield of rice as affected by the application of organic and inorganic fertilizers in both seasons. There were highly significant differences among the treatments at 1% level. It was observed that the treatment T₈ (3 t ha⁻¹ Poultry Manure + 50% NPK) gave the highest grain yield (796.10g) in dry and (666.4 g) in wet seasons. Similar yield was observed from treatment T₁₀ (1.5 t ha⁻¹ poultry manure + 75% NPK) in both seasons. In both seasons the lowest grain yield was given by the control treatment T₁ ranges from (324.83 g) to (317.2 g). Combined application of manure and chemical fertilizer at different levels on rice plants greatly influenced the grain yield due to the increased in plant height, tiller number, panicle length, grains per panicle and 1000 seed weight in a study by Alam [20]. This may have resulted from the presence of

important nutrient sources for plant development and productivity. According to [14] by Mahmud, organic fertilizers supply nutrients through microbial activity, whereas chemical fertilizers make nutrients rapidly available by allowing them to be easily accessed in soil solution. Similarly, it was reported by [20, 8] Alam and Singh that applying nitrogenous fertilizer and organic manure combined application enhanced rice grain production.

Table 4: Mean value of yield and yield components parameters of rice as affected by organic and inorganic fertilizers during dry season, 2023

Treatments	No. of Panicle hill ¹	No. of Spikelet panicle ¹	Filled grain %	1000-grain weight (g)	Panicle Length(cm)	Yield (g/m ²)
T ₁	8.46 d	104.49 e	62.66 d	17.47	20.18 f	324.83 g
T ₂	13.16 ab	173.11 a	82.43 a	18.32	24.66 ab	720.13ab
T ₃	10.42 d	127.10 d	67.98cd	18.27	22.69 de	429.90 ef
T ₄	11.0 cd	129.63cd	68.86 bcd	17.92	22.50 e	453.34def
T ₅	12.0 bc	133.37 c	69.93 bcd	17.96	23.09 cde	488.69 e
T ₆	12.33 bc	133.85 c	78.32 abc	17.90	23.74 bcd	557.42 de
T ₇	12.84 ab	154.02 b	78.96 ab	18.30	24.02 bc	621.91 bc
T ₈	14.0 a	175.67 a	83.14 a	18.64	25.90 a	796.10 a
T ₉	13.0 ab	163.52ab	79.47 ab	17.92	24.44 b	702.96 b
T ₁₀	13.26 ab	174.75 a	82.79 a	18.32	24.66 ab	762.51 a
LSD _{0.05}	1.49	17.61	11.09	1.01	1.19	108.71
Pr>F	**	**	**	ns	**	**
CV%	7.21	6.95	8.57	3.28	2.94	10.78

Means followed by different letters in the same column are not significantly different by LSD test at 5% level. * significant difference at 5% level, **significant difference at 1% level, ^{ns} non-significant difference

Treatments	No. of panicles hill ¹	No. of spikelets panicle ⁻¹	Filled grain %	1000-grain weight (g)	Panicle length (cm)	Yield (g/m ²)
T ₁	7.50 e	80.53 f	51.57 c	19.10	21.54 c	317.2 e
T ₂	9.58 abc	142.72 b	68.91 ab	19.50	24.35 a	649.7ab
T ₃	8.26 cde	111.85 de	62.02 bc	18.87	21.63 c	415.4 d
T ₄	7.66 de	98.10 ef	64.227 b	19.44	21.71 c	433.5 d

T ₅	9.08 abcd	118.10 cd	60.91 bc	19.14	22.31 bc	445.7 cd
T ₆	9.2 abc	132.48 bc	66.87 ab	19.18	22.75 abc	545.9 bc
T ₇	8.75 bcde	149.37 ab	68.37 ab	19.19	23.75 ab	583.1ab
T ₈	10.43 a	166.74 a	77.23 a	19.76	24.11 a	666.4 a
T ₉	9.58 abc	139.76 b	66.32 b	19.25	22.86 abc	596.7ab
T ₁₀	10.00 ab	166.27 a	71.17 ab	19.42	23.65 ab	664.5 a
LSD _{0.05}	1.57	18.30	10.56	0.84	1.74	111.43
Pr>F	*	**	**	ns	*	**
CV%	10.19	8.17	9.37	2.54	4.45	12.17

Table 5: Mean value of yield and yield components parameters of rice as affected by organic and inorganic fertilizers during wet season, 2023

Means followed by different letters in the same column are not significantly different by LSD test at 5% level. * significant difference at 5% level, **significant difference at 1% level, ^{ns} non-significant difference

4. Conclusions

From the results of the experiment, it may be concluded that combined application of (50% RRF and poultry manure 3 t ha⁻¹) were significantly influenced among different treatments that the grain yields as well as the yield attributing parameters such as number of tillers hill⁻¹, panicle length, number of panicles hill⁻¹, number of spikelets panicle⁻¹, filled grain percent except 1000 grain weight. The maximum grain yield was resulted (796.10 g m²) in dry season and (666.35 g m²) in wet season at T₈ treatment (50% RRF + 3 t ha⁻¹ PM). The higher yield of T₈ and T₁₀ were observed in both seasons followed by T₂, T₇ and T₉. From the economic point of view, farmers can use the combination of organic manures and reduced rate of inorganic fertilizers to boost the yield of rice as well as to maintain and improve soil health. In summary, the combination of organic and inorganic fertilizers can reduce the use of inorganic fertilizers. Other combination of organic manures and chemicals fertilizer may be necessary to be tested for obtaining good agricultural practice for sustainable rice production in Myanmar.

REFERENCES

- Liu W, Liu J, Triplett L, Leach JE, Wang GL. Novel insights into rice innate immunity against bacterial and fungal pathogens. Annual review of phytopathology. 2014 Aug 4;52:213-41.
- Masarirambi MT, Mandisodza FC, Mashingaidze AB, Bhebhe E. Influence of plant population and seed tuber size on growth and yield components of potato (*Solanum tuberosum*). Int. J. Agric. Biol. 2012 Aug 1;14(4):545-549.

- Mahajan A, Bhagat RM, Gupta RD. Integrated nutrient management in sustainable rice-wheat cropping system for food security in India. SAARC Journal of Agriculture; 2008;6(2):149- 163.
- Satyanarayana V, Prasad PV, Murthy VRK, Boote KJ. Influence of integrated use of farmyard manure and inorganic fertilizers on yield and yield components of irrigated lowland rice. Journal of plant nutrition;2002; 25:2081- 2090.
- Sharada. P, Sujathamma P. Effect of Organic and Inorganic Fertilizers on the Quantitative and Qualitative Parameters of Rice (*Oriza Sativa L.*). Curr Agri Res; 2018;6(2).
- Ghosh BC, Bhat R. Environmental hazards of nitrogen loading in wetland rice fields. Environ Pollut; 1998; 102:123-126.
- Shukla BD, Misra AK, Gupta R.K. Application of nitrogen in production and post-production systems of agriculture and its effect on environment in India. Environ. Pollut. 1998;102: 115-122.
- Singh RB. Environmental consequences of agricultural development: a case study for the Green Revolution State of Haryana, India. Agr. Ecosys. Environ; 2000;82: 97-103.
- Ramalakshmi ChS, Rao PC, Sreelatha T, Mahadevi M, Padmaja G, Rao PV, Sireesha A. Nitrogen use efficiency and production efficiency of rice under rice-pulse cropping system with integrated nutrient management. Journal of Rice Research; 2012; 5(1&2):42-51.
- Abdul-Rahman, A.B.I. The combined effect of organic and inorganic fertilizers on growth of rice plants. United Nations University Land Restoration Training Programme [finalproject] <https://www.grocentre.is/static/gro/publication/719/document/abdul;2019>.
- Moe, K., Moh, S.M. and Htwe, A.Z. Effects of integrated organic and inorganic fertilizers on yield and growth parameters of rice varieties. Rice Sci; 2019;26(5): 309-318rahman2019.pdf
- Belay, A., Classens, A.S., Wehner, F.C. and De, B.J.M. Influence of Residual Manure on Selected Nutrient Elements and Microbial Composition of Soil under Long-Term Crop Rotation. South African Journal of Plant and Soil;2001;18: 1-6.
- Siavoshi, M., Nasiri, A., Lawere, S.L. Effect of organic fertilizer on growth and yield components in rice (*Oryza sativa L.*). J. Agril. Sci;2011;3: 217-22
- Mahmud, A.J., Shamsuddoha, A.T.M. and Haque, M.N. Effect of Organic and Inorganic Fertilizer on the Growth and Yield of Rice (*Oryza sativa L.*). Nat. Sci. 2016;14(2): 45-54.
- Sarker, D., Mazumder, S., Kundu, S., Akter, F. and Paul, S.K. Effect of poultry manure incorporated with nitrogenous and sulfur fertilizer on the growth, yield, chlorophyll and nutrient contents of rice var. BRRI dhan33. Bangladesh Agron. J. 2015;18(1): 99-111.
- Arief, M., M. Tasneem., F. Bashir, G. Yaseen, and R.M. Iqbal. Effect of integrated use of organic manures and inorganic fertilizers on yield and yield components of rice. Journal of Agricultural Research. 2014;52(2): 200-203.
- Yang, C.M., Yang, L., Yang, Y. and Ouyang, Z.. Rice root growth and nutrient uptake as influenced by organic manure in continuously and alternately flooded paddy soils. Agricultural Water Management, 2004;70(1): 67-81.
- A. Kustera, "Keragaman Genotipe dan Fenotipe Galur-Galur Padi Hibrida Di Desa Kahuman, Polanharjo Klaten," Skripsi. Jurusan/Program Studi Agronomi. Fakultas Pertanian. Universitas Sebelas Maret. Surakarta;2008.
- Alim, M. A. "Effect of organic and inorganic sources and doses of nitrogen fertilizer on the yield of boro rice." Journal of Environmental Science and Natural Resources 5, no. 1 2012: 273-282.
- Alam, M.M., Arifunnahar, M. and Khatun, M.M. Effect of different organic and inorganic fertilizers on the growth and yield of boro rice (BRRI dhan29). Int. J. Res. Sci. Innov. (IJRSI). 2021;8(1): 176-180.

