

# Elucidating drought tolerance potential of maize (*Zea mays* L.) inbred lines through polyethylene glycol induced drought stress

## Abstract

Drought is one of the most distressing environmental stresses restraining the crop production. Drought undermines the plant growth from seedling to maturity and the studies have unveiled that the harmful impact of drought results in crops due to the damages perceived during crucial stages of development namely, germination, seedling development and flowering stages. Biometric elements and indices at an early growth stage could be employed in designing the selection criteria to figure out the drought tolerant genotypes. Breeding for drought-prone environments is constrained by lack of suitable selection indices of drought stress resistance. The present study was conducted to determine the reliability of *in-vitro* screening method for initiating drought breeding programme. Poly ethylene glycol (PEG) has been used often as abiotic stress inducer in many studies to screen drought tolerant germplasm. The present investigation was carried out to reveal the drought tolerance potential of 103 maize inbred lines by *in-vitro* screening under three levels of PEG levels viz., 0% as control, 10% and 20% and evaluating their effect on the root parameters like primary root length, number of seminal roots, number of lateral roots and root biomass. The seedlings exhibited a significant variation for all the traits analyzed. All the root parameters had highest value under control and had significant decline with increasing PEG concentrations (0% < 10% < 20%). The inbreds showing the best level of drought tolerance at all levels of PEG induced stress can be used as a source of drought tolerance for the improvement of drought tolerant hybrids. The variation among maize inbreds for these traits was found to be an ideal indicator to screen the drought tolerant genotypes at early growth stages.

Keywords: Drought, drought tolerance, PEG (polyethylene glycol), primary root length, lateral roots, root biomass

## Introduction

“Maize (*Zea mays* L.) is one of the important food and industrial crops grown extensively in major part of the world and it ranks third after wheat and rice” (Godfray et al., 2010 and Kyenpia et al., 2009). “Maize is of worldwide importance as a food, feed and as a source of diverse industrially important products” (Prasanna, 2012). “Due to the growing demand for dairy and meat products in developing countries and the decline in rice production in China and India, maize has been projected to become the most important crop by 2030” (Salvi et al., 2007).

“Maize is essential for global food security. The current trends of climatic changes increase water scarcity and reduce maize productivity by 15- 30%” (Magar *et al.*, 2019). There are several abiotic factors limiting maize production in different parts of the world. Drought is one of the most distressing environmental stresses restraining the crop production. These trends, coupled with an expansion of cropping into marginal production areas, are generating increasingly drought-prone maize production environments and persuades the need for drought tolerance in maize crop.

“Drought undermines the plant growth from seedling to maturity and the studies have unveiled that the harmful impact of drought results in crops due to the damages perceived during crucial stages of development namely, germination, seedling development and flowering stages” (Nivethitha *et al.*, 2020). Biometric elements and indices at an early growth stage could be employed in designing the selection criteria to figure out the drought tolerant genotypes. Breeding for drought-prone environments is constrained by lack of suitable selection indices of drought stress resistance.

“Polyethylene glycol (PEG-6000) treatment can be applied to identify local maize varieties as sensitive or tolerant to drought stress before germination” (Mustamuet *et al.*, 2023). “This method can be used to identify the best varieties for drought-prone areas and help in conservation efforts. PEG concentrations could act as a limiting factor by affecting maize plant growth during the germination and seedling stages” (Dani and Siswoyo, 2019 and Petcu *et al.*, 2018). “The upsurge in concentration of PEG caused a decrease in germination percentage, seedling vigour in certain crop plants” (Khodarahmpour, 2011). “Poly ethylene glycol (PEG) has been used often as abiotic stress inducer in many studies to screen drought tolerant germplasm” (Ahmad *et al.*, 2015; Jatoi *et al.*, 2014)). “The germplasm which is showing better performance can be considered as drought tolerant. It is an established fact that tolerance at maturity is demonstrated by the tolerance at immature stage of plant. Therefore, it is necessary to develop effective screening criteria at early growth stages to get the maximum yield” (Tahir *et al.*, 2023). The present study was conducted to determine the reliability of *in-vitro* screening method for initiating drought breeding programme. The aim of the present study was to investigate the effects of PEG induced stress on root traits of maize (*Zea mays*L.) inbreds to screen them for drought tolerance.

## **Materials and Methods**

This study was carried out at Division of Genetics and Plant Breeding, SKUAST-K. 103 maize (*Zea mays* L.) inbreds were used to study the effect of PEG induced stress on root traits to screen them for drought tolerance. In this experiment with polyethylene glycol, PEG-6000 (HIMEDIA) was used in three concentrations viz., control (0%), 10% and 20%. Four seeds of each genotype were surface sterilized with 0.5% NaOCl for one minute, rinsed thoroughly with distilled water and were put in petri plates containing moist filter paper with different

concentrations of PEG-6000 and allowed to germinate in a germinator at 25<sup>0</sup>C and 75% humidity in darkness. Primary root length, number of seminals, number of laterals and root biomass was measured after seven days. The design used was factorial CRD with three replications.

## Results and Discussion

The results of this study reveal that different concentrations of PEG-6000 (0-20%) had significant effect on the root traits of maize inbreds (Table-1). Analysis of variance and mean comparison revealed that there were significant differences between drought stress levels and genotypes. Analysis of variance for various root traits scored under PEG 6000 shows that mean square due to genotypes, PEG levels and genotypes x PEG levels was significant for all the traits (Table 2). The mean performance results also revealed that the root traits under different stress levels were different.

The results of *in-vitro* screening of maize inbreds under three levels of PEG levels *viz.*, 0% as control, 10% and 20% are explained below under appropriate headings:

### Primary Root length (cm):-

The data on primary root length under different levels of PEG-6000 was;

- **0% Level (Control):-** Under controlled conditions primary root length had a mean value of 13.42 with highest value recorded in IMR- 55 and IMR- 416 (19.00) followed by IMR- 566 and IMR- 577 (18.33) each and was lowest in IMR- 4 and IMR- 43(7.00).
- **10% Level:-** Under 10% the primary root length had a mean value of 7.16 with highest value recorded in IMR- 7 and IMR- A (13.00) followed by IMR- 671 (12.00) and IMR- 5, IMR- 11, IMR- 566, IMR- 579 and IMR- B (11.00) and was lowest in IMR- 4, IMR- 76 and IMR- 220(3.33).
- **20% Level:-** Under 20% the primary root length had a mean value of 2.92 with highest value recorded in IMR- 7 (6.00) followed by IMR- A (5.67) and IMR- 116, IMR- 566 and IMR- 671 (5.00) and was lowest in IMR- 4, IMR- 43, IMR- 76, IMR- 128, IMR- 220, IMR- 445 and IMR- 541(1.33).

### Number of Seminals:

The data on number of seminals under different levels of PEG-6000 was;

- **0% Level (Control):-** Under controlled conditions the number of seminals had a mean value of 5.86 with highest value recorded in IMR- 60 and IMR- 667 (9.00) followed by IMR- 565(8.67) and IMR- 134 and IMR- A (8.33) and was lowest in IMR- 292(3.33).
- **10% Level:-** Under 10% the number of seminals had a mean value of 3.92 with highest value recorded in IMR- 60 and IMR- 565 (7.00) followed by IMR- 151(6.67) and IMR- 134, IMR- 385, IMR- 576 and IMR- A(6.33) each and and was lowest in IMR- 11, IMR- 41, IMR- 87, IMR- 116, IMR- 217, IMR- 227, IMR- 245, IMR- 292, IMR- 368, IMR-

376, IMR- 413, IMR- 419, IMR- 579(2.00).

- **20% Level:-** Under 20% the number of seminals had a mean value of 2.05 with highest value recorded in IMR- 439 and IMR- 667(4.00) followed by IMR- 134, IMR- 385, IMR- 565 (3.33) and IMR- 7, IMR- 98, IMR- 268, IMR- 569, IMR- 580, IMR- 609 (3.00) and was lowest in IMR- 572(1.00).

### **Number of Laterals**

The data on number of laterals under different levels of PEG-6000 was;

- **0% Level (Control):-** Under controlled conditions the number of laterals had a mean value of 29.77 with highest value recorded in IMR- 565 (63.00) followed by IMR- 155 (62.67) and IMR- 60 (62.00) and was lowest in IMR- 247(9.67).
- **10% Level:-** Under 10% the number of laterals had a mean value of 16.64 with highest value recorded in IMR- 565(42.00) followed by IMR- 63, IMR- B (37.00) and IMR- 50, IMR- 60(36.00) and was lowest in IMR- 224 (2.33).
- **20% Level:-** Under 20% the number of laterals had a mean value of 8.04 with highest value recorded in IMR- 565 (24.00) followed by IMR- 63 (21.00) and IMR- 8, IMR- 50, IMR- 60 (19.00) and was lowest in IMR-224, IMR-268 and IMR-386 (1.33).

### **Root Biomass(g):-**

The data on root biomass under different levels of PEG-6000 was

- **0% Level (Control):-** Under controlled conditions the root biomass had a mean value of 0.37 with highest value recorded in IMR- 63(0.63) followed by IMR- 384 (0.62) and IMR- 121 and IMR- 574 (0.61) and was lowest in IMR- 20 and IMR- 248(0.16).
- **10% Level:-** Under 10% the root biomass had a mean value of 0.28 with highest value recorded in IMR- 574 (0.56) followed by IMR- 671(0.53) and IMR- 60 (0.52) and was lowest in IMR- 20(0.09).
- **20% Level:-** Under 20% the root biomass had a mean value of 0.17 with highest value recorded in IMR- 671(0.44) followed by IMR- 11, IMR- 574 (0.43) and IMR- 121(0.41) and was lowest in IMR- 152 (0.02).

In the present study all the root parameters including primary root length, number of seminals, number of laterals and root biomass decreased with increasing PEG concentrations from 0-20%. All the root parameters had highest value under control and had significant decline effect with increasing PEG concentrations (0% < 10% < 20%). Similar results were reported by (Fatima *et al.*, 2023) in maize who reported that increasing PEG concentrations led to a decline in key parameters such as germination rate, root elongation and shoot development. Bukhari, *et al.* (2021) reported that the concentration of PEG 6000 affected the response of corn varieties to germination and seed vigor. Polyethylene glycol (PEG) creates osmotic stress and could be used to examine the effect of water stress on seed germination (Hellal *et al.*, 2018). Polyethylene

glycol is the best solute that is known of imposing a low water stress that is reflective of the type of stress imposed by a drying soil (Badr *et al.*, 2020). The effect of PEG in physiological processes has also been reported in maize by (Majid *et al.*, 2020). In a water deficit environment, development of a crop mainly depends on the germination of seeds and the establishment of seedlings (Zafar *et al.*, 2015). The selection method against drought stress using PEG in the vegetative phase was used to simulate maize tolerance to drought stress in the vegetative phase (Suhartono, 2021). (The germination process consists of enzymatic hydrolysis of stored food and the formation of new tissues (Alvi *et al.*, 2022). Therefore, it is necessary to develop effective screening criteria at early growth stages to get the maximum yield (Tahir *et al.*, 2023). It is an established fact that tolerance at maturity is demonstrated by the tolerance at immature stage of plant. Root system with the ability of better growth under stress conditions can be considered as tolerant germplasm (Abdel-Raheem *et al.*, 2007). Thus there is a scope to identify genotypes that have tolerance to drought at the primary growth stage. In our study, IMR- 7 recorded lowest reduction in primary root length, IMR- 439 and IMR- 667 recorded lowest reduction in number of seminal roots, IMR- 565 recorded lowest reduction in number of laterals and IMR- 671 recorded lowest reduction in root biomass across different concentrations of PEG-6000. Such inbreds having genetic potential to maintain the higher growth under stress conditions were identified as drought tolerant.

## **Conclusion**

The variation among maize inbreds for these traits in the present study was found to be an ideal indicator to screen the drought tolerant genotypes at early growth stages. These results have significant implications for increasing maize production in drought prone areas. The inbreds showing the better level of drought tolerance at all levels of PEG induced stress could be used as a source of drought tolerance for the improvement of drought tolerant hybrids .

UNDU

**Table-1: *In vitro* response of maize (*Zea mays* L.) inbreds to different levels of PEG 6000**

| Inbreds | Primary root length(cm) |       |      | Number of Seminals |      |      | Number of Laterals |       |       | Root Biomass(g) |      |      |
|---------|-------------------------|-------|------|--------------------|------|------|--------------------|-------|-------|-----------------|------|------|
|         | Control                 | 10%   | 20%  | Control            | 10%  | 20 % | Control            | 10%   | 20%   | Control         | 10%  | 20%  |
| IMR 3   | 15.00                   | 5.33  | 2.33 | 6.00               | 4.00 | 1.67 | 26.00              | 12.00 | 6.00  | 0.21            | 0.12 | 0.03 |
| IMR 4   | 7.00                    | 3.33  | 1.33 | 7.00               | 5.00 | 3.00 | 22.00              | 13.00 | 9.00  | 0.25            | 0.17 | 0.06 |
| IMR 5   | 17.00                   | 11.00 | 3.33 | 6.00               | 4.00 | 2.00 | 18.00              | 9.00  | 2.00  | 0.54            | 0.47 | 0.41 |
| IMR 7   | 18.00                   | 13.00 | 6.00 | 8.00               | 6.00 | 3.00 | 25.00              | 16.00 | 7.00  | 0.45            | 0.38 | 0.25 |
| IMR 8   | 11.00                   | 6.67  | 2.67 | 6.00               | 4.00 | 1.67 | 45.00              | 34.00 | 19.00 | 0.37            | 0.29 | 0.18 |
| IMR 11  | 15.67                   | 11.00 | 3.67 | 4.00               | 2.33 | 1.33 | 33.00              | 21.00 | 10.00 | 0.57            | 0.51 | 0.43 |
| IMR 19  | 16.00                   | 6.67  | 2.67 | 5.00               | 3.00 | 1.33 | 31.00              | 20.00 | 11.00 | 0.24            | 0.14 | 0.08 |
| IMR 20  | 10.00                   | 4.33  | 1.67 | 4.67               | 2.67 | 1.33 | 24.00              | 13.00 | 5.00  | 0.17            | 0.09 | 0.04 |
| IMR 26  | 12.00                   | 5.67  | 2.33 | 3.67               | 2.67 | 1.67 | 15.00              | 9.00  | 3.00  | 0.34            | 0.26 | 0.13 |
| IMR 34  | 11.00                   | 6.33  | 2.00 | 6.33               | 4.33 | 2.33 | 32.00              | 21.00 | 10.00 | 0.34            | 0.28 | 0.18 |
| IMR 41  | 16.00                   | 8.33  | 3.33 | 4.00               | 2.33 | 1.00 | 18.00              | 11.00 | 6.00  | 0.23            | 0.16 | 0.09 |
| IMR 42  | 11.00                   | 6.00  | 2.33 | 5.67               | 3.67 | 1.67 | 20.00              | 10.00 | 5.00  | 0.52            | 0.43 | 0.31 |
| IMR 43  | 7.00                    | 3.67  | 1.33 | 4.67               | 2.67 | 1.33 | 25.00              | 14.00 | 7.00  | 0.45            | 0.36 | 0.21 |
| 1MR 50  | 11.00                   | 5.33  | 2.33 | 4.67               | 2.67 | 1.33 | 50.00              | 36.00 | 19.00 | 0.31            | 0.25 | 0.13 |
| 1MR 55  | 19.00                   | 9.33  | 3.00 | 5.33               | 3.33 | 1.33 | 25.00              | 13.00 | 7.00  | 0.20            | 0.13 | 0.08 |
| 1MR 60  | 14.00                   | 10.00 | 4.67 | 9.00               | 7.00 | 4.67 | 62.00              | 36.00 | 19.00 | 0.30            | 0.20 | 0.12 |
| 1MR 63  | 14.00                   | 8.67  | 3.67 | 6.00               | 4.00 | 2.33 | 58.33              | 37.00 | 21.00 | 0.63            | 0.52 | 0.43 |
| 1MR 76  | 8.00                    | 3.33  | 1.33 | 6.00               | 4.00 | 2.00 | 33.00              | 24.00 | 13.00 | 0.23            | 0.16 | 0.09 |
| 1MR 77  | 17.00                   | 8.00  | 3.00 | 7.00               | 4.67 | 2.33 | 36.00              | 26.00 | 11.00 | 0.40            | 0.31 | 0.15 |
| 1MR 82  | 15.00                   | 6.67  | 2.33 | 7.33               | 5.33 | 2.67 | 40.00              | 32.00 | 15.00 | 0.26            | 0.19 | 0.11 |
| 1MR 87  | 14.00                   | 6.67  | 2.67 | 4.33               | 2.33 | 1.33 | 22.00              | 11.67 | 5.00  | 0.54            | 0.45 | 0.24 |
| 1MR 91  | 13.00                   | 6.33  | 2.33 | 4.33               | 2.67 | 1.33 | 33.00              | 21.00 | 10.00 | 0.20            | 0.13 | 0.03 |
| 1MR 97  | 17.00                   | 9.67  | 4.00 | 5.00               | 3.00 | 1.33 | 42.00              | 25.00 | 12.00 | 0.54            | 0.47 | 0.24 |
| 1MR 98  | 16.00                   | 7.00  | 3.33 | 8.00               | 6.00 | 3.00 | 34.00              | 21.00 | 10.00 | 0.47            | 0.36 | 0.25 |

|         |       |      |      |      |      |      |       |       |       |      |      |      |
|---------|-------|------|------|------|------|------|-------|-------|-------|------|------|------|
| 1MR 103 | 12.00 | 6.67 | 3.00 | 7.00 | 5.33 | 2.33 | 48.00 | 32.00 | 14.00 | 0.37 | 0.28 | 0.20 |
| 1MR 114 | 16.00 | 9.67 | 4.33 | 6.67 | 3.67 | 1.33 | 26.00 | 16.00 | 11.00 | 0.45 | 0.39 | 0.25 |
| 1MR 115 | 16.00 | 8.33 | 4.00 | 5.67 | 3.67 | 1.67 | 35.00 | 12.00 | 4.00  | 0.39 | 0.25 | 0.12 |
| 1MR 116 | 18.00 | 9.67 | 5.00 | 4.33 | 2.33 | 1.33 | 44.00 | 28.00 | 14.00 | 0.59 | 0.50 | 0.33 |
| 1MR 121 | 13.00 | 8.00 | 4.33 | 5.00 | 4.00 | 2.00 | 42.00 | 24.00 | 11.00 | 0.61 | 0.51 | 0.41 |
| 1MR 127 | 15.00 | 9.00 | 3.33 | 4.33 | 3.33 | 1.33 | 30.00 | 18.00 | 8.00  | 0.23 | 0.17 | 0.11 |
| 1MR 128 | 9.00  | 4.33 | 1.33 | 6.67 | 5.00 | 2.33 | 45.00 | 31.00 | 18.00 | 0.39 | 0.31 | 0.14 |
| 1MR 132 | 10.00 | 5.00 | 3.00 | 5.67 | 3.00 | 1.67 | 24.00 | 11.00 | 5.00  | 0.25 | 0.18 | 0.11 |
| 1MR 133 | 14.00 | 4.33 | 1.67 | 6.67 | 4.67 | 2.67 | 23.00 | 12.00 | 4.00  | 0.53 | 0.47 | 0.26 |
| 1MR 134 | 12.00 | 7.33 | 3.67 | 8.33 | 6.33 | 3.33 | 34.00 | 22.00 | 12.00 | 0.44 | 0.37 | 0.22 |
| 1MR 137 | 15.33 | 7.67 | 2.67 | 5.00 | 3.00 | 1.33 | 22.00 | 11.00 | 6.00  | 0.53 | 0.48 | 0.30 |
| 1MR 145 | 9.67  | 5.67 | 2.33 | 7.33 | 5.33 | 2.33 | 32.00 | 16.00 | 11.00 | 0.46 | 0.38 | 0.28 |
| 1MR 149 | 10.67 | 5.33 | 2.00 | 6.33 | 4.67 | 2.33 | 38.00 | 22.67 | 13.00 | 0.35 | 0.28 | 0.16 |
| 1MR 151 | 9.00  | 5.00 | 1.67 | 9.00 | 6.67 | 2.67 | 40.00 | 23.00 | 9.00  | 0.44 | 0.34 | 0.21 |
| 1MR 152 | 16.00 | 5.00 | 2.67 | 6.00 | 5.00 | 2.00 | 25.67 | 10.67 | 3.67  | 0.21 | 0.12 | 0.02 |
| 1MR 155 | 17.33 | 8.33 | 3.33 | 7.67 | 4.67 | 2.67 | 62.67 | 29.33 | 13.00 | 0.29 | 0.19 | 0.09 |
| 1MR 209 | 12.00 | 4.67 | 1.67 | 5.67 | 3.67 | 2.67 | 36.67 | 20.00 | 6.00  | 0.39 | 0.22 | 0.11 |
| 1MR 217 | 11.00 | 6.00 | 3.33 | 4.33 | 2.33 | 1.33 | 42.67 | 20.67 | 8.67  | 0.45 | 0.30 | 0.15 |
| 1MR 220 | 8.67  | 3.33 | 1.33 | 5.33 | 3.33 | 2.33 | 27.67 | 11.67 | 4.00  | 0.23 | 0.15 | 0.05 |
| 1MR 222 | 8.00  | 4.33 | 1.67 | 4.67 | 2.67 | 1.67 | 34.67 | 14.67 | 5.00  | 0.24 | 0.15 | 0.05 |
| 1MR 223 | 10.33 | 4.67 | 1.67 | 4.67 | 2.67 | 1.67 | 21.33 | 7.33  | 2.33  | 0.51 | 0.37 | 0.17 |
| 1MR 224 | 13.67 | 7.33 | 3.33 | 4.67 | 3.00 | 2.00 | 10.00 | 2.33  | 1.33  | 0.37 | 0.24 | 0.09 |
| 1MR 227 | 12.33 | 7.67 | 4.67 | 4.33 | 2.33 | 1.33 | 13.33 | 6.00  | 1.67  | 0.30 | 0.16 | 0.08 |
| 1MR 230 | 13.33 | 5.67 | 2.67 | 5.00 | 3.00 | 2.00 | 14.33 | 5.33  | 2.67  | 0.25 | 0.16 | 0.10 |
| 1MR 245 | 14.00 | 8.67 | 4.00 | 4.33 | 2.33 | 1.33 | 25.33 | 11.00 | 4.00  | 0.22 | 0.15 | 0.08 |
| 1MR 247 | 11.67 | 6.67 | 2.67 | 5.33 | 3.33 | 2.33 | 9.67  | 3.67  | 1.67  | 0.24 | 0.15 | 0.07 |
| 1MR 248 | 8.33  | 4.67 | 2.33 | 5.33 | 3.33 | 2.33 | 25.67 | 9.67  | 3.33  | 0.17 | 0.11 | 0.06 |
| 1MR 268 | 12.00 | 6.33 | 3.33 | 6.00 | 4.00 | 3.00 | 12.67 | 5.67  | 1.33  | 0.33 | 0.18 | 0.09 |
| 1MR 272 | 18.00 | 8.33 | 3.67 | 5.00 | 3.00 | 1.67 | 34.33 | 16.33 | 9.67  | 0.34 | 0.17 | 0.10 |
| 1MR 292 | 11.00 | 5.67 | 2.67 | 3.33 | 2.33 | 1.33 | 14.00 | 2.67  | 1.67  | 0.22 | 0.12 | 0.08 |

|         |       |       |      |      |      |      |       |       |       |      |      |      |
|---------|-------|-------|------|------|------|------|-------|-------|-------|------|------|------|
| 1MR 368 | 9.00  | 3.67  | 1.67 | 3.67 | 2.33 | 1.33 | 17.33 | 7.33  | 2.33  | 0.50 | 0.28 | 0.12 |
| 1MR 376 | 15.33 | 6.33  | 2.33 | 3.67 | 2.33 | 1.33 | 25.33 | 11.33 | 2.67  | 0.46 | 0.28 | 0.14 |
| 1MR 377 | 15.00 | 6.67  | 2.67 | 7.33 | 4.33 | 2.33 | 48.67 | 18.67 | 5.33  | 0.31 | 0.19 | 0.08 |
| 1MR 379 | 15.67 | 8.33  | 2.67 | 6.67 | 4.00 | 2.00 | 27.00 | 10.00 | 4.00  | 0.20 | 0.10 | 0.04 |
| 1MR 384 | 11.33 | 4.67  | 1.67 | 5.67 | 3.67 | 2.67 | 59.33 | 27.33 | 14.00 | 0.62 | 0.36 | 0.16 |
| 1MR 385 | 15.00 | 5.00  | 2.00 | 8.67 | 6.33 | 3.33 | 36.00 | 22.00 | 11.00 | 0.22 | 0.16 | 0.10 |
| 1MR 386 | 17.00 | 9.67  | 3.67 | 5.00 | 3.00 | 2.00 | 11.67 | 4.67  | 1.33  | 0.24 | 0.13 | 0.06 |
| 1MR 387 | 18.00 | 7.67  | 2.67 | 6.00 | 3.67 | 2.67 | 16.67 | 6.67  | 2.67  | 0.42 | 0.21 | 0.10 |
| 1MR 413 | 11.00 | 5.67  | 3.67 | 4.33 | 2.33 | 1.33 | 20.33 | 9.33  | 3.67  | 0.25 | 0.11 | 0.07 |
| 1MR 414 | 15.33 | 9.00  | 4.00 | 5.33 | 3.33 | 2.33 | 11.33 | 4.33  | 2.67  | 0.53 | 0.31 | 0.16 |
| 1MR 415 | 17.33 | 9.00  | 3.67 | 4.67 | 2.67 | 1.67 | 12.67 | 6.67  | 2.67  | 0.25 | 0.13 | 0.06 |
| 1MR 416 | 19.00 | 10.00 | 4.00 | 4.67 | 2.67 | 1.67 | 40.67 | 18.67 | 8.67  | 0.55 | 0.29 | 0.13 |
| 1MR 417 | 17.00 | 9.00  | 3.67 | 5.33 | 3.33 | 1.33 | 34.33 | 13.67 | 9.33  | 0.47 | 0.25 | 0.07 |
| 1MR 419 | 11.33 | 7.00  | 2.67 | 4.33 | 2.33 | 1.33 | 48.33 | 18.33 | 8.33  | 0.39 | 0.22 | 0.06 |
| 1MR 420 | 18.00 | 9.33  | 3.67 | 6.33 | 4.33 | 2.33 | 15.33 | 5.33  | 2.33  | 0.45 | 0.23 | 0.08 |
| 1MR 424 | 15.00 | 7.33  | 3.33 | 6.67 | 3.67 | 1.67 | 42.67 | 17.67 | 12.67 | 0.60 | 0.34 | 0.13 |
| 1MR 425 | 17.00 | 8.00  | 3.33 | 7.00 | 5.00 | 2.33 | 24.00 | 14.00 | 7.00  | 0.50 | 0.37 | 0.24 |
| 1MR 429 | 11.00 | 5.67  | 2.33 | 4.67 | 2.67 | 1.67 | 43.00 | 27.00 | 13.00 | 0.46 | 0.39 | 0.23 |
| 1MR 439 | 11.00 | 6.00  | 2.00 | 8.00 | 5.00 | 4.00 | 31.00 | 20.00 | 11.00 | 0.22 | 0.16 | 0.11 |
| 1MR 445 | 9.67  | 4.67  | 1.33 | 4.67 | 3.33 | 1.33 | 46.00 | 32.00 | 15.00 | 0.24 | 0.19 | 0.11 |
| 1MR 447 | 8.67  | 4.33  | 1.67 | 5.33 | 3.33 | 1.33 | 34.00 | 22.00 | 10.00 | 0.52 | 0.46 | 0.33 |
| 1MR 450 | 9.00  | 4.67  | 2.33 | 7.33 | 5.33 | 2.33 | 24.00 | 13.00 | 6.00  | 0.36 | 0.29 | 0.18 |
| 1MR 451 | 12.00 | 6.67  | 2.33 | 5.33 | 4.00 | 2.33 | 14.33 | 9.00  | 4.00  | 0.31 | 0.25 | 0.14 |
| 1MR 526 | 13.00 | 7.00  | 2.67 | 6.33 | 5.33 | 2.33 | 12.67 | 7.00  | 2.00  | 0.26 | 0.19 | 0.10 |
| 1MR 534 | 10.67 | 7.00  | 3.00 | 5.33 | 3.67 | 1.33 | 16.33 | 9.00  | 3.00  | 0.23 | 0.17 | 0.07 |
| 1MR 540 | 13.00 | 8.00  | 3.33 | 5.67 | 3.67 | 1.67 | 19.67 | 10.00 | 5.00  | 0.37 | 0.30 | 0.16 |
| 1MR 541 | 8.00  | 4.33  | 1.33 | 6.67 | 5.00 | 2.33 | 27.00 | 14.00 | 7.00  | 0.34 | 0.29 | 0.17 |
| 1MR 564 | 10.00 | 4.67  | 1.67 | 6.33 | 4.33 | 2.33 | 17.00 | 8.00  | 3.00  | 0.36 | 0.24 | 0.15 |
| 1MR 565 | 11.67 | 6.00  | 2.33 | 8.67 | 7.00 | 3.33 | 63.00 | 42.00 | 24.00 | 0.33 | 0.26 | 0.12 |
| 1MR 566 | 18.33 | 11.00 | 5.00 | 4.67 | 3.33 | 1.33 | 38.00 | 23.00 | 12.00 | 0.23 | 0.17 | 0.09 |

|                     |   |             |             |  |      |      |  |              |             |  |      |      |
|---------------------|---|-------------|-------------|--|------|------|--|--------------|-------------|--|------|------|
| 1MR 569             | 10.33   | 6.00        | 2.67        | 8.00   | 6.00 | 3.00 | 40.00  | 28.00        | 14.00       | 0.49   | 0.41 | 0.29 |
| 1MR 571             | 9.33  | 5.00        | 2.00        | 5.33   | 3.33 | 1.67 | 27.00  | 18.00        | 10.00       | 0.47   | 0.42 | 0.30 |
| 1MR 572             | 17.00   | 10.00       | 3.67        | 4.00   | 2.67 | 1.00 | 34.00  | 21.00        | 8.00        | 0.31   | 0.27 | 0.14 |
| 1MR 573             | 15.00   | 8.00        | 2.67        | 5.33   | 3.33 | 1.50 | 20.00  | 11.00        | 4.00        | 0.20   | 0.13 | 0.07 |
| 1MR 574             | 15.67   | 9.00        | 3.67        | 7.67   | 5.67 | 1.67 | 21.00  | 13.00        | 7.00        | 0.61   | 0.56 | 0.43 |
| 1MR 576             | 11.00   | 6.00        | 2.33        | 7.67   | 6.33 | 3.33 | 15.00  | 9.00         | 3.00        | 0.23   | 0.18 | 0.06 |
| 1MR 577             | 18.33   | 10.00       | 3.67        | 5.33   | 3.33 | 1.67 | 23.00  | 12.00        | 6.00        | 0.43   | 0.37 | 0.25 |
| 1MR 579             | 18.00   | 11.00       | 4.33        | 4.67   | 2.33 | 1.33 | 19.00  | 10.00        | 3.00        | 0.26   | 0.20 | 0.11 |
| 1MR 580             | 12.00   | 7.33        | 3.33        | 8.00   | 6.00 | 3.00 | 15.00  | 9.00         | 5.00        | 0.28   | 0.19 | 0.11 |
| 1MR 582             | 11.67   | 7.00        | 2.33        | 5.00   | 2.67 | 1.67 | 25.00  | 14.00        | 7.00        | 0.52   | 0.47 | 0.34 |
| 1MR 583             | 15.00   | 8.00        | 3.67        | 6.33   | 4.33 | 2.67 | 14.00  | 11.00        | 6.00        | 0.31   | 0.26 | 0.23 |
| 1MR 585             | 18.00   | 8.67        | 2.67        | 4.67   | 3.33 | 1.33 | 35.00  | 24.00        | 14.00       | 0.56   | 0.48 | 0.36 |
| 1MR 609             | 17.00   | 10.00       | 3.67        | 8.00   | 6.00 | 3.00 | 26.00  | 13.00        | 8.00        | 0.47   | 0.41 | 0.33 |
| 1MR 667             | 14.00   | 9.00        | 3.33        | 9.00   | 7.00 | 4.00 | 27.00  | 11.00        | 6.00        | 0.39   | 0.32 | 0.25 |
| 1MR 668             | 11.00   | 7.00        | 2.00        | 5.33   | 3.33 | 1.33 | 32.00  | 21.00        | 13.00       | 0.47   | 0.39 | 0.33 |
| 1MR 671             | 17.33   | 12.00       | 5.00        | 6.33   | 4.33 | 2.33 | 48.00  | 29.00        | 15.00       | 0.59   | 0.53 | 0.44 |
| 1MR A               | 16.00   | 13.00       | 5.67        | 8.33   | 6.33 | 3.33 | 26.33  | 18.00        | 7.00        | 0.24   | 0.18 | 0.09 |
| 1MR B               | 18.00   | 11.00       | 4.33        | 5.00   | 3.00 | 1.33 | 58.00  | 37.00        | 18.00       | 0.51   | 0.45 | 0.35 |
| 1MR C               | 12.00   | 7.00        | 2.33        | 7.67   | 5.67 | 2.67 | 21.00  | 12.00        | 6.00        | 0.30   | 0.24 | 0.17 |
| <b>Mean</b>         | <b>13.42</b>  | <b>7.16</b> | <b>2.92</b> | <b>5.86</b>  | 3.92 | 2.05 | <b>29.77</b>   | <b>16.64</b> | <b>8.04</b> | <b>0.37</b>  | 0.28 | 0.17 |
| <b>C.D(p ≤0.05)</b> | Genotype =1.217<br>PEG Levels =0.207<br>Genotype × PEG Levels = 2.108 |             |             | Genotype = 0.875<br>PEG Levels =0.149<br>Genotype × PEG Levels = 1.515 |      |      | Genotype = 1.203<br>PEG Levels =0.205<br>Genotype × PEG Levels = 2.084 |              |             | Genotype = 0.014<br>PEG Levels =0.002<br>Genotype × PEG Levels = 0.025 |      |      |

**Table-2: Analysis of variance for traits scored under different levels of PEG 6000 in maize (*Zea mays* L.) inbreds**

| Source of variation    | d.f | Primary root length (cm) | Number of Seminals | Number of Laterals | Root Biomass (g) |
|------------------------|-----|--------------------------|--------------------|--------------------|------------------|
| Genotypes              | 102 | 37**                     | 11**               | 677**              | 0.119**          |
| PEG Levels             | 2   | 8615**                   | 1126.6**           | 37041**            | 3.286**          |
| Genotypes × PEG Levels | 204 | 6**                      | 0.8**              | 66**               | 0.005**          |
| Error                  | 618 | 2                        | 0.9                | 2                  | 0                |

\*\* Significant at 5% level

## References

- Abdel-Raheem, A. T., Ragab, A. R., Kasem, Z. A., Omar, F. D. and Samera, A. M. 2007. In vitro selection for tomato plants for drought tolerance via callus culture under polyethylene glycol (PEG) and mannitol treatments. *African Crop Science Conference Proceedings*8(8): 2027-2032.
- Ahmad, Z., Waraich, E. A., Ahmad, R., Iqbal, M. A. and Awan, M. I. 2015. Studies on screening of maize (*Zeamays* L.) hybrids under drought stress conditions. *Journal of Advanced Botany and Zoology*2: 1-5.
- Alvi, A. K., Ahmad, M. S. A., Rafique, T., Naseer, M., Farhat, F., Tasleem, H. and Nasim, A. 2022. Screening of maize (*Zeamays* L.) genotypes for drought tolerance using

- photosynthetic pigments and antioxidative enzymes as selection criteria. *Pakistan Journal of Botany***54**(1): 33-44.
- Badr, A., Hana H.S., Rasha A.T and Borner , A. 2020. Screening for drought tolerance in Maize (*Zea mays* L.) germplasm using germination and seedling traits under simulated drought conditions. *Plants*, 9: 565
- Bukhari, B., Sabaruddin, Z.,Sufardi, S. and Syafruddin, S. 2021. Drought test resistance of maize varieties through PEG 6000. In *IOP Conference Series: Earth and Environmental Science* (Vol. 644, No. 1, p. 012040). IOP Publishing.
- Dani, A. R. H. and Siswoyo, T. A. 2019. Impact of drought stress during germination on antioxidant capacities and antioxidant enzymes activities of Madura local maize (*Zeamays* L.) seeds. *Agricultural Sciences***10**(11): 1506-1516.
- Fatima, Z., Awais, M., Waheed, A., Ali, A., Hashmi, Q. A., Abbas, M. and Sufyan, A. 2023. Effect of Polyethylene Glycol (PEG) on Germination and Seedling Parameters of Maize Varieties. *Asian Journal of Research in Crop Science*, 8(4), 410-420.
- Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F. and Toulmin, C. (2010). Food security: the challenge of feeding 9 billion people. *Science***327**: 812-818.
- Hellal, F. A., H. M. El-Shabrawi, M. Abd El-Hady, I. A. Khatab, S. A. A. El-Sayed, and C. Abdelly. 2018. Influence of PEG induced drought stress on molecular and biochemical constituents and seedling growth of Egyptian barley cultivars. *Journal of Genetics Engineering and Biotechnology***16**(1): 203-212.
- Jatoi, S. A., Latif, M. M., Arif, M., Ahson, M., Khan, A. and Siddiqui, S. U. 2014. Comparative assessment of wheat landraces against polyethylene glycol simulated drought stress. *Science Technology and Development* **33**: 1-6.


- Khodarahmpour, Z. 2011. Effect of drought stress induced by polyethylene glycol (PEG) on germination indices in corn (*Zea mays* L.) hybrids. *African Journal of Biotechnology* **10**(79): 18222-18227.
- Kyenpia, E. O., Namo, O. A. T., Gikyuu, S. W. and Ifenkwe, O. P. 2009. A comparative study of the biochemical composition of some varieties of maize (*Zea mays* L.) grown in Nigeria. *Nigeria Journal of Botany* **22**: 291-296.
- Magar, M. M., Parajuli, A., Shrestha, J., Koirala, K. B. and Dhital, S. P. 2019. Effect of PEG induced drought stress on germination and seedling traits of maize (*Zea mays* L.) lines. *Türk TarımveDoğaBilimleriDergisi***6**(2): 196-205.
- Majid, A., Dar, Z. A., Zaffar, G., Lone, F. A., Kumar, I. S., Sofi, P. A. and Rashid, M. 2020. Effect of PEG-6000 induced drought stress on seed germination in Maize (*Zea mays*). *SKUAST Journal of Research*, **22**(1), 40-44.
- Mustamu, N. E., Tampubolon, K., Basyuni, M., Al-Taey, D. K., Janabi, H. J. K. A. and Mehdizadeh, M. 2023. Drought stress induced by polyethylene glycol (PEG) in local maize at the early seedling stage. *Heliyon***9**(9): 213-231.
- Nivethitha, T., Ravikesavan, R., KumariVinodhana, N.,and Senthil, N. 2020. Deciphering drought tolerance potential of sweet corn genotypes through polyethylene glycol induced drought stress. *Electronic Journal of Plant Breeding***11**(01), 217-223.
- Petcu, E., Martura, T., Ciocăzanu, I., Iordan, H. L., Băduț, C. and Urechean, V. 2018. The Effect of water stress induced with PEG solution on maize seedlings. *Romanian Agricultural Research* **35**: 21-28.
- Prasanna, B. M. 2012. Diversity in global maize germplasm: characterization and utilization. *Journal of Biosciences***37**(5), 843-855.
- Salvi, S., Sponza, G., Morgante, M., Tomes, D., Niu, X., Fengler, K. A., Meeley, R., Ananiev, E. V., Svitashv, S., Bruggemann, E. and Li, B. 2007. Conserved noncoding genomic

sequences associated with a flowering-time quantitative trait locus in maize. *Proceedings of the National Academy of Sciences* **104**: 11376-11381.

Suhartono, A. A. 2021. Selection of maize plants resistant to drought stress in the vegetative phase using polyethylene glycol (PEG 6000). *AMA Agric Mech Asia Afr Lat Am*, 52(1), 2255-2261.

Tahir, S., Zafar, S., Ashraf, M. Y., Perveen, S. and Mahmood, S. 2023. Evaluation of drought tolerance in maize (*Zea mays L.*) using physiological indices. *Pakistan Journal of Botany* **55**(3): 843-849.

Zafar, S., Yasin Ashraf, M., Niaz, M., Kausar, A. and Hussain. J. 2015. Evaluation of wheat genotypes for salinity tolerance using physiological indices as a screening tool. *Pakistan Journal of Botany* **47**(2): 397-405.



**Fig1 - Roots of maize(*Zea mays*) inbreds under different PEG concentrations**