

Study on spatial distribution and occurrence of pink boll worm on cotton

ABSTRACT:

An investigation was carried out to “Study on spatial distribution and occurrence of pink boll worm on cotton” under Agricultural Research Station (ARS), Adilabad, Professor Jayashankar Telangana State Agricultural University during Kharif, 2018 and 2019. The main objective of this study was to determine distance and migration of pink boll worm (PBW) adults from ginning mill to cropping area of different location through geographic information system (GIS) tools and the results revealed that spatial distribution of pink bollworm was found to be low in the villages which are 15 kms away from the ginning mill i.e. pheromone catches (1515 and 1267) and PBW larvae/20 bolls (13.76 % and 15.00%) and villages which are 5kms away from the ginning mill was found to be high incidence i.e. pheromone catches (2185 and 1929) and PBW larvae/20 bolls (38.14 % and 28.10%) during Kharif 2018 and 2019, respectively.

KEY WORDS: Kharif, pink boll worm, spatial distribution, ginning mills, migration, GIS tools, pheromone catches, PBW larvae.

INTRODUCTION:

Cotton is one of the most commercially important fiber crop in the world, cultivated in tropical and subtropical regions of more than 70 countries. In India, cotton is mainly cultivated in three distinct agro-ecological zones North, Central and South. It is an important raw material for the Indian textile industry and plays a key role in the national economy in terms of both employment generation and foreign exchange. In India, cotton is cultivated in 119.78 lakh hectare with a production of 365 lakh bale of seed cotton (Sathyan *et al.*, 2016).

The cotton species recognized in the world are about 50 of which 4 are cultivated. Two of these (*Gossypium arboreum* and *G. herbaceum*) are diploids, and two (*G. hirsutum* and *G. barbadense*) are tetraploids. More than 80% of the world's cotton area is covered by tetraploids. However, diploid cottons are cultivated in Asia and the Middle East. India is the only country where all the cultivated species and some of their hybrid combinations are commercially grown (khadi *et al.*, 2015).

Insect pest attack is one of the most important limiting factors in the successful cultivation of this crop. Amongst various causes of low yield, losses due to insect pests are

Formatted: Font: Not Italic

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

one of the important factors in cotton. Many insect pests are encountered on cotton crop from germination to harvesting. Among the important key pests of cotton the sucking pests viz., leafhopper, *Amrasca Amrasca devastans* (Distant), aphid *Aphis gossypii* (Glover), whitefly *Bemisia tabaci* (Gennadius) and thrips, *Thrips tabaci* (Linnman) cause severe damage and serious threat to the crop at early stage of the crop growth and can also affect the crop stand and yield of cotton.

Comment [Dr1]: Check this

Formatted: Highlight

Pink bollworm has become apparent as a threat to cotton cultivation in south and central cotton growing zones of India where the pest has developed resistance to Cry1Ac and Cry2Ab expressing cotton also developing resistance to insecticides and infesting late season cotton. The pink bollworm is assuming a major pest status even in some regions of northern India where there are ginning and oil extraction units which are procuring cotton seeds from central and south Indian cotton states where pink bollworm has demonstrated resistance to Cry toxins in the field. So there is possibility of resistant pink bollworm infesting Bt cotton in North India through movement of seed. The evolution of resistance and pest adaptation to Bt crops containing Cry1Ac and Cry2Ab has been observed recently. The development of resistance is due to multiple factors such as absence of refuge or supply of fraudulent refuge, mono cropping, cultivation of long duration hybrids, extended cropping season (Naik *et al.*, 2020).

MATERIALS AND METHODS:

The present experiment was carried out at Agriculture research station, Adilabad during Kharif, 2018 and 2019, in order to study on spatial distribution and occurrence of pink boll worm on cotton based on distance from ginning mills and to find out the effect of distance on spatial distribution of pink boll worm on cotton crop through GIS tools. Data has been taken at seven days interval from boll formation stage from three different locations *i.e.*, 5km, 10km, and 15km from ginning mills and selected five villages from each location.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

Comment [Dr2]: Give a space between numerical value and unit of measurement

Monitoring of pink bollworm population using sex pheromone traps: The adult male moth population of main lepidopterous pest, pink bollworm of cotton crop is being monitored with installation of sex pheromone traps at selected villages and at ginning mill and also the larvae of *P. gossypiella* were collected from green bolls.

RESULTS AND DISCUSSION:

The observations recorded on population dynamics of pink boll worm of cotton at eight different dates *i.e.*, 15/11/18, 22/11/18, 29/11/18, 05/12/18, 12/12/18, 19/12/18,

26/12/18, 02/01/19 during kharif, 2018 and 20/11/19, 27/11/19, 04/12/19, 11/12/19, 18/12/19, 24/12/19, 31/12/19, 08/01/20 during kharif, 2019 are presented in Tables 1- 6, respectively.

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Spatial distribution and occurrence of pink bollworm in Adilabad district by using GIS tools (5km) during kharif, 2018 and 2019

Formatted: Font: Not Italic

The incidence of pink bollworm at boll formation stages of crop growth was observed in terms of pheromone trap catches and pink bollworm larvae per 20 bolls.

During kharif, 2018 high incidence of pink bollworm in pheromone trap catches and pink bollworm larvae per 20 bolls in five villages at an distance of 5km from ginning mills i.e., ponnari (420 and 5.63%), pochera (427 and 9.38%), anukunta (436 and 7.50%), piperwada (441 and 7.50%) and kajjarla (460 and 8.13%) pheromone traps catches and pink bollworm larvae, respectively.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, English (United States)

During kharif, 2019 incidence of pink bollworm in pheromone trap catches and pink bollworm larvae per 20 bolls was high in five villages at an distance of 5km from ginning mills i.e., ponnari (390 and 7.50%), pochera (367 and 6.25%), anukunta (384 and 4.37%), piperwada (387 and 4.37%) and kajjarla (401 and 5.62%) Pheromone traps catches and PBW larvae, respectively.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, English (United States)

Spatial distribution and occurrence of pink bollworm in Adilabad district by using GIS tools (10km) during kharif, 2018 and 2019

Formatted: Font: Not Italic

The incidence of pink bollworm at boll formation stages of crop growth was observed in terms of Pheromone trap catches and larvae per 20 bolls.

During kharif, 2018 high incidence of pink bollworm in pheromone trap catches and pink bollworm larvae per 20 bolls in five villages at an distance of 10km from ginning mills i.e., sunkidi (352 and 6.25%), seethagondi (332 and 4.38%), Bhoraj (379 and 5.63%), mediguda (391 and 8.13%) and yalaguda (407 and 3.75%) pheromone traps catches and pink bollworm larvae, respectively.

Comment [Dr3]: Check this

Comment [Dr4]: Check this

Formatted: Font: (Default) Times New Roman, 12 pt, English (United States),

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, English (United States)

During kharif, 2019 high incidence of pink bollworm in Pheromone trap catches and PBW larvae per 20 bolls in five villages at an distance of 10km from ginning mills i.e., sunkidi (297 and 5.00%), seethagondi (308 and 4.37%), Bhoraj (321 and 4.37%), mediguda (326 and 2.50%) and yalaguda (340 and 3.12%) Pheromone traps catches and PBW larvae, respectively.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, English (United States)

Spatial distribution and occurrence of pink bollworm in Adilabad district by using GIS tools (15km) during kharif, 2018 and 2019:

Formatted: Font: Not Italic

The incidence of pink bollworm at boll formation stages of crop growth was

observed in terms of pheromone trap catches and pink bollworm larvae per 20 bolls.

During kharif, 2018 high incidence of pink bollworm in pheromone trap catches and PBW larvae per 20 bolls in five villages at an distance of 15km from ginning mills i.e., jainath(270 and 6.25%), niral (283 and 4.38%), talamadugu (303 and 1.25%), gudihatnoor (326 and 0.63%) and tantoli (333 and 1.25%) Pheromone traps catches and PBW larvae, respectively.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, English (United States)

During kharif, 2019 high incidence of pink bollworm in Pheromone trap catches and PBW larvae per 20 bolls in five villages at an distance of 15km from ginning mills i.e., jainath (237 and 5.00%), niral (242 and 3.75%), talamadugu (256 and 2.50%), gudihatnoor (261 and 1.25%) and tantoli (271 and 2.50%) Pheromone traps catches and PBW larvae, respectively.

Comment [Dr5]: Only scientific names should be italicized.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, English (United States)

From the above study during kharif, 2018-19 spatial distribution of pink bollworm was found to be low in the villages which are 15 kms away from the ginning mill i.e. pheromone catches were 1515 and pink bollworm larvae/20 bolls 13.76 %, villages which are 10kms away from the ginning mill was found to be average incidence i.e. pheromone catches were 1861 and pink bollworm larvae/20 bolls 28.14 % and villages which are 5kms away from the ginning mill was found to be high incidence i.e. pheromone catches were 2185 and pink bollworm larvae/20 bolls 38.14 %.

Formatted: Font: Not Italic

During 2019-20, from the above study spatial distribution of pink bollworm was found to be low in the villages which are 15kms away from the ginning mill i.e. pheromone catches were 1267 and pink bollworm larvae/20 bolls 15.00 %, villages which are 10kms away from the ginning mill was found to be average incidence i.e. pheromone catches were 1592 and pink bollworm larvae/20 bolls 19.30 % and villages which are 5kms away from the ginning mill was found to be high incidence i.e. pheromone catches were 1929 and pink bollworm larvae/20 bolls 28.11 %.

Discussion

The flight potential and propensity of *Pectinophoragossypiella*(Saunders) in the laboratory were measured using a 32-channel, computer-monitored flight-mill system. Females flew significantly farther within a 72-hour flight period than males did. According to Wu et al. (2006), the mean accumulated flight distance and flight duration of 1-day-old female individuals in a tethered-flight test were 41.25 ± 7.76 km and 23.87 ± 2.55 hours, respectively, whereas for male individuals, the corresponding parameters were 23.46 ± 2.13 km and 14.12 ± 1.12 hours. The flight ability of adults was significantly positively correlated with pupal weight.

Comment [Dr6]: All scientific name should be italicized

Formatted: Font: Italic

Milonas *et al.*, (2016) reported that in Central Greece, spatial and temporal variations in population dynamics were examined. In 2007, captures of male *H. armigera* increased in late July and declined at the end of August. For *P. gossypiella* male captures remained at low levels during summer, increased late in August, peaked at mid of September and declined towards the end of the season. Spatial analysis produced a spatial trend map over space, a temporal stability map over time which could lead in separating the field into management zones and direct control to areas that exhibit high densities of the pest population and are stable over time.

Conclusion

We have found during the Kharif seasons of 2018 and 2019, villages situated 5 km away from the ginning mills exhibited high incidence rates of pink bollworm, as evidenced by the higher numbers of pheromone trap catches and pink bollworm larvae per 20 bolls. In contrast, villages located 15 km away showed lower incidence rates. The findings of this study contribute valuable insights for cotton growers and pest management authorities to develop more effective strategies for managing pink bollworm infestations and safeguarding cotton yields.




Fig 1. Map showing study location

Table 1: Spatial distribution and occurrence of Pink bollworm in Adilabad district by using GIS tools (5km) during kharif, 2018

Formatted: Font: Not Italic

Ponnari	Pheromone catches	PBW larvae/ 20bolls
15/11/18	48	1
22/11/18	50	1
29/11/18	52	1
05/12/18	43	1
12/12/18	52	1
19/12/18	52	1
26/12/18	59	2
02/01/19	64	1
TOTAL	420	9
PER CENT	-	5.63
Pochera	Pheromone catches	PBW larvae/ 20bolls
15/11/18	52	2
22/11/18	51	2
29/11/18	48	3
05/12/18	39	2
12/12/18	54	1
19/12/18	56	1
26/12/18	62	2
02/01/19	65	2
TOTAL	427	15
PER CENT	-	9.38
Anukunta	Pheromone catches	PBW larvae/ 20bolls
15/11/18	49	0
22/11/18	52	1
29/11/18	50	1
05/12/18	42	2
12/12/18	55	3
19/12/18	61	3
26/12/18	58	2
02/01/19	69	0
TOTAL	436	12
PER CENT	-	7.50
Piperwada	Pheromone catches	PBW larvae/ 20bolls
15/11/18	51	0
22/11/18	53	1
29/11/18	53	1
05/12/18	49	2
12/12/18	58	3
19/12/18	63	3
26/12/18	55	2
02/01/19	59	0
TOTAL	441	12
PER CENT	-	7.50
Kajjarla	Pheromone catches	PBW larvae/ 20bolls

15/11/18	52	1
22/11/18	54	3
29/11/18	55	3
05/12/18	52	1
12/12/18	56	1
19/12/18	64	2
26/12/18	60	1
02/01/19	67	1
TOTAL	460	13
PER CENT	-	8.13

Table 2: Spatial distribution and occurrence of Pink bollworm in Adilabad district by using GIS tools (10km) during kharif, 2018

Formatted: Font: Not Italic

Sunkidi	Pheromone catches	PBW larvae/ 20bolls
15/11/18	36	3
22/11/18	39	2
29/11/18	40	2
05/12/18	42	0
12/12/18	46	0
19/12/18	47	0
26/12/18	50	1
02/01/19	52	2
TOTAL	352	10
PER CENT	-	6.25
Seethagondi	Pheromone catches	PBW larvae/ 20bolls
15/11/18	39	1
22/11/18	40	4
29/11/18	39	0
05/12/18	39	0
12/12/18	39	0
19/12/18	43	0
26/12/18	44	1
02/01/19	49	1
TOTAL	332	7
PER CENT	-	4.38
Bhoraj	Pheromone catches	PBW larvae/ 20bolls
15/11/18	41	0
22/11/18	42	1
29/11/18	46	1
05/12/18	48	1
12/12/18	49	1
19/12/18	50	2
26/12/18	51	2
02/01/19	52	1
TOTAL	379	9
PER CENT	-	5.63

Mediguda	Pheromone catches	PBW larvae/ 20bolls
15/11/18	42	2
22/11/18	39	3
29/11/18	47	2
05/12/18	52	2
12/12/18	53	2
19/12/18	52	1
26/12/18	53	1
02/01/19	53	0
TOTAL	391	13
PER CENT	-	8.13
Yapalguda	Pheromone catches	PBW larvae/ 20bolls
15/11/18	43	1
22/11/18	46	2
29/11/18	48	1
05/12/18	50	0
12/12/18	50	1
19/12/18	53	1
26/12/18	58	0
02/01/19	59	0
TOTAL	407	6
PER CENT	-	3.75

Table 3: Spatial distribution and occurrence of Pink bollworm in Adilabad district by using GIS tools (15km) during kharif, 2018

Formatted: Font: Not Italic

Jainath	Pheromone catches	PBW larvae/ 20bolls
15/11/18	25	1
22/11/18	28	1
29/11/18	30	0
05/12/18	32	1
12/12/18	35	0
19/12/18	39	1
26/12/18	40	0
02/01/19	41	2
TOTAL	270	6
PER CENT	-	6.25
Nirala	Pheromone catches	PBW larvae/ 20bolls
15/11/18	28	2
22/11/18	30	0
29/11/18	32	1
05/12/18	33	1
12/12/18	36	0
19/12/18	38	1
26/12/18	42	1
02/01/19	44	1

TOTAL	283	7
PER CENT	-	4.38
Talamadugu	Pheromone catches	PBW larvae/ 20bolls
15/11/18	29	1
22/11/18	34	1
29/11/18	35	0
05/12/18	39	0
12/12/18	38	0
19/12/18	40	0
26/12/18	43	0
02/01/19	45	0
TOTAL	303	2
PER CENT	-	1.25
Gudihatnoor	Pheromone catches	PBW larvae/ 20bolls
15/11/18	30	0
22/11/18	36	1
29/11/18	39	0
05/12/18	42	0
12/12/18	45	0
19/12/18	42	0
26/12/18	45	0
02/01/19	47	0
TOTAL	326	1
PER CENT	-	0.63
Tantoli	Pheromone catches	PBW larvae/ 20bolls
15/11/18	32	1
22/11/18	39	1
29/11/18	40	0
05/12/18	41	0
12/12/18	43	0
19/12/18	44	0
26/12/18	46	0
02/01/19	48	0
TOTAL	333	2
PER CENT	-	1.25

Table 4: Spatial distribution and occurrence of Pink bollworm in Adilabad district by using GIS tools (5km) during kharif, 2019

Formatted: Font: Not Italic

Ponnari	Pheromone catches	PBW larvae/ 20bolls
20/11/19	18	1
27/11/19	32	1
04/12/19	41	1
11/12/19	48	1
18/12/19	58	1
24/12/19	60	2
31/12/19	64	3
08/01/20	69	2

TOTAL	390	12
PER CENT	-	7.50
Pochera	Pheromone catches	PBW larvae/ 20bolls
20/11/19	15	0
27/11/19	26	1
04/12/19	44	1
11/12/19	52	2
18/12/19	54	1
24/12/19	56	1
31/12/19	58	2
08/01/20	62	2
TOTAL	367	10
PER CENT	-	6.25
Anukunta	Pheromone catches	PBW larvae/ 20bolls
20/11/19	19	0
27/11/19	30	0
04/12/19	49	2
11/12/19	50	1
18/12/19	54	2
24/12/19	58	0
31/12/19	60	1
08/01/20	64	1
TOTAL	384	7
PER CENT	-	4.37
Piperwada	Pheromone catches	PBW larvae/ 20bolls
20/11/19	14	1
27/11/19	28	1
04/12/19	40	1
11/12/19	53	1
18/12/19	58	1
24/12/19	61	1
31/12/19	65	0
08/01/20	68	1
TOTAL	387	7
PER CENT	-	4.37
Kajjarla	Pheromone catches	PBW larvae/ 20bolls
20/11/19	16	0
27/11/19	32	1
04/12/19	45	1
11/12/19	51	2
18/12/19	59	2
24/12/19	62	1
31/12/19	67	1
08/01/20	69	1
TOTAL	401	9
PER CENT	-	5.62

Table 5: Spatial distribution and occurrence of Pink bollworm in Adilabad district by using GIS tools (10km) during kharif, 2019

Formatted: Font: Not Italic

Sunkidi	Pheromone catches	PBW larvae/ 20bolls
20/11/19	9	0
27/11/19	20	1
04/12/19	34	1
11/12/19	40	1
18/12/19	45	1
24/12/19	47	0
31/12/19	50	2
08/01/20	52	2
TOTAL	297	8
PER CENT	-	5.00
Seethagondi	Pheromone catches	PBW larvae/ 20bolls
20/11/19	11	1
27/11/19	18	0
04/12/19	30	1
11/12/19	41	0
18/12/19	48	2
24/12/19	50	1
31/12/19	52	1
08/01/20	58	1
TOTAL	308	7
PER CENT	-	4.37
Bhoraj	Pheromone catches	PBW larvae/ 20bolls
20/11/19	8	0
27/11/19	19	1
04/12/19	32	0
11/12/19	45	1
18/12/19	49	0
24/12/19	53	2
31/12/19	56	1
08/01/20	59	2
TOTAL	321	7
PER CENT	-	4.37
Mediguda	Pheromone catches	PBW larvae/ 20bolls
20/11/19	8	0
27/11/19	17	0
04/12/19	36	1
11/12/19	42	0
18/12/19	48	1
24/12/19	54	1
31/12/19	59	0
08/01/20	62	1
TOTAL	326	4
PER CENT	-	2.50
Yapalguda	Pheromone catches	PBW larvae/ 20bolls

20/11/19	7	0
27/11/19	21	0
04/12/19	38	1
11/12/19	50	1
18/12/19	46	1
24/12/19	55	1
31/12/19	56	1
08/01/20	67	0
TOTAL	340	5
PER CENT	-	3.12

Table 6: Spatial distribution and occurrence of Pink bollworm in Adilabad district by using GIS tools (15km) during kharif, 2019

Formatted: Font: Not Italic

Jainath	Pheromone catches	PBW larvae/ 20bolls
20/11/19	4	0
27/11/19	12	0
04/12/19	26	1
11/12/19	32	1
18/12/19	36	1
24/12/19	39	2
31/12/19	42	1
08/01/20	46	2
TOTAL	237	8
PER CENT	-	5.00
Nirala	Pheromone catches	PBW larvae/ 20bolls
20/11/19	5	0
27/11/19	14	0
04/12/19	21	0
11/12/19	30	1
18/12/19	38	1
24/12/19	41	1
31/12/19	45	2
08/01/20	48	1
TOTAL	242	6
PER CENT	-	3.75
Talamadugu	Pheromone catches	PBW larvae/ 20bolls
20/11/19	5	0
27/11/19	14	0
04/12/19	27	1
11/12/19	34	0
18/12/19	38	1
24/12/19	42	0
31/12/19	47	1
08/01/20	49	1
TOTAL	256	4

PER CENT	-	2.50
Gudihatnoor	Pheromone catches	PBW larvae/ 20bolls
20/11/19	7	0
27/11/19	10	0
04/12/19	23	0
11/12/19	36	1
18/12/19	40	0
24/12/19	46	0
31/12/19	49	1
08/01/20	50	0
TOTAL	261	2
PER CENT	-	1.25
Tantoli	Pheromone catches	PBW larvae/ 20bolls
20/11/19	10	0
27/11/19	18	1
04/12/19	27	0
11/12/19	39	1
18/12/19	35	0
24/12/19	43	1
31/12/19	48	0
08/01/20	51	1
TOTAL	271	4
PER CENT	-	2.50

Table 7: Spatial distribution and occurrence of Pink bollworm PBW in Adilabad district by using GIS tools (5, 10 and 15km)

Distances (2018-19)	Pheromone catches	PBW larvae in percent
5 km distance total (five villages)	2185	38.1
10 km distance total (five villages)	1861	28.1
15 km distance total (five villages)	1515	13.7

Distances (2019-20)	Pheromone catches	PBW larvae in percent
5 km distance average (five villages)	1929	28.1
10 km distance average (five villages)	1592	19.3
15 km distance average (five villages)	1267	15.0

Reference:

Chinna Babu Naik, V., Pratik, P., Pusadkar, Sandesh, T., Waghmare, Raghavendra K. P.,

- Sandhya Kranthi, Sujit Kumbhare, V. S.,Nagrare, Rishi Kumar, Tenguri Prabhulinga1, NandiniGokte-Narkhedkar&Waghmare, V. N. 2020. Evidence for population expansion of Cotton pink bollworm *Pectinophoragossypiella* (Saunders) (Lepidoptera: Gelechiidae) in India. *Scientific Reports*. 10:4740.
- Huaiheng, Wu., Kongming, Wu., Dengyuan Wang and Yuyuan Guo. 2006. Flight Potential of Pink Bollworm, *Pectinophoragossypiella* Saunders (Lepidoptera: Gelechiidae). *Environmental Entomology*. 35(4): 887-893.
- Khadi,B.M., Santhy,V. and Yadav, M.S. 2015. Cotton: An Introduction, Biotechnology in Agriculture and Forestry 65, DOI 10.1007/978-3-642-04796-1-1.
- Milonas,P., Gogou, C., Papadopoulou , A., Fountas, S.,Liakos, V., Papadopoulos,N. T. 2016. Spatio-Temporal Distribution of *Helicoverpaarmigera* (Hübner) (Lepidoptera: Noctuidae) and *Pectinophoragossypiella* (Saunders) (Lepidoptera: Gelechiidae) in a Cotton Production Area. *Neotropical Entomology*. 45, 240-251.
- Sathyan, T.,Murugesan, N., Elanchezhyan, K., Arockia Stephen Raj, J. and Ravi, G. 2016. Efficacy of Synthetic Insecticides against sucking insect pests in cotton, *Gossypiumhirsutum* L. *International Journal of Entomology Research*. 1(1): 16-21.