

Review Article

Potential of plant based metallophytes for phytoremediation in agriculture

Comment [NS1]: There is no action word in this title. What type of research are you carrying out? Is it a review study or what?

Abstract

Metallophytes are the unique group of plants that have evolved to thrive in metal rich environments and have drawn plenty of attention. Remediating heavy metal contaminated places with plants is an effective choice due to phytoremediation, an environmentally friendly method that uses plants to mitigate the pollution. Metallophytes are ideal options for phytoremediation applications due to their inherent traits such as hyper-accumulation, efficient metal absorption and tolerance mechanisms impacting both the plant and soil. These plants absorb and translocate heavy metals, detoxifying the soil while accumulating them in tissues. This reduces metal toxicity in soil and holds potential for resources recovery. The role of metallophytes in phytoremediation is analysed in this review with particular focus given to their ways of metal absorption, translocation and detoxification. Metallophytes have high metal tolerance and accumulation capacities due to their unique physiological and biochemical adaptations including enhanced metal sequestration in vacuoles, metal chelation by phytochelatins and activation of anti-oxidant defence systems. This review also highlights the significance of metallophytes in enhancing the soil health, reducing metal bioavailability, and promoting the ecological sustainability as well as their potential for restoring contaminated ecosystems. Utilizing the unique capabilities of metallophytes obtained from plants possesses enormous possibilities to minimise the negative effects of heavy metal pollution, protect ecosystems, and promote sustainable development for future generations. Eventually, it outlines future research approaches that aim to enhance metallophytes based phytoremediation strategies, widen their implementation and include them in holistic approaches for environmental restoration and sustainable land management.

Comment [NS2]: Please state the methodology employed in this study.

No results as well as recommendations. Revise

Comment [NS3]: No need for this space (paragraph)

Keywords: Metallophytes, phytoremediation, environmental restoration, heavy metals, soil health, ecosystem

Introduction

Due to massive economic development and quick expansion in numerous industries including agriculture, the environment is becoming more polluted (Shubham et al., 2023). Toxic metals which originate from natural and man-made sources might be referred to as environmental contaminants. Abiotic factors chiefly soil, water, air and biotic communities such as plants, animals and anthropogenic activities are put at risk by a number of environmental processes including burning trash, converting coal and creating synthetic businesses. A major issue preventing plants from growing in many situations is metal toxicity in soils. Certain metals like zinc and copper are considered to be micronutrients in low concentrations but become hazardous if present in larger concentrations, whereas, other metals like lead and aluminium are exclusively recognized for their toxicity. Certain soils naturally contain high traces of metals. For instance, serpentinite regions are found

contaminated with enormous amount of trace metals which includes high level of Cr, Ni and associated metals (Mg, Pb, Co and Zn) with other elements (Pal *et al.*, 2005). In addition, acid rains can cause metal poisoning indirectly by increasing the availability of most metals for plant absorption due to low soil pH. The Pb and Zn were mainly adsorbed on colloidal particles and were transported during simulated acid rain. The mobility of Zn was increased at low pH (Singh & Agrawal, 2007). Therefore, a significant constraint on agriculture may arise from the elevated availability of aluminium in some acidic soils.

Comment [NS4]: Write in full

The world is facing several challenges today due to rapid and reckless industrialization, intense farming practices, hazardous mining operations and waste disposal (Bhatti *et al.*, 2018; Shubham *et al.*, 2021). The health of people, animals and plants is also impacted by the continuous accumulation of metals in terrestrial systems. According to a Central Pollution Control Board (CPCB) research, 80 per cent of hazardous waste in India including heavy metals is produced in the states of Gujarat, Maharashtra and Andhra Pradesh (Marg, 2011).

The usage of pesticides, fertilisers, municipal and compost wastes, as well as, the emission of heavy metals from metalliferous mines and smelting businesses have all resulted in the contamination of large regions of land with heavy metals (Yang *et al.*, 2005). Even though the earth's crust contains a large number of heavy metals at different concentrations by nature, the issue comes when these metals are released into the environment in excess as a result of man-made or natural processes. Based on their density ($>5 \text{ g/cm}^3$), the 53 elements that make up the D-block have been classified as "heavy metals" (Jarup, 2003). Only 19 elements, including the macronutrients C, O, H, Mg, S, N, Ca, P, and K and the micronutrients Cu, Zn, Mn, Fe, Mo, B, Ni, Co, Cl, and B, were chosen for basic metabolism during the development of angiosperms (Ernst, 2006). Furthermore, silicon (Si) is regarded as a helpful element and has been implicated in certain plants structural preservation (Epstein, 1999). Plant physiological and biochemical activities, including chlorophyll production, photosynthesis, DNA synthesis, protein modifications, redox reactions in the chloroplast and mitochondrion, sugar metabolism and nitrogen fixation are significantly influenced by macro and micro-nutrients. In many plants, visual symptoms are caused by metal poisoning. Leaves may get discoloured and root development is frequently inhibited. Deficiencies in some nutrients (e.g., magnesium or calcium deficiency due to aluminium competition for root absorption) may appear as symptoms. Toxic metals can also be harmful when they attach to proteins and other substances because they share similarities with essential metals. Both essential and non-essential heavy metals often have similar harmful effects on plants including reduced biomass accumulation, chlorosis, growth and photosynthetic inhibition, altered water balance and nutrient absorption and senescence which eventually results in plant death. Since, heavy metals are so persistent in nature they not only have negative effects on plants but also constitute a hazard to human health. One of the most hazardous heavy metals is lead (Pb) which has around 150–5000 year soil retention period and has been shown to sustain high concentrations for up to 150 years (Yang *et al.*, 2005).

Comment [NS5]: Too old

According to McLaughlin *et al.* (1999) a contaminated food chain serves as the main pathway for the heavy metals to enter the tissues of animals and humans and thus increasing their susceptibility to a variety of illnesses including dermatitis and cancer. If appropriate action is not done at the appropriate time, this issue might get worse. Heavy metal toxicity in plants can occur in four different ways. These comprise: (a) Structural and functional similarities that cause competition for absorption at the root surface *e.g.* As and Cd compete with P and Zn, respectively (b) Direct interactions between heavy metals and the sulphhydryl

groups (-SH) of functional proteins which disrupt and degrade the structure and function of the proteins (c) Removal of necessary cations from specific binding sites that results in a collapse in function (d) Production of reactive oxygen species (ROS) which in turn damages the macromolecules. It has been well documented that plants may adapt to soils that are enriched in heavy metals (Antonovicset al., 1971). Plants that have modified their metabolism to enable growth and reproduction in metalliferous soils, such as *Agrostis capillaries* and *Silene vulgaris* have found success. It is possible for heavy metals like Cd, Pb, Al and Ag to be non-essential micronutrients in metal-enriched soils, yet necessary micronutrients like Mn, Zn, Cu, Fe and Mo may be present (Woolhouse, 1983). The soil characteristics at a certain location influence the kind and level of resistance to heavy metals. High atomic weight elements with a density at least five times that of water are known as heavy metals. These elements generally occur naturally. It is non-biodegradable and continues to accumulate in the environment, reaching toxic levels in the soil as a result of various natural processes like soil formation, rock weathering, volcanic eruptions as well as anthropogenic activities like excessive application of chemicals in agriculture, industry, mining and effluents from various sources etc. In the environment, heavy metals are quite persistent. Because of their slow movement and restricted breakdown once they are incorporated into the soil, they tend to accumulate over time. As a result, areas exposed to heavy metals over prolonged periods of time are becoming more contaminated, which presents long-term risks to crop production (Pehoiuet al., 2020). Such soils are remediated by various methods such as removal, isolation, incineration, solidification/stabilization, thermal treatment, solvent extraction and chemical oxidation. Currently, in situ techniques, phytoremediation techniques are preferred because they are more cost-effective and cause less environmental disruption (Favas et al., 2014). Using plants and associated soil microbes to reduce the harmful effects of pollutants (HM's) in the environment is known as phytoremediation. This technique uses metal accumulating plants referred as Metallophytes. Some metallophytes are Indian mustard, broad leaf plant like *ribwort plantain*, *Arabidopsis arenosa*(Singh et al., 2016).There are many pros of using metallophytes in phyto-remediation of heavy metal pollute soils as they are economically feasible, eco-friendly, prevents erosion and establishment of vegetation in barren lands.

Comment [NS6]: Outdated

Comment [NS7]: Outdated

Metallophytes

Metallophytes plants can be used in phytoremediation because of their capacity to absorb the significant concentrations of heavy metals from the soil. The capacity of metallophytes to withstand and even accumulate the dangerous concentrations of heavy metals in their tissues without suffering noticeably negative consequences is one of their most remarkable characteristics. There are different categories of metallophytes named metal excluders, metal hyper-accumulators and metal indicators (Devi & Kumar, 2020). Metal excluders are such metallophytes which accumulates heavy metals in the root from the particular contaminated soil, but the metals movement into the aerial plants components is restricted (Sheoranet al., 2011; Malik et al., 2012; Shubhamet al., 2023). According to Baker and Walker, (1990), plants that accumulate metals in their above ground tissues are known as metal indicators and the amounts of metals in these tissues typically corresponds to the levels of metals in the soil. Plant species classified as hyper-accumulators are capable of absorbing and utilizing large concentrations of heavy metals >100 to 10,000 µg/g (or much higher) relative to the plants dry weight without facing any harmful effects.

Comment [NS8]: Write in full

Metallophytes are classified into two types *i.e.* hyper-accumulators and non-hyper accumulators. Heavy metals from soils can be absorbed by hyper accumulators through their roots and then transferred to shoots and leaves. On the other hand, plant species classified as non-hyper accumulators are capable of storing heavy metals in their underground parts and are unable to transfer them to their shoots or leaves with a few species allowing restricted translocation. The heavy metals that these hyper accumulators take up from the soil are not stored in the roots instead they go up the shoots and build up in the above ground organs at concentrations that are 100–1000 times higher than those found in non-hyper accumulating species. Three characteristics mark them apart from non-hyper accumulators. These include increased heavy metal absorption capacity, root-to-shoot heavy metal translocation and detoxification and the storage of heavy metals (Singh *et al.*, 2016). Approximately, 45 per cent of hyper accumulators are members of angiosperm families of these, 25 per cent are members of the *Brassicaceae* family. Members of the *Asteraceae*, *Caryophyllaceae*, *Fabaceae*, *Cyperaceae*, *Poaceae*, *Cunoniaceae*, *Lamiaceae* and numerous other families are among the other families. Research indicates that the rhizosphere and endosphere of hyper-accumulator plants have a high diversity of bacteria and fungus (Kafle *et al.*, 2022; Ganiet *al.*, 2024). There is lots of research supporting *E. crassipes* ability to remove the heavy metals (Sarkar *et al.*, 2017).

Heavy metals toxicity in soil due to agronomic practices

- a) **Sources of contamination:** Two prominent ways that heavy metals might end up in agricultural soils are through chemical fertilisers and insecticides that contain the metals. Using contaminated irrigation water from industrial or municipal sources. The concentration of heavy metals (HM's) in agricultural soil grew in the following order, according to a study conducted in the agricultural soils of Jhansi, India: Cd, Co, Mn, Cu, Pb, Ni, and Zn. The Food and Agriculture Organisation/World Health Organisation (FAO/WHO), the United States Environmental Protection Agency (USEPA), and Indian norms are the research areas for concentrations of Cu, Zn, Pb, and Ni. The mean Cd content in the studied region was greater than the allowed limits established by the FAO/WHO and USEPA (Gupta *et al.*, 2021). Use of sewage sludge, also known as bio-solids, which may include significant levels of heavy metals from human waste and industrial discharge. Heavy metals are released into the atmosphere by industry and vehicle emissions.
- b) **Accumulation and persistence:** Because of their slow rates of degradation and restricted mobility, heavy metals have a tendency to accumulate in soil over time. Heavy metals may accumulate gradually and may contaminate crops after being added to the soil, where they can remain for extended periods of time.
- c) **Impacts on soil health:** High levels of heavy metals in soil can have detrimental effects on soil health through two different processes: disrupting soil microbial communities and nutrient cycle mechanisms. Lowering water retention and penetration as well as reducing the soil's structure. Decreasing the fertility of the soil and impeding plant growth.
- d) **Heavy metal absorption by plants:** Plants raised in polluted soils have the potential to absorb heavy metals through their roots, which can result in a build-up of the metals in edible plant parts. A study indicated that the concentration of DTPA-extractable and total heavy metals in flood plain soils around the Sutlej river was higher along the trans-boundary. In soil samples, 18 per cent of samples surpassed the limits of total Cd during both pre and post-monsoon seasons. The PLI values for total metal concentration were above one in 60 per cent of the soil samples. The

Comment [NS9]: State the sources

Comment [NS10]: State the sources

concentration of Co, Pb and Cd were higher in rice and wheat grains in contrast to their safe limits (Setia et al., 2021). Because heavy metals can bio-accumulate and bio-magnify throughout the food chain, eating polluted crops can put humans and animals at risk for health problems.

- e) **Implications for human health:** Exposure to heavy metals can be harmful to human health if they ingest polluted food, drink contaminated water, or breathe in contaminated dust. In India, high concentrations of As and Cr in urban soils exceed threshold values and represent serious non-carcinogenic health concerns for both adults and children. Residents may be at risk for cancer from Cr and As, while Pb levels usually remain within advised ranges. These health risks in urban areas require immediate attention as well as preventative measures (Adimalla, 2020).
- f) **Mitigation measures:** A number of measures such as soil testing to determine heavy metal levels and direct remediation activities can be used to lessen the pollution of agricultural soils with heavy metals. Adoption of integrated pest control and organic farming as sustainable farming methods to lessen the dependency on chemical inputs. Heavy metal immobilisation, extraction or degradation in polluted soils can be achieved using soil amendments and phytoremediation methods; it is a plant-based remediation which is an affordable and environment friendly approach to remove heavy metal and organic pollutant contamination from soil and water. Implementing the suitable waste management procedures to reduce the amount of heavy metals that are introduced into agricultural systems.

Comment [NS11]: Your citations are not uniform. Some are italicized while some are not.

Comment [NS12]: Where are your sources

Effects of heavy metals on crops

The intricate reactions between the main harmful ions and other necessary or non-essential ions produce toxic damage to cells and tissues when plants assimilate and accumulate high concentrations of heavy metals. Plant development does not require heavy metals (Ahmadpouret al., 2012; Shubhamet al., 2022). Research from Northern India suggests that leafy vegetables, such as fenugreek and spinach, have a higher concentration of heavy metals than fruit vegetables, including eggplant and chilli (Gupta et al., 2021). Since arsenic-polluted groundwater has been used to irrigate crops, especially rice crops (*Oryzasativa L.*), Bangladesh has greater soil arsenic levels. Amin et al., (2013) conducted a study to examine the accumulation of Cu, Ni, Zn, Cr, Fe, Mn, Co, and Pb in waste water-irrigated red onions (*Allium cepa*), garlic (*Allium sativum*), tomatoes (*Solanumlycopersicum*), and eggplants (*Solanummelongena*). Waste water irrigation significantly increased the degree of heavy metal pollution in crops when compared to tube well water. *A. cepa* that was irrigated with tube-well water did not have a manganese build up 50 times greater (28.05 mg/kg) than the severely polluted onions (Kgopaet al., 2018).

Comment [NS13]: The subheading is not well captured in the write-up. Please review

Effects of heavy metals on soils in India

Heavy metals such as lead, mercury, cadmium, arsenic and chromium pose a major danger to the stability of ecosystems and the health of the soil. Heavy metal accumulation can negatively impact crop productivity, soil quality and agricultural product quality all of which have an impact on the health of people, animals and the environment (Nagajyoti et al., 2010). Large areas of land might get contaminated by heavy metals released by waste incinerators, industrial smelters, municipal compost or sludge, pesticides, fertilisers, and industrial effluent. In the Surat industrial region, soil samples were extracted from the top 10 cm of the soil. The study revealed that the soils in the area were highly polluted, with a greater amount of toxic elements than normally seen. The study area's soils had heavy metal loadings of

Comment [NS14]: Please settle down and review this subject matter with recent citations. There are thousands of recent research works on this. The citations you have here simply mean that you did not research widely. Please kindly review.

139.0 mg/kg for Zn, 317.9 mg/kg for Sr, 380.6 mg/kg for V, 51.3 mg/kg for Co, 137.5 mg/kg for Ba, 305.2 mg/kg for Cr, and 51.3 mg/kg for Ni (Krishna & Govil, 2007). In another study conducted at Andhra Pradesh, environmental geochemistry studies were carried out in and around the Patancheru industrial growth region so as to determine the extent of chemical contamination in the soil. The data indicate how severely contaminated the area's soils are, with quantities of harmful substances two to three times higher than average. Above their usual distribution, a number of heavy metals, such as Cr, V, Fe, As, Cd, Se, Ba, Zn, Sr, Mo, and Cu, are discovered in the soil. The research region's soils showed heavy-metal loadings of 240 mg/kg, 235 mg/kg, 1,350 mg/kg, and 500 mg/kg for Cr, V, Ba, Cd, and Cu, respectively (Govilet *et al.*, 2001). To determine the impact of pollution in the study region in and around the Pali industrial growing area of Rajasthan, an environmental geochemistry study was conducted. Soil samples around the Pali industrial district were evaluated for Pb, Cr, Cu, Zn, Sr, and V using a Philips PW 2440 X-ray fluorescence spectrometer. Most of the samples were taken in the vicinity of the Bandi River and small streams that surrounded industrial areas. The soil around the industrial zone was found to contain significantly higher amounts of the metals Pb (293 mg/kg), Cr (240 mg/kg), Cu (298 mg/kg), Zn (1,364 mg/kg), Sr (2,694 mg/kg), and V (377 mg/kg) than their typical distribution in soil (Krishna & Govil, 2004). The microbial population in the Baula-Nuasahi mining belts in Keonjhar District, Orissa, was found to be decreased and the overburden soils exhibited low concentrations of nutrients (N, P, and K). It turns out that the agricultural lands nearby had become unsuitable for crop production due to the metal ions that had seeped into them (Dal *et al.*, 2011). It was found that the main sources of chromium contamination were seepage water and overburden dumps (Dhal and Pinaki, 2014).

To assess the extent of heavy metal pollution carried by urbanisation and industrial development, the soils of Hyderabad's Balanagar industrial area were examined. Significant contamination was found with amounts of toxic substances higher than those present in soils with a normal distribution. Copper (Cu), nickel (Ni), lead (Pb), zinc (Zn), cobalt (Co), vanadium (V) and chromium (Cr) concentrations ranged from 82.2 to 2,264 mg/kg, 31.3 to 1,040 mg/kg, 34.3 to 289.4 mg/kg, 57.5 to 1,274 mg/kg, 67.5 to 5,819.5 mg/kg, 8.6 to 54.8 mg/kg, and 66.6 to 297 mg/kg, respectively (Machendaret *et al.*, 2011). Toxic heavy metals pose a major danger to many countries worldwide in Karnataka's chromite mining districts, where mining operations have historically occurred. The soil pollution around chromite mines was analysed in this study using a variety of indicators to evaluate contamination levels. The heavy metal concentrations in soil samples from 57 locations close to Tagdur and Jambur, two closed and active chromite mines, and residential areas near Chikkondanahalli in the Nuggihalli Schist Belt of Karnataka, India, were examined using wavelength dispersive X-ray fluorescence spectrometry. The studied area showed elevated levels of cobalt (Co), nickel (Ni), and chromium (Cr) compared to the Soil Quality Guideline's recommended limits. The higher enrichment factor of these metals indicates an increased risk of heavy metal contamination, perhaps as a result of past mining operations (Krishna *et al.*, 2013).

Conclusion

The utilization of metallophytes derived from plants in phytoremediation is a sustainable and long-term way of minimizing the degree of heavy metal contamination in soils. These unique adaptations enable these specialized plants to thrive in metalliferous situations and effectively eliminate the metals from the environment around them. By reducing the bio-availability and toxicity of heavy metals, metallophytes significantly contribute to environmental restoration through processes such as the absorption of metals,

Comment [NS15]: This is not a conclusion. And this could be attributed to the fact that the subject matter has not been dealt with, no methodology, nothing. Please kindly revise.

translocation and increased antioxidant defences. Also the metallophytes offer a variety of benefits over conventional remediation techniques when used in phytoremediation. Phytoremediation is an environmentally friendly method that does not add to the existing pollutants in the environment, in contrast to chemical treatments. Metallophytes are excellent options for cleaning up the extremely contaminated regions that are challenging to restore using traditional approaches, as they are able to survive in hard environments where other plants might struggle. Their ability to adapt and bounce back from heavy metal contamination makes metallophytes a vital asset in efforts to restore ecosystems. Phytoremediation capability helps to ensure the sustainability of natural resources and lowers the environmental pollution, thus being consistent with the aims of the circular economy and sustainable resource management. The implementation of metallophytes into phytoremediation strategies offers a holistic approach to environmental conservation, emphasising the importance of biodiversity conservation and ecosystem resilience, in addition to offering an effective approach for cleaning the contaminated environments. As this field of research advances, further study of metallophyte species and their interactions with metals will help us better understand phytoremediation processes and enhance the effectiveness of this strategy for the global issue of heavy metal pollution.

References

- Pal A, Dutta S, Mukherjee PK, Paul AK. Occurrence of heavy metal resistance in microflora from serpentine soil of Andaman. *Journal of Basic Microbiology: An International Journal on Biochemistry, Physiology, Genetics, Morphology, and Ecology of Microorganisms*. 2005 Jun;45(3):207-18.
- Singh A, Agrawal M. Acid rain and its ecological consequences. *Journal of Environmental Biology*. 2007 Jan;29(1):15.
- Bhatti SS, Kumar V, Sambyal V, Singh J, Nagpal AK. Comparative analysis of tissue compartmentalized heavy metal uptake by common forage crop: a field experiment. *Catena*. 2018 Jan 1;160:185-93.
- Marg BZ. Hazardous metals and minerals pollution in India: Sources, toxicity and management. A Position Paper, Indian National Science Academy, New Delhi. 2011 Aug.
- Yang X, Feng Y, He Z, Stoffella PJ. Molecular mechanisms of heavy metal hyperaccumulation and phytoremediation. *Journal of trace elements in medicine and biology*. 2005 Jun 27;18(4):339-53.
- Järup L. Hazards of heavy metal contamination. *British medical bulletin*. 2003 Dec 1;68(1):167-82.
- Ernst WH. Evolution of metal tolerance in higher plants. *For Snow Landsc Res*. 2006;80(3):251-74.
- Epstein E. Silicon. *Annual review of plant biology*. 1999 Jun;50(1):641-64.

Comment [NS16]: Most of your references are outdated.

More so, your referencing style is not uniform. Which system of referencing did you adopt?

Majority of your references were not cited. Where did you get those references from? You can only reference the works cited. Up to four pages of references, how and why?

Yang X, Feng Y, He Z, Stoffella PJ. Molecular mechanisms of heavy metal hyperaccumulation and phytoremediation. *Journal of trace elements in medicine and biology*. 2005 Jun 27;18(4):339-53.

McLaughlin MJ, Parker DR, Clarke JM. Metals and micronutrients–food safety issues. *Field crops research*. 1999 Jan 1;60(1-2):143-63.

Shubham, Uday Sharma & Rajesh Kaushal (2023) Potential of Different Nitrification Inhibitors on Growth of Late Sown Cauliflower Var. Pusa Snowball K-1 and Behavior of Soil NH₄⁺ and NO₃⁻ in *TypicEutrochrept* Under Mid Hills of NW Himalayas, *Communications in Soil Science and Plant Analysis*, 54:10, 1368-1378, DOI: [10.1080/00103624.2022.2146130](https://doi.org/10.1080/00103624.2022.2146130)

Shubham, Uday Sharma & Rajesh Kaushal (2023) Effect of nitrification inhibitors on quality, yield and economics of cauliflower cv. PSB K1 in *TypicEutrochrept* under mid hills of North Western Himalayas, *Journal of Plant Nutrition*, 46:17, 4096-4109, DOI: [10.1080/01904167.2023.2220741](https://doi.org/10.1080/01904167.2023.2220741)

Shubham, Uday Sharma, Rajesh Kaushal and Yash Pal Sharma. 2022. Effect of Forest Fires on Soil Carbon Dynamics in Different Land Uses under NW Himalayas. *Indian Journal of Ecology* 49(6): 2322-2329. DOI: <https://doi.org/10.55362/IJE/2022/3828>.

Shubham, Uday Sharma and ArvindChahal. 2021. Effect of forest fire on ammonification and nitrification: A study under chir pine (*Pinusroxburghii*) forest areas of Himachal Pradesh. *Indian Journal of Ecology* 48(2): 376-380.

Antonovics J, Bradshaw AD, Turner RG. Heavy metal tolerance in plants. In *Advances in ecological research* 1971 Jan 1 (Vol. 7, pp. 1-85). Academic Press.

Woolhouse HW. Toxicity and tolerance in the responses of plants to metals. In *Physiological plant ecology III: Responses to the chemical and biological environment* 1983 Jan 1 (pp. 245-300). Berlin, Heidelberg: Springer Berlin Heidelberg.

Pehoiu G, Murarescu O, Radulescu C, Dulama ID, Teodorescu S, Stirbescu RM, Bucurica IA, Stanescu SG. Heavy metals accumulation and translocation in native plants grown on tailing dumps and human health risk. *Plant and Soil*. 2020 Nov;456:405-24.

Favas PJ, Pratas J, Varun M, D'Souza R, Paul MS. Phytoremediation of soils contaminated with metals and metalloids at mining areas: potential of native flora. *Environmental risk assessment of soil contamination*. 2014 Mar 26;3:485-516.

Singh S, Parihar P, Singh R, Singh VP, Prasad SM. Heavy metal tolerance in plants: role of transcriptomics, proteomics, metabolomics, and ionomics. *Frontiers in plant science*. 2016 Feb 8;6:165395.

Devi P, Kumar P. Concept and application of phytoremediation in the fight of heavy metal toxicity. *Journal of Pharmaceutical Sciences and Research*. 2020 Jun 1;12(6):795-804.

Sheoran V, Poonia P, Trivedi SK. Metal pollution in soil and plants near copper mining site. *Int. J. Geol. Earth Environ. Sci*. 2011;1(1):2277-081.

Malik S, Thavamani P, Beer M, Megharaj M, Naidu R. Microbial activity and diversity in long-term mixed contaminated soils with respect to polyaromatic hydrocarbons and heavy metals. *Journal of environmental management*. 2012 May 30;99:10-7..

Baker AJ, Walker PL. Ecophysiology of metal uptake by tolerant plants. Heavy metal tolerance in plants: evolutionary aspects. 1990 Jun 2;2:155-65.

Kafle A, Timilsina A, Gautam A, Adhikari K, Bhattarai A, Aryal N. Phytoremediation: Mechanisms, plant selection and enhancement by natural and synthetic agents. *Environmental Advances*. 2022 Jul 1;8:100203.

Gani A, Hussain A, Pathak S, Banerjee A. An empirical investigation on the elimination of heavy metals using bioremediation method for selected plant species. *Physics and Chemistry of the Earth, Parts A/B/C*. 2024 Jun 1;134:103568.

Sarkar SR, Majumdar A, Barla A, Pradhan N, Singh S, Ojha N, Bose S. A conjugative study of *Typhalatifolia* for expunge of phyto-available heavy metals in fly ash ameliorated soil. *Geoderma*. 2017 Nov 1;305:354-62.

Gupta N, Yadav KK, Kumar V, Cabral-Pinto MM, Alam M, Kumar S, Prasad S. Appraisal of contamination of heavy metals and health risk in agricultural soil of Jhansi city, India. *Environmental Toxicology and Pharmacology*. 2021 Nov 1;88:103740.

Setia R, Dhaliwal SS, Singh R, Kumar V, Taneja S, Kukal SS, Pateriya B. Phytoavailability and human risk assessment of heavy metals in soils and food crops around Sutlej river, India. *Chemosphere*. 2021 Jan 1;263:128321.

Adimalla N. Heavy metals pollution assessment and its associated human health risk evaluation of urban soils from Indian cities: a review. *Environmental geochemistry and health*. 2020 Jan;42(1):173-90.

Ahmadpour P, Ahmadpour F, Mahmud TM, Abdu A, Soleimani M, Tayefeh FH. Phytoremediation of heavy metals: A green technology. *African Journal of Biotechnology*. 2012;11(76):14036-43.

Gupta A, Sharma V, Sharma K, Kumar V, Choudhary S, Mankotia P, Kumar B, Mishra H, Moulick A, Ekielski A, Mishra PK. A review of adsorbents for heavy metal decontamination: Growing approach to wastewater treatment. *Materials*. 2021 Aug 20;14(16):4702.

Amin H, Arain BA, Amin F, Surhio MA. Phytotoxicity of chromium on germination, growth and biochemical attributes of *Hibiscus esculentus* L. *American Journal of Plant Sciences*. 2013 Nov 21;2013.

Kgopa PM, Mashela PW, Manyevere A. Accumulation of heavy metal in onion (*Allium cepa*) plants irrigated with treated wastewater under field conditions. *Research on Crops*. 2018;19(1):62-7.

Nagajyoti PC, Lee KD, Sreekanth TV. Heavy metals, occurrence and toxicity for plants: a review. *Environmental chemistry letters*. 2010 Sep;8:199-216.

Krishna AK, Govil PK. Soil contamination due to heavy metals from an industrial area of Surat, Gujarat, Western India. *Environmental monitoring and assessment*. 2007 Jan;124:263-75.

Govil P, Reddy G, Krishna A. Contamination of soil due to heavy metals in the Patancheru industrial development area, Andhra Pradesh, India. *Environmental geology*. 2001 Dec;41:461-9.

Krishna AK, Govil PK. Heavy metal contamination of soil around Pali industrial area, Rajasthan, India. *Environmental Geology*. 2004 Dec;47:38-44.

Dhal B, Das NN, Pandey BD, Thatoi HN. Environmental quality of the Boula-Nuasahi chromite mine area in India. *Mine Water and the Environment*. 2011 Sep;30:191-6.

Dhal PK, Sar P. Microbial communities in uranium mine tailings and mine water sediment from Jaduguda U mine, India: a culture independent analysis. *Journal of Environmental Science and Health, Part A*. 2014 May 12;49(6):694-709.

Machender G, Dhakate R, Prasanna L, Govil PK. Assessment of heavy metal contamination in soils around Balanagar industrial area, Hyderabad, India. *Environmental Earth Sciences*. 2011 Jul;63:945-53.

Krishna AK, Mohan KR, Murthy NN, Periasamy V, Bipinkumar G, Manohar K, Rao SS. Assessment of heavy metal contamination in soils around chromite mining areas, Nuggihalli, Karnataka, India. *Environmental earth sciences*. 2013 Sep;70:699-708.

UNDER PEER REVIEW