

IMMUNOGLOBULIN BASED THERAPEUTIC APPROACH FOR PARVO VIRUS INFECTION IN DOG - A CASE REPORT

ABSTRACT

Canine parvo viral infection is a highly contagious disease in dog with high mortality rate in untreated cases. This study represents a case report of an unvaccinated 4 months old German shepherd puppy presented with a history of profuse foetid diarrhoea, vomiting, inappetence and depression for 2 days. Physical examination of the dog revealed sub normal temperature (99.5°F), abdominal pain, bloat, severe dehydration and debility. Haematology showed low RBC count, low Hb level, low Hct value, lower MCV and mild lymphopenia. The dog was diagnosed as canine parvoviral infection based on the result of rapid antigen test and the characteristic clinical findings. The line of treatment was given along with immunoglobulins. The pup showed good response from 3rd day of the treatment and finally recovered successfully. Thus, administration of specific immunoglobins along with standard treatment regimen in canine parvo virus infection can be effective in reducing mortality in dogs.

The method should more detailed in abstract section

Keywords: Canine Parvovirus Infection, Rapid antigen test, Immunoglobulin

1. INTRODUCTION

Canine parvovirus (CPV) infection caused by Canine parvo virus 2 (CPV2) is a highly contagious and a serious life-threatening viral disease in dog with high morbidity (100%) and varying mortality up to 10-18% in adults and 91% in pups if left untreated [1, 2, 3, 4]. CPV mainly occurs in two prominent clinical forms: (i) haemorrhagic gastroenteritis with vomiting, foul smelling bloody diarrhoea and severe leukopenia affecting all age group of dogs [5, 6, 7], (ii) myocarditis and subsequent heart failure in pups up to 3 months of age [8]. CPV 2 is extensively shed in faeces for 7–12 days or possibly for long-term duration, which help it to spread rapidly among the population [9]. Prevalence of CPV infection in Assam was 25.28 to 62.86% [10, 11, 12] with 41.86% in 4-6 months age group (Deka *et al.*, 2013), 51.85% in 7-12 months of age group, 50.18% in male and was highest in German spitz breed with 55.21% [11]. CPV infection was recorded higher in unvaccinated dogs with 30.15%

compared to 9.52% in vaccinated dogs [4]. Season-wise, occurrence of CPV was recorded highest in pre-monsoon (43.18%) followed by winter (34.62%) [4].

2. PRESENTATION OF CASE

One four months old, weaned German shepherd puppy was brought with a history of profuse foetid smell diarrhoea, vomiting, dull, depressed and inappetence for 2 days. During physical examination, the puppy had sub normal temperature (99.5°F), severe dehydration, sunken eyes, severe debilitation, abdominal pain and bloat. The puppy along with its mother was unvaccinated and not dewormed regularly. Blood sample was collected and sent for haematology. The faeces were collected and subjected for rapid immunochromatographic based antigen (Ag) test (Quickvet, Ubio) for detection of canine parvo virus Ag. The affected puppy had low RBC count ($4.2 \times 10^6/\text{mm}^3$), low Hb level (8.3 g/dl), low Hct value (21.2 %), lower MCV (21.2 %) and mild lymphopenia (9.3%). Based on the result of rapid antigen test and clinical findings, the diagnosis was made as canine parvo virus infection and accordingly treatment had been commenced.

The case which showed in this section including the visual report such as some pictures in jpg

3. DISCUSSION

The treatment was started with intra venous (IV) fluid replacement therapy that included 5% DNS and Ringer's lactate infusion to restore the fluid & electrolyte loss. Pantoprazole was administered IV @ 1 mg per kg body weight (b. w.) OD for 10 days along with Ondansetron @ 0.1 mg per kg b. w. IV BID for 7 days. Gentamicin was given @ 10 mg per kg b. w. slow IV OD for 7 days to lower the possibility of bacterial translocation across the disrupted intestinal epithelium in the gastrointestinal tract. Metronidazole was injected @ 10 mg per kg b. w. slow IV for 7 days to lower the intestinal parasites which may exacerbate the infection by increasing intestinal cell turnover and subsequent viral replication [13]. Tranexamic acid was also given slow IV @ 10 mg per kg b. w. of the animal. Based on the severity of the infection, purified hyperimmune immunoglobulin (Immunoglobulinum anti parvovirus canis NLT 1024 HIU, Canglob ® -P) was administered IV @ 0.4 ml per 1 kg of body weight of the animal for 5 days. The pup showed good response from 3rd day of treatment. Injecting specific immunoglobulins at the clinical phase of canine parvo virus infection can improve the clinical symptoms, reduce mortality and subsequently facilitate rapid recovery of the patient [14, 15, 16]. Intravenous administration of Immunoglobulin, IgY derived from chicken egg yolk had been reported to inhibit the propagation of CPV infection and thus reduce the fatality [15]. In another study, freeze-dried immunoglobulin G was found to diminish clinical symptoms and shorten hospitalization duration in naturally acquired CPV enteritis [17].

The discussion should be focused the novelty of studi in case report

4. CONCLUSION

Based on the present case study, it may be concluded that early administration of specific immunoglobins in canine parvo virus infection can be effective in ameliorating the clinical symptoms and reducing mortality.

ACKNOWLEDGEMENT

The authors declare no sponsorship to carry out the study.

AUTHORS' CONTRIBUTIONS

This work was carried out in association with all authors.

COMPETING INTERESTS

Authors declare that there are no competing interests regarding this work.

REFERENCES

1. Appel MJG, Scott FW and Carmichael LE. 1979. Isolation and immunization studies of a canine parvo-like virus from dogs with hemorrhagic enteritis. *Vet Res.*; 105:156–159.
2. Black JW, Holscher MA, Powell HS and Byerly CS. 1979. Parvoviral enteritis and panleucopenia in dogs. *J Med Small Anim Clin.*;74:47–50.
3. Nandi S & Kumar M. 2010. Canine Parvovirus: Current Perspective Indian J. Virol. 21(1):31–44 DOI 10.1007/s13337-010-0007-y.
4. Deka D, Phukan A and Sarma DK. 2013. Epidemiology of parvovirus and corona virus infections in dogs in Assam. *Environmental Science, Medicine*, 90(9):49-51.
5. Appel MJG, Cooper BJ, Greisen H and Carmichael LE. 1978. Status report: canine viral enteritis. *J Am Vet Med Assoc.*; 173:1516–1518.
6. Woods CB, Pollock RVH and Carmichael LE. 1980. Canine parvoviral enteritis. *J Am Anim Hosp Assoc.*; 16:171–179.
7. Shackelton LA, Parrish CR, Truyen U and Holmes EC. 2005. High rate of viral evolution associated with the emergence of carnivore parvovirus. *Proc Natl Acad Sci USA.*; 102:379–384.
8. Hayes MA, Russell RG, Mueller RW and Lewis RJ. 1979. Myocarditis in young dogs associated with a parvovirus-like agent. *Can Vet J.*; 20:126–132.
9. Cho HS, Song JE and Park YS. Diagnosis of the canine parvovirus in fecal samples by in situ hybridization. *Indian Vet J.* 2004; 81:855–859.
10. Phukan A, Deka D and Boro PK. 2004. Occurrence of canine parvovirus infection in and around Guwahati. *The Indian Journal of Animal Sciences*74(9):930-931.
11. Phukan A, Baishya B, Deka D and Boro PK. 2010. Prevalence of canine parvovirus infection in Assam. *The Indian veterinary journal*, 87(10):972-974.
12. Deka D, Phukan A and Sarma DK. 2014. Diagnosis of canine viral gastroenteritis. *The Indian Veterinary Journal*, 91(3):36-39.
13. Douglass Macintire. Treatment of severe parvoviral enteritis (Proceedings). 2008, August.

14. Nguyen SV, Umeda K, Yokoyama H, Tohya Y, Kodama Y. Passive protection of dogs against clinical disease due to Canine parvovirus-2 by specific antibody from chicken egg yolk. *The Can. J of Vet. Res* 2006; 70:62-64.
15. Suartini GAA, Sendow I. Prospek Pemanfaatan Imunoglobulin Y untuk Terapi Infeksi Canine Parvovirus pada Anjing. *WARTAZOA* 2015;25N(2):55-64.
16. Rishikesavan R, Palanivel KM, Saravanajayam M. Successful treatment of canine parvoviral infection with imunoglobulins in a pup. *The Pharma Innovation Journal*. 2021;10(1):27-8.
17. Macintire D, Smith-Carr S, Jones R, et al. Treatment of dogs naturally infected with canine parvovirus with lyophilized canine IgG. *Proceedings 17th Annual Conference of the American College of Veterinary Internal Medicine, Chicago, USA, June 10–13, 1999*, p 721.

UNDER PEER REVIEW