

Impact of dolutegravir as first-line antiretroviral treatment on the tolerance and virological suppression among PLHIV in the southern region of Senegal

Abstract:

Introduction??????

Aims: To determine the virological suppression and the tolerance on patients under a dolutegravir-based regimen by comparison with patients on a regimen without dolutegravir after at least 6 months of treatment.

Materials and methods: A retrospective, descriptive, analytical and comparative study covering data from 469 patients with 219 treatment-naïve patients under a dolutegravir based regimen and 250 patients pretreated with triple therapies not including dolutegravir, followed at the HIV treatment center of Kolda from January 1st, 2017 to December 31st, 2022. The inclusion criteria were: Patients HIV-1 and/or HVI-2 positive, aged > 15 years, with a viral load achieved after at least 6 months of ARV treatment. Undetectable threshold is defined as a viral load < 50 copies/ml. data analysis??

Results: The proportions of patients screened at an advanced clinical stage of AIDS (WHO stage 3 or 4) were significant in both groups of patients under dolutegravir and without dolutegravir respectively (63.47% versus 59.20%, P= 0.344). The mean duration on ARV treatment was significantly higher in patients pretreated with therapeutic regimens not including dolutegravir (55.67±11.93 versus 28.95±7.80) (P < 0.0001) (Table 2). The dolutegravir-based regimen combined tenofovir, lamivudine and dolutegravir while the majority (90%) of pretreated patients were on an ARV regimen combining TDF, lamivudine, Efavirenz in first line (Table). Fewer patients presented side effects due to triple therapy in the dolutegravir-based regimen group compared to the non-dolutegravir group (2.74% vs. 3.20%; p=0.77). The rate of undetectable viral load was slightly higher in the dolutegravir-based regimen group, compared to the group without dolutegravir but the difference was not statistically significant (97.72% vs 97.60%; p = 0.933). There was less Virological failure (viral load >1000 copies/ml) in the dolutegravir group than in the group without dolutegravir (1 versus 3). The mean of the body mass index did not differ in the two groups. The weight gain was lower in the dolutegravir based regimen group but the difference was not statistically significant 6,7± 8,2 kg versus 7,2 ± 6,2 kg. The occurrence of obesity was not significantly different in the two groups [23.74%] versus [23.20%], P = 0.89).

Conclusion: The results of our study show a high rate of virological success, and good tolerance of the dolutegravir-based regimen in a decentralized setting.

Keywords: Dolutéglavir, Treatment, PLHIV, Senegal

Introduction:

Sub-Saharan Africa remains the most heavily affected region by HIV in the world, holding more than two-thirds (67%) of all people living with HIV [1].

Comment [JO1]: This is a comparative study (DTG vs non DTG containing regimen. This needs to be clear from the topic. Reframe as « Impact of Dolutegravir-based regimen on tolerance and virologic suppression, compared to non-dolutegravir regimen among PLHIV in Southern Senegal

Comment [JO2]: Delete colon » : »

Comment [JO3]: Include introduction maybe in one or two sentences

Comment [JO4]: A retrospective cohort analysis involving 469 patients of which 219 were treatment-naïve individuals initiated on Dolutegravir (DTG) based regimen and 250 were previously treated with non-DTG based triple therapy

Comment [JO5]: Inclusion criteria unnecessary at this point, Just state...The study included HIV infected patients aged 15 years and above who had been on antiretroviral therapy after at least 6 months

Comment [JO6]: Why did you only include those who had achieved viral suppression? Did you mean patients on HIV treatment after at least 6 months of ART.

Comment [JO7]: Include data analyses, how was the data analysed? e.g technique used, e.g chi square, tool used, e.g stata vs 13

Comment [JO8]: Delete, should NEVER appear in an abstract

Comment [JO9]: delete

Comment [JO10]: what were these side effects? mention them

Comment [JO11]: The whole abstract could be re-organized to bring out clearly the comparisons between DTG based and non DTG based regimen. Include statistical analysis method used

With the significant progress in antiretroviral therapy, HIV infection has become a chronic infection which, failing to be eradicated can be controlled. Better tolerated and/or more effective ARVs on sensitive or resistant viruses have been developed in recent years, belonging either to old classes or to new classes, such as integrase inhibitors and CCR5 inhibitors.

The effectiveness of integrase inhibitors has been established in studies of treatment-naïve and pretreated patients [3,4]. Dolutegravir as a first-line treatment has shown an improved safety profile compared to EFV which is a non-nucleoside reverse transcriptase inhibitor (NNRTI) [4]. It is estimated that around 15 million people will be taking Dolutegravir by 2025 and it will replace first-line Efavirenz-based regimens [5,6]. These ARVs have a beneficial role not only in terms of individual health but also in the drastic reduction of contagiousness when plasma viral replication is controlled under treatment. This fundamental data regarding the possible reduction of transmission associated with the individual benefits of earlier treatment led to the concept of TasP (Treatment as Prevention) and Test and Treat [2].

Senegal is one of the first countries in sub-Saharan Africa to set up an access program to ARV treatments in 1998. It is also one of the first countries in Africa to have decided to make ARV treatments free in 2003. ???references

In 2020, Senegal began a transition to Dolutegravir-based treatment regimens in accordance with WHO recommendations. This regimen was first initiated in newly infected patients then gradually in patients under other protocols. After three years of switching to dolutegravir, we conducted this study with the aim of comparing the tolerance and effectiveness of these new protocols to those pre-existing regimens, especially based on NNRTI in a decentralized environment.

Methodology :

The study took place in the HIV outpatient treatment unit at Kolda's health center (UTA) from January 1st, 2017 to December 31st, 2022.

It is a descriptive, analytical and comparative observational study based on data from patients who initiated their first-line treatment with the dolutegravir-based ARV regimen between April 2020 and December 2022. Our comparison group consisted of PLHIV pretreated with an ARV regimen without dolutegravir. The duration of exposure to dolutegravir had to be at least 6 months. Each patient included had to have a virological assessment available after at least 6 months of ARV treatment. We excluded pretreated PLHIV who had switched to dolutegravir from the comparison group. Patients whose files were unusable due to lack of information were not included.

Comment [JO12]: Two sentences are too short to make a paragraph. Also include references

Comment [JO13]: In the Methodology section split it into two.
1. Study design and participants
2. Data analysis
Then add Ethical Consideration

Comment [JO14]: Not clear, edit please

Comment [JO15]: The patients in DTG group were followed up for 3 years, those in non DTG group follow up period is not indicated

At what point was the outcome (viral suppression, side effects measured) have in mind that some experience blips
What was the definition of viral suppression in this study? i.e. after how many viral load measurements did you consider it as suppressed. How did you address blips

Did you experience virologic failure, if yes, how many per treatment arm?

The viral load was considered undetectable when it was less than 50 copies per milliliter of blood. Viral suppression was a viral load below 1000 copies per milliliter of blood.

Patients' medical records in the center's cohort served as the source of data collection. The data was collected on a survey sheet. Each survey sheet was assigned an anonymous number. To meet our objective, the epidemiologic, clinical parameters (age, sex, level of education, duration of HIV infection, clinical stage at discovery of HIV, weight, presence of opportunistic infection, side effects) and virological parameters (viral load) were studied. A Chi test allowed the comparison of qualitative variables. The significance threshold was set at 5%.

Data were entered into Excel 2016 after cleaning and coding, then analyzed using the Stata version 13 software, before categorizing the different variables according to the defined methods.

Ethical Consideration????

Results :

A total of 469 PLHIV were included in this study, with 219 who initiated treatment with a dolutegravir-based regimen and 250 with other regimens. The patients' average age did not differ significantly between the two groups (41.63 ± 12.89 versus 39.56 ± 12.98 , $P=0.085$) respectively. We also noted in both groups a female predominance (73.52% versus 78.00%, $P= 0.257$), a significant proportion of married people (77.31% versus 72.29%, $P= 0.214$), and patients who haven't shared their HIV status with their partners (27.06% versus 34.60%, $P= 0.083$). The majority of patients were illiterate (50.68% versus 57.20%, $P = 0.158$). The proportions of comorbidities found: the proportion of diabetes did not differ in the two groups (7.76% versus 8.00%, $P = 0.924$), however hypertension was significantly higher among patients on ARV based on dolutegravir compared to patients on other regimens (21.00% versus 11.60%; $P: 0.006$). The proportions of patients presenting weight loss ($BMI < 18.5 \text{ kg/m}^2$) at inclusion did not differ significantly between the two groups (47.03% versus 43.20%, $P = 0.405$). HIV-HBV co-infection was similar in both groups (5.48% versus 5.60%, $P= 0.545$). The proportions of patients screened at an advanced clinical stage of AIDS (WHO stage 3 or 4) were significant in both groups (63.47% versus 59.20%, $P= 0.344$). The mean duration on ARV was significantly higher in patients pretreated with therapeutic regimens not including dolutegravir (28.95 ± 7.80) versus 55.67 ± 11.93) ($P < 0.0001$) (Table 2). The dolutegravir-based regimen combined tenofovir, lamivudine and

Comment [JO16]:

Considering appropriate statistical technique when comparing the two treatment groups e.g T test ANOVA, logistic regression to make your study robust
The Chi square test used in the entire study is not advisable

Comment [JO17]: Add ethical consideration

Comment [JO18]: Include the results table within the results

Comment [JO19]: The table (i believe table 2) supporting this narrative is missing hence it is difficult to comment

dolutegravir while the majority (90%) of pretreated patients were on an ARV regimen combining TDF, lamivudine, Efavirenz as first line treatment (Table 1). Fewer patients experienced side effects related to triple therapy in the dolutegravir-based regimen group compared to the non-dolutegravir group (2.74% vs 3.20%; $p=0.77$).

The undetectable viral load rate was slightly higher in the dolutegravir-based regimen group compared to the non-dolutegravir regimen group, but the difference was not statistically significant (97.72% vs 97.60%; $p = 0.933$). Virological failure (Viral load > 1000 copies/ml) was less frequent in the dolutegravir group than in the group without dolutegravir (1 versus 3). Mean body mass index did not differ in the two groups. The weight gain was lower in the dolutegravir based regimen group but the difference was not statistically significant $6,7 \pm 8,2$ kg versus $7,2 \pm 6,2$ kg. The occurrence of obesity was not significantly different in the two groups [23.74%] versus [23.20%], $P = 0.89$.

Discussion

In this study, we compared data on virological suppression and tolerability of dolutegravir based regimens of 219 treatment-naive patients with those of 250 patients pretreated with a non-dolutegravir regimen for HIV-1 and HIV-2 in decentralized and resource-limited settings.

We observed in both groups a female predominance (75.9%) and an average age of 40 years. These results are in agreement with data from the CNLS (National council for the fight against AIDS) 2021 in Senegal [13]. This female predominance has also been found in several studies in Africa: [7, 8, 9, 10, 11, 12]. Indeed, HIV infection mostly affects the sexually active age group corresponding to the population of young adults due to their exposure to certain risk factors, especially economic insecurity and the relaxation of social control, notably in urban areas. However, a meta-analysis of 13 cohorts by Egger [14] bringing together European and North American countries suggested a male predominance. This difference between the countries of the North and the South finds two explanations: on one hand, the predominance of heterosexual transmission in the South associated with female vulnerability; on the other hand the relatively high role of homosexuality and injection drug use in the North in the transmission of the disease.

Most patients in both groups were detected at an advanced clinical stage of AIDS (WHO stage 3 or 4) (63.47% versus 59.20%, $P = 0.344$). This diagnostic delay has been observed in several African studies [10, 11]. The advanced clinical stage is a vulnerability factor for patients both to the disease itself and

Comment [JO20]: Which statistical technique was used to make comparisons if it is difficult to make a comment .

Comment [JO21]: I find this statement « misplaced ». The statement above it looked at gender hence you need to focus on that and not age group

Comment [JO22]: Majority of HIV infection in both groups were detected at an advanced clinical stage

Comment [JO23]: This is no longer the case now

the opportunistic conditions. It illustrates a diagnostic delay which remains frequent in our developing countries; the reasons for this delay can be explained by poverty, lack of information, the attitude of hiding the disease because of the risk of stigmatization and discrimination from the populations, the geographical and especially financial inaccessibility to healthcare structures [10]. This despite the freeness and efforts made to decentralize the screening in Senegal.

The proportion of weight loss (BMI < 18.5 kg/m²) at inclusion was significant and did not differ significantly between the two groups (47.03% versus 43.20%, P = 0.405). Studies have revealed that in the very advanced stages of AIDS there is a reduction in nutritional intake, an increase in energy expenditure linked to basal metabolism due to viral replication [15,16,17]. This viral presence triggers proteolysis which leads to muscle damage, thus causing an imbalance in the nitrogen balance and an exaggerated loss of lean tissue [15].

According to the serological profile, HIV type-1 is predominant with 88.91% (n = 469). This high prevalence of HIV-1 is explained by its greater virulence, which is characterized by greater transmissibility both sexually and from mother to child. This is how HIV-1 constitutes the serotype solely responsible for the current pandemic [18].

After monitoring patients on ARV for at least six months, we observed a progressive increase in weight gain in both groups. This increase was slightly greater in the group without dolutegravir with a mean weight gain of 7.2 ± 6.2 and a BMI of 22.42 ± 4.5 kg/m² compared to a mean weight gain of 6.7 ± 8.2 kg and BMI 22.19 ± 4.4 in the dolutegravir group but the difference was not statistically significant (p: 0.58). This variation in weight gain in the two groups could be due to the fact that the average duration of ARV treatment is longer in the group without dolutegravir (55.67 ± 11.93 versus 28.95 ± 7.80; (P < 0.0001). The usage of anthropometrics parameters including BMI as a marker of the effectiveness of ARV treatment has been described and remains important in settings with limited resources [19,20] [21, 22]. This weight gain in our study is close to the one found in patients followed in South Africa [21] [23]. However, several studies have found the occurrence of significant weight gain under ARV based on dolutegravir in Bouaké[22,24,23,25]. Dolutegravir would therefore play a role in patients weight gain. According to the literature, various mechanisms explain weight gain in PLHIV. Weight gain has recently appeared as a main side effect of antiretroviral treatment, potentially increasing the associated cardiovascular risks and having a negative impact on the quality of life of people living with HIV. This weight gain undoubtedly has a

Comment [JO24]: This is no longer the case now as most infections are promptly identified. If you were doing this study in the last 15 years this would hold, even the references is 2009, >10 years old. Consider revising this statement to depict the current trends, use newer references less than 5 years old. You could argue it based on UNAIDS first 95% initiative (95% of PLHIV knowing their status)

Comment [JO25]: Avoid repeating the results in the discussion

multifactorial origin and is of variable magnitude depending on the anti-integrase, the associated ARVs, the sex, the previous weight, the CD4 nadir and ethnic origin. We studied, in patients switching from triple therapy based on dolutegravir (DTG) to dual therapy, the evolution of weight gain [26]. Indeed, HIV infection and ARV treatment itself would lead to intestinal lesions causing an increase in fungal translocation and therefore an increase in the Beta D glucan (BDG) marker. A higher BDG would be associated with greater fat gains under antiretroviral treatment [27].

Retention for 12 months was significant among patients newly initiated to ARV treatment including dolutegravir in our cohort. This finding was observed in a therapeutic trial where people initiated on dolutegravir were more likely to be retained in care for 12 months (adjusted RR 1.09, 95% CI 1.04 to 1.14; adjusted risk difference 5.2%, 2.2 to 8.4). This high rate of retention of patients on dolutegravir-based ARV could be explained by increased tolerance [28,29,30]. Indeed, in our study, fewer patients experienced side effects linked to triple therapy in the dolutegravir-based regimen group compared to the group without dolutegravir (2.74% vs 3.20%; $p = 0.77$).

The virological success rate (measured by the proportion of participants with a viral load <50 copies per milliliter after at least 6 months on ARV) was slightly higher in the dolutegravir-based regimen group, compared to the non-dolutegravir regimen group however there was not a statistically significant difference (97.72% vs 97.60%; $p = 0.933$). This virological success rate observed in our study was higher compared to those found in several others: 88% in the group having received dolutegravir, abacavir and lamivudine in the SINGLE2 trial [31] and 90% in the group that received dolutegravir and two NRTIs in the FLAMINGO trial [32]. Studies have shown viral suppression was impaired in participants with a high baseline viral load. However, in our study, the baseline viral load was not measured in our patients before starting ARV. This could explain the difference in virological success rates. Furthermore, an improvement in viral suppression over 12 months with dolutegravir was observed in treatment-naïve patients in two clinical trials in Africa. [28,33]. Public health data from Brazil showed good safety outcomes [34] and better viral suppression over 12 months in people initiated on dolutegravir compared to those on efavirenz [35], but the proportion on dolutegravir was low and there was no assessment of retention in care.

In our study we did not look for resistance to ARVs, particularly to dolutegravir. Studies from Malawi, Lesotho, and Uganda suggest low levels of HIV resistance mutations to dolutegravir and high levels of viral suppression in

people who transitioned to dolutegravir, but did not compare results with those of people remaining on dolutegravir-free regimens [35,36,37]. A retrospective cohort study of 3,108 people from four African countries have found that people who transitioned to dolutegravir had better viral suppression than those who remained on the same first- or second-line treatment.[38] . Dolutegravir-based regimens are associated with a reduced need to switch to other antiretrovirals and there is a lower risk of developing major antiretroviral resistance mutations compared to efavirenz-based regimens [39,40].

Conclusion:

This study showed that patients on treatment based on dolutegravir have a high rate of virological success, a low rate of virological failure and good tolerance similar or better than other regimens mainly with efavirenz. Other **diets** even though the improvements in terms of retention in care and viral suppression are modest, constitutes an asset for achieving the 95-95-95 objectives in our context.

Table 1 : Demographic characteristics of DTG and other diets

Characteristics	DTG diet	Other diets	P-value
Gender			
Female	161 (73.52%)	195 (78.00%)	0,257
Male	58 (26.48%)	55 (22.00%)	
Age (years)			
<35	59 (26.94%)	72 (28.80%)	
35 - 49	91 (41.55%)	85 (34.00%)	
>50	69 (31.51%)	93 (37.20%)	
Sharing of HIV viral status with partner			
Yes	159 (72.94%)	155 (65.40 %)	0.083
No	59 (27.06%)	82 (34.60%)	
Clinical stage WHO			0,344
1	41 (18, 72%)	45 (18, 00%)	
2	39 (17, 81%)	57 (22, 80%)	
3	118 (53, 88 %)	125 (50, 00%)	
4	21 (9, 59 %)	23 (9, 20%)	
Co-morbidity			
High blood pressure	46 (21.00%)	29 (11.60%)	0.006

Comment [JO26]: Remove colon « : »

Comment [JO27]: What does « diet » mean ?

Comment [JO28]: Are these demographic characteristics of DTG ? Change to

Demographic Characteristics of Respondents

Comment [JO29]: I fail to understand the term « Diet » what does it mean ? Did you mean drug / regimen ?

Comment [JO30]: HIV status disclosure

Comment [JO31]: Change to WHO Stage at enrolment

Comment [JO32]: Change to Hypertension

Diabetes	17 (7.76 %)	20 (8.00%)	0.924
Dual infection HIV-HBV	12 (5, 48%)	14 (5, 60%)	
HIV profile			
Type 1	194 (88.58%)	223 (89.20%)	0.576
Type 2	18 (08.21%)	17 (06.8%)	
Type 1+2	7 (03.17%)	10 (04.00%)	
BMI at admission			
Mean	22.19 ±4.4	22.42 ± 4.5	
< 18,5	103 (47.03%)	108 (43.20%)	0.443
18,5- 24,9	87 (39.73%)	114 (45.60 %)	
25 – 29,9	23 (10.50%)	24 (9.60%)	
30 – 34,9	6 (2.74%)	3 (1.20 %)	
35 – 39,9	0 (0.00 %)	1 (0.40%)	
Mean ARV treatment duration	28.95± 7.8	55.67± 11.93	0.0000
Virological success <50 copies/ml	212 (96.80%)	241 (96.40%)	0.810
Virological failure >1000 copies/ml	1 (0.46%)	3 (1.20%)	
Sides effects			
Sides effects sides	6 (2.74%)	8 (3.20%)	
Dizziness	1 (0.46%)	1 (0.40%)	
Acute diarrhea	1 (0.46%)	1 (0.40%)	
Vomiting	2 (0.91%)	0 (0.00%)	
Nausea	0 (0.00%)	2 (0.80%)	
Headache	0 (0.00%)	1 (0.40%)	
Insomnia	2 (0.91%)	2 (0.80%)	
Skin rash	0 (0.00%)	1 (0.40%)	
ARV regimen			
TDF+3TC+DTG	219 (100%)	0	
TDF+3TC+EFV	0 (0.00%)	225 (90%)	
TDF+3TC+LPv	0 (0.00%)	12 (4,80%)	
AZT+3TC+LPv	0 (0.00%)	8 (3,20%)	
AZT+3TC+NVP	0 (0.00%)	4 (1,60%)	
ABC+3TC+EFV	0 (0.00%)	1 (0,40%)	
Patients after 12months			
Regularly followed	172 (78.54%)	245 (98.00 %)	

Comment [JO33]: HIV-HBV Co-infection

Comment [JO34]: How did you address blips ?

Comment [JO35]: Delete « s »

Comment [JO36]: Edit please

Lost of sight	0 (0,00%)	4 (1.60%)	
Transfers	47 (21.46%)	1 (0.40%)	
Mean BMI after at least 12 months of ARV treatment	22.19 ±4.4	22.42±4.5	0.581
Meanweight gain	6,7± 8,2	7,2 ± 6,2	0,503

Comment [J037]: Meanweight gain after how long

References:

1 UNAIDS. Report on global HIV/AIDS epidemic. 2023

2 Pialoux G, Lert F. Prevention et reduction des risques dans les groupes à haut risque vis-à-vis du VIH et des IST. Rapport « Reduction des risques sexuels.VIH&IST », 25 novembre 2009 (http://www.vih37.org/IMG/pdf/rapport_lert-pialoux.pdf).

3 Aboud M, Kaplan M, Lombaard J et al. Superior efficacy of dolutegravir (DTG) plus 2 nucleoside reverse transcriptase inhibitors (NRTIs) compared with lopinavir/ritonavir (LPV/RTV) plus 2NRTIs in second-line treatment: interim data from the DAWNING study. International AIDS Society Conference on HIV Science. July 2017. Paris, France. Abstract TUAB0105LB.

4 Walmsley S, Baumgarten A, Berenguer J et al. Brief report: Dolutegravir plus abacavir/lamivudine for the treatment of HIV-1 infection in antiretroviral therapy naïve patients: week 96 and week 144 results from the SINGLE randomised clinical trial. *J Acquir Immune Defic Syndr* 2015; 70: 515–519

5 Clinton Health Access Initiative. ARV market report; 2017. Available at: https://clintonhealthaccess.org/content/uploads/2017/09/2017-ARV-Market-Report_Final-2.pdf (accessed March 2018).

6 World Health Organization. Transition to new antiretrovirals in HIV programmes; 2017. Available at: <http://apps.who.int/iris/bitstream/10665/255888/1/WHO-HIV-2017.20-eng.pdf?ua=1> (accessed March 2018).

7 Dorward J, Sookrajh Y, Khubone T, van der Molen J, Govender R, Phakathi S, Lewis L, Bottomley C, Maraj M, Lessells RJ, Naidoo K, Butler CC, Van Heerden R, Garrett N. Implementation and outcomes of dolutegravir-based first-line antiretroviral therapy for people with HIV in South Africa: a retrospective cohort study. *Lancet HIV*. 2023 May;10(5):e284-e294

8 Koy, T., Mukumbi, H., Muteba, G. L., Donnen, P., & Wilmet-Dramaix, M. (2014). Profil comparatif et évolutif des personnes infectées par le virus de l'immunodéficience humaine traitées aux antirétroviraux à Kinshasa, République Démocratique du Congo. *Pan African Medical Journal*, 19.

- 9 Matthew P, Fox M and Rosen S. Patient retention in antiretroviral therapy Programs up to three years on treatment in sub-Saharan Africa: 2007-2009, systematic review. *Tropical medicine and International Health*. June 2010; 15 (s1): 1-15
- 10 Manga NM, Diop SA, Ndour CT, Dia NM, Mendy A, Coudec M, et al. [Late diagnosis of HIV infection in the Fann, Dakar clinic of infectious diseases: testing circumstances, therapeutic course of patients, and determining factors]. *Med Mal Infect*. 2009;39:95-100
- 11 Fortes Déguénonvo L, Manga NM, Diop SA, DiaBadiane NM, Seydi M, Ndour CT, et al. Profil actuel des patients infectés par le VIH hospitalisés à Dakar (Sénégal). *Bull Société Pathol Exot*. déc 2011;104(5):366-70
- 12 . Chas J, Hema A, Slama L, Kabore F, Lescure FX, Fontaine C, et al. The Day Hospital of the University Hospital, Bobo Dioulasso: An Example of Optimized HIV Management in Southern Burkina Faso. *PloS one* 2015;10:e0125588
- 13 CNLS. Rapport annuel 2021 : <https://www.cnls-senegal.org/wp-content/uploads/2022/06/Rapport-CNLS-2021-1.pdf>
- 14 Egger M, Sterne J, Phillips A. Prognostic importance of initial response in HIV1 infected patients starting potent antiretroviral therapy: analysis of prospective studies. *Lancet* 2003; 362: 679-86.
- 15 .Macallan DC, Margaret A, Nurlan MC, Milne E, Graham CA, Garlick PJ. Whole-body protein turnover from leucine kinetics and the response to nutrition in human immune deficiency virus infection. *Am J Clin Nutr*. 1995;61(4):818-826. [PubMed] [Google Scholar]
- 16 .Yarasheski X, Kevin E, Jeffrey J, Zachwieja, Jennifer Gischler, Jan Crowley, Mary M Horgan, William G Powderly. Increased plasma Gln and Leu rate and inappropriately low muscle protein synthesis rate in AIDS wasting. *J. Physiol*. 1998;275(38):E577-83. [Article PMC gratuit] [PubMed] [Google Scholar]
- 17 . Mulligan KVV, Tai, Chamberlan M. Energy expenditure in Human Immunodeficiency virus infection. *N Engl J Med*. 1997;336:70-71. [PubMed] [Google Scholar]
- 18 OMS. Rapport VIH 2022.
- 19 . World Health Organization. Working Document on Monitoring and Evaluating of National ART Program in the Rapid Scale up to 3. Disponible et consulté le 25 Août 2013
- 20 Messou E, Gabillard D, Moh R, Inwoley A, Sorho S, Eholié S, Rouet F, Seyler C, Danela C, Anglaret X. Anthropometric and immunological success of antiretroviral therapy and prediction of virological success in west African adults. *Bulletin of the World Health Organization*. June 2008, 86 (6): 435-442.

21 Hurley E, Coutsooudis A, Giddy J, Knight S E, Loots E, Esterhuizen T M. Weight evolution and perceptions of adults living with HIV following initiation of antiretroviral therapy in a South African urban setting. *SAMJ*. 2011; 101(9): 645-650.

22 Kadiané-Oussou NJ, Koné D, Aba YT, Yapo MT, Karidioula JM, Tiéoule SC, Kra O. Devenir à douze mois des patients infectés par le virus de l'immunodéficience 1 initiant le dolutégravir à Bouaké (Côte D'Ivoire) de 2020 à 2021. *Rev Mali Infect Microbiol* 2023, Vol 18 N°2

23 Venter WDF, Sokhela S, Simmons B, Moorhouse M, Fairlie L, Mashabane N, et al. Dolutegravir with emtricitabine and tenofovir alafenamide or tenofovir disoproxil fumarate versus efavirenz, emtricitabine, and tenofovir disoproxil fumarate for initial treatment of HIV1 infection (ADVANCE): week 96 results from a randomized, phase 3, non-inferiority trial. *Lancet HIV* 2020; 7: e666-676

24. Norwood J, Turner M, Bofill C, Rebeiro P, Shepherd B, Bebawy S, et al. Brief Report: Weight Gain in Persons With HIV Switched From Efavirenz-Based to Integrase Strand Transfer Inhibitor-Based Regimens. *J Acquir Immune Defic Syndr* 2017; 76: 527-31.

25. Masson E. EM-Consulte. [cited August 22, 2023]. Weight gain induced by dolutegravir in Cameroon: 96-week follow-up of the ANRS 12313 - NAMSAL trial. Available at: <https://www.emconsulte.com/article/1383214/prise-de-poids-induite-par-le-dolutegravir-au-came>

26 M.-A. Khuong-Josses, B. Frison, Q. Bougault, M. Poupard, N. Sayre, A. Boussairi. Does switching to dual therapy (dolutegravir + lamivudine) have an influence on the weight of patients on dolutegravir-based triple therapy? 21st National Infectiology Days / Medicine and infectious diseases 50 (2020) S31–S199

27. Dirajlal-Fargo S, Moser C, Rodriguez K, El Kamari V, Funderburg N, Bowman E, et al. Changes in the Fungal Marker β -D-Glucan After Antiretroviral Therapy and Association With Adiposity. *Open Forum Infectious Diseases* 2019; 6.

28 Venter WDF, Moorhouse M, Sokhela S, et al. Dolutegravir plus two different prodrugs of tenofovir to treat HIV. *N Engl J Med* 2019; 381: 803–15.

29 Walmsley SL, Antela A, Clumeck N, et al. Dolutegravir plus abacavir-lamivudine for the treatment of HIV-1 infection. *N Engl J Med* 2013; 369: 1807–

30. Kouanfack C, Mpoudi-Etame M, OmgbaBassega P, et al. Dolutegravir-based or low-dose efavirenz-based regimen for the treatment of HIV-1. *N Engl J Med* 2019; 381:816–26.

singl

31. Walmsley SL, Antela A, Clumeck N, et al. Dolutegravir plus abacavir–lamivudine for the treatment of HIV-1 infection. *N Engl J Med* 2013;369:1807-18.

32. Molina JM, Clotet B, van Lunzen J, et al. Once-daily dolutegravir versus darunavir plus ritonavir for treatment-naive adults with HIV-1 infection (FLAMINGO): 96 week results from a randomized, open-label, phase 3b study. *Lancet HIV* 2015;2(4): e127-e136.

33 Dorward J, Sookrajh Y, Khubone T, van der Molen J, Govender R, Phakathi S, Lewis L, Bottomley C, Maraj M, Lessells RJ, Naidoo K, Butler CC, Van Heerden R, Garrett N. Implementation and outcomes of dolutegravir-based first-line antiretroviral therapy for people with HIV in South Africa: a retrospective cohort study. *Lancet HIV*. 2023 May;10 (5):e284-e294

34 Batista CJB, Correa RG, Evangelista LR, et al. The Brazilian experience of implementing the active pharmacovigilance of dolutegravir. *Medicine (Baltimore)* 2019; 98:e14828.

35 Meireles MV, Pascom ARP, Duarte EC, McFarland W. Comparative effectiveness of first-line antiretroviral therapy: results from a large real-world cohort after the implementation of dolutegravir. *AIDS* 2019; 33:1663–68.

36 Schramm B, Temfack E, Descamps D, et al. Viral suppression and HIV-1 drug resistance 1 year after pragmatic transition to dolutegravir first-line therapy in Malawi: a prospective cohort study. *Lancet HIV* 2022; 9:e544–53.

37 Nabitaka VM, Nawaggi P, Campbell J, et al. High acceptability and viral suppression of patients on dolutegravir-based first-line regimens in pilot sites in Uganda: a mixed-methods prospective cohort study. *PLoS One* 2020; 15: e0232419.

38 Esber A, Dear N, Shah N, et al. Virologic impact of the dolutegravir transition: prospective results from the multinational African Cohort Study. J Acquir Immune Defic Syndr 2022; 91:285–89.

39 Walmsley SL, Antela A, Clumeck N, et al. Dolutegravir plus abacavir-lamivudine for the treatment of HIV-1 infection. N Engl J Med 2013; 369:1807–18.

40 Wainberg MA, Han YS. Will drug resistance against dolutegravir in initial therapy ever occur? Front Pharmacol 2015; 6:90

UNDER PEER REVIEW