

INFLUENCE OF BIOFERTIGATION AND CRUDE EDIBLE OILS APPLICATION ON GROWTH CHARACTERISTICS OF PAPAYA (*Carica papaya* L.) cv. RED LADY

ABSTRACT

An experiment was carried out to know the influence of biofertilization and crude edible oils application on growth, floral and fruit characteristics of papaya (*Carica Papaya* L.) cv. Red Lady was carried out at Department of Fruit Science, Dr. YSRHU-College of Horticulture, Venkataramanannagudem, West Godavari district of Andhra Pradesh during 2021-22 and 2022-23. The experiment was carried out in a Factorial RBD with three replications consisted of two factors viz., biofertilizers at three levels (B₁: 5 ml each of NFB + PSB + KRB; B₂: 2.5 ml each of NFB + PSB + KRB; B₃: Without biofertilizer) and crude edible oil at five levels (C₁: Ground nut oil @0.5%; C₂: Castor oil @0.5%; C₃: Mustard oil @0.5%; C₄: Ground nut oil + Castor oil + Mustard oil @0.5%; C₅: No crude oil spray) with fifteen treatment combinations. Among the treatments studied, the results revealed that the maximum plant height, stem girth, number of leaves per plant at 360 DAT were registered in the plants applied with 5 ml each of biofertilizers NFB+ PSB + KRB along with ground nut oil + castor oil + mustard oil @0.5% (B₁C₄).

Keywords: Papaya, Biofertilization, edible oils, NFB, PSB and KRB.

Introduction

Papaya (*Carica papaya* L.) is an evergreen herbaceous commercial fruit crop of tropical and subtropical regions and belongs to the family Caricaceae with chromosome number 2n=18. Papaya is a dicotyledon, polygamous and diploid species with geographical origin of Southern Mexico and Costa Rica (Candolle, 1884). Papaya is highly responsive to application of fertilizers. Though, chemical fertilizers fulfil the major requirement of the crop, their excessive and unbalanced use may lead to ecological hazards, depletion of physico-chemical properties.

Use of biofertilizers in papaya supplies plant nutrients to an optimum level besides maintenance of soil fertility for sustainable desired crop productivity and fruit quality through optimization of benefits from all possible sources in an integrated manner. Liquid biofertilizers have the ability to fix/solubilize the plant nutrients in soil through biological process. They are environmental friendly and play a significant role in crop production (Verma *et al.*, 2011).

N-fixing bacteria possess unique potential of fixing atmospheric nitrogen either by living symbiotically or non-symbiotically or to transform native soil nutrients from non-usable to usable form through biological process (Marwaha, 1995). Azotobacter have also been found to promote synthesis of growth promoting substances like auxins, gibberellins, cytokinins and antibiotic metabolites which in turn improved resistance against biotic and

abiotic stress (Awasthi *et al.*, 1998). Phosphate solubilizing bacteria play an important role in supplementing phosphorus to plants (Bhattacharya and Jain, 2000). The principal mechanism of mineral phosphate solubilization is the production of organic acids and acid phosphatases which play a major role in the mineralization of phosphorous in the soil (Shankar *et al.*, 2013). Most of the mineral potassium of the soil is unavailable for plant uptake (Sparks and Huang, 1985). Potassium releasing bacteria (KRB) mediate the releasing of potassium bearing minerals through production and excretion of organic acids (Ullaman *et al.*, 1996).

A lot of organic substances have been found effective in the maintenance of soil and plant health thus giving positive results for growth, yield and quality of different crops. Traditionally the organic edible oil is used as a fertilizer for farming to get high quality yields with low cost per crop (Vankineni and Muthusami, 2016). Recently farmers are adopting the practice of application of crude edible oils obtained from ground nut, castor and mustard seeds *etc.*, to get good growth and higher yield with better quality in some of the fruit crops. There is a need to analyse the effect of foliar application of crude edible oils on growth, yield and quality of papaya.

Materials and Methods

The current investigation, “Influence of biofertiligation and crude edible oils application on growth characteristics of papaya (*Carica Papaya* L.) cv. Red Lady was carried out at Department of Fruit Science, Dr. YSRHU-College of Horticulture, Venkataramanannagudem, West Godavari district of Andhra Pradesh during 2021-22 and 2022-23. The experiment was laid out in a Factorial Randomized Block Design (FRBD) with three replications consisted of two factors *viz.*, biofertilizers at three levels (B₁: 5 ml each of NFB + PSB + KRB; B₂: 2.5 ml each of NFB + PSB + KRB; B₃: Without biofertilizer) and crude edible oil at five levels (C₁: Ground nut oil @0.5%; C₂: Castor oil @0.5%; C₃: Mustard oil @0.5%; C₄: Ground nut oil + Castor oil + Mustard oil @0.5%; C₅: No crude oil spray) with fifteen treatment combinations.

First dose of NFB, PSB and KRB liquid biofertilizers were applied at 15 days after transplanting. Second and third doses of liquid biofertilizers were applied respectively at three months and six months after transplanting. Foliar sprayings of different concentrations of crude edible oils were taken up at three intervals on first month, four months and seven months after transplanting. The method of application of liquid biofertilizers was done with the help of drip system which was connected to small tank and drip liners and crude edible oils were applied to the plants by spraying.

Results and Discussion

GROWTH PARAMETER

Plant height (cm)

The influence of biofertiligation and foliar application of crude edible oils and their interaction on plant height was presented in table 1. Significant differences were noticed among the biofertiligation treatments at all the stages of observations recorded. The plant height was gradually increased from 90 to 360 DAT. The plant height was maximum (106.60, 159.53, 207.42 and 257.89 at 90, 180, 270 and 360 DAT respectively) in B₁ (5 ml

each of biofertilizers NFB (*Azotobacter chroococcum*) + PSB (*Bacillus megaterium*) + KRB (*Bacillus mucilaginosus*) followed by B₂ (2.5 ml each of NFB + PSB + KRB biofertilizers) (87.06, 145.93, 193.84 and 242.39 at 90, 180, 270 and 360 DAT respectively). The plant height was minimum (73.13, 119.23, 162.17 and 222.92 at 90, 180, 270 and 360 DAT respectively) in B₃ (without biofertilization).

The foliar application of crude edible oils had significantly influenced the plant height in papaya. The plant height was gradually increased from 90 to 360 DAT. The plant height was more (93.51, 145.38, 194.49 and 249.46 at 90, 180, 270 and 360 DAT respectively) in C₄ (ground nut oil + castor oil + mustard oil @0.5%) treatment, followed by C₁ (ground nut oil @0.5%) treatment at 90 DAT (90.22) and 180 DAT (143.36) and C₄ was on par with C₃ (mustard oil @0.5%) treatment at 270 DAT (190.47) and C₄ followed by C₃ at 360 DAT (242.82). The plant height was minimum (84.84, 136.15, 178.61 and 230.48 at 90, 180, 270 and 360 DAT respectively) in C₅ treatment (without crude edible oil spray).

The interaction between the biofertilization and crude edible oils spray treatments was found significant for plant height at all the stages of observations recorded except at 90 DAT. The interaction of biofertilization and crude edible oils spray (B₁C₄) (the combination of 5 ml each of biofertilizers NFB + PSB + KRB and 0.5% of ground nut oil + castor oil + mustard oil) recorded the maximum plant height (163.13, 215.43 and 268.53 at 180, 270 and 360 DAT respectively) followed by B₁C₁ (the combination of 5 ml each of biofertilizers NFB + PSB + KRB and 0.5% of ground nut oil) at 180 DAT (160.93) and B₁C₃ (the combination of 5 ml each of biofertilizers NFB + PSB + KRB and mustard oil @0.5%) at 270 DAT (210.54) and at 360 DAT (257.58). The plant height was minimum in B₃C₅ (without biofertilization and crude edible oils spray) at all stages of observations recorded (109.37, 145.52 and 209.48 at 180, 270 and 360 DAT respectively).

Though, plant height was genetically controlled trait, it could be manipulated with the application of biofertilizers and crude edible oils spray. The increase in the plant height with the application of biofertilizer along with crude edible oils might be due to better uptake and translocation of nitrogen to the growing points. The fact behind increase in plant height was due to application of nitrogen as it encourages vegetative growth through the formation of new cells, cell division, cell elongation and cell development (Azcon and Barea, 1975). The biofertilizers also help in fixing atmospheric nitrogen and also solubilization of phosphorous and potassium nutrients fixed to the soil particles by releasing organic acids through their metabolic process (Verma *et al.*, 2011). This resulted in vigorous growth of root system, which ultimately helped in better absorption of nutrients from soil and utilization of both nitrogen applied and fixed by NFB was reflected in terms of plant height. The results were in accordance with findings of Bharathi *et al.* (2021), Kumar *et al.* (2017) and Sourabh *et al.* (2018) in guava and Sharma *et al.* (2016) in mango cv. Amrapali, Amar *et al.* (2020), Divya and Bindu (2021) in papaya.

The crude edible oil sprays improved the plant height by supplying both primary and secondary macronutrients. As evident from the above the positive significant increase in plant height by the organic amendments used in the present investigation *i.e.* crude edible oils

reminds the fact that these substances are directly or indirectly associated with maintenance of plant health. Similar results were reported by Dayeswari *et al.* (2019) in Ber.

Stem girth (cm)

The influence of biofertiligation and foliar application of crude edible oils and their interaction on stem girth was presented in table 2. Significant differences were noticed among the biofertiligation treatments at all the stages of observations recorded. The stem girth was gradually increased from 90 DAT to 360 DAT. The stem girth was maximum (12.28, 21.95, 30.02 and 35.77 at 90, 180, 270 and 360 DAT respectively) in B₁ (5 ml each of biofertilizers NFB + PSB + KRB) followed by B₂ (2.5 ml each of NFB + PSB + KRB biofertilizers) (10.66, 19.34, 23.95 and 29.60 at 90, 180, 270 and 360 DAT respectively). The stem girth was minimum (8.52, 16.67, 20.33 and 26.59 at 90, 180, 270 and 360 DAT respectively) in B₃ (without biofertiligation).

The foliar application of crude edible oils had significantly influenced the stem girth in papaya. The stem girth was gradually increased from 90 DAT to 360 DAT. The stem girth was maximum (11.24, 20.25, 25.88 and 31.93 at 90, 180, 270 and 360 DAT respectively) in C₄ (ground nut oil + castor oil + mustard oil @0.5%) followed by C₁ (ground nut oil @0.5%) treatment at 90 DAT (10.63), 180 DAT (19.63) and 270 DAT (25.30) and C₂ (castor oil @0.5%) at 360 DAT (30.89). The stem girth was minimum (9.59, 18.12, 23.46 and 29.24 at 90, 180, 270 and 360 DAT respectively) in C₅ treatment (without crude edible oil spray).

The interaction between the biofertiligation and crude edible oils spray treatments was found significant for stem girth at all the stages of observations recorded except at 90 DAT and 360 DAT. The interaction of biofertiligation and crude edible oils spray *i.e.* B₁C₄ - the combination of 5 ml each of biofertilizers NFB + PSB + KRB and ground nut oil + castor oil + mustard oil @0.5% had recorded the maximum stem girth (23.06 and 31.36 at 180 and 270 DAT respectively) followed by B₁C₁ (The combination of 5 ml each of biofertilizers NFB + PSB + KRB and ground nut oil @0.5%) at 180 DAT (22.51) and 270 DAT (31.02). The stem girth was minimum in B₃C₅ (without biofertiligation and crude edible oils spray) at all stages of observations recorded (15.84 and 19.66 at 180 and 270 DAT respectively).

The stem girth is a genetically controlled trait, it could be manipulated with the application of growth-related compounds. The increase in stem girth of papaya plants was noticed with the combined application of biofertilizers and crude edible oils might be attributed to high rate of nitrogen mineralization through NFB, release of potassium nutrient through KRB microorganisms leads to continuous supply of available nutrients from organic and inorganic form improved the stem girth by activation of lateral meristems.

The present results were in consonance with findings of Suneetha and Ramachandrudu (2010) in oil palm; Kundu *et al.* (2011) in mango cv. Amarapali, Bhatnagar and Singh (2015) in custard apple cv. Arka Sahan; Amit *et al.* (2019) in guava, Dayeswari *et al.* (2019) in ber, Amar *et al.* (2020), Divya and Bindu (2021) in papaya.

Number of leaves per plant

The influence of biofertiligation and foliar application of crude edible oils and their interactions on number of leaves was presented in table 3. Significant differences were noticed among the biofertiligation treatments at all the stages of observations recorded. The number of leaves was gradually increased from 90 DAT to 360 DAT. The number of leaves was maximum (18.08, 25.14, 30.73 and 36.12 at 90, 180, 270 and 360 DAT respectively) in B₁ (5 ml each of biofertilizers NFB + PSB + KRB) followed by B₂ (2.5 ml each of NFB + PSB + KRB biofertilizers) (15.32, 19.94, 26.14 and 33.94 at 90, 180, 270 and 360 DAT respectively). The number of leaves was minimum (13.06, 16.51, 22.53 and 27.34 at 90, 180, 270 and 360 DAT respectively) in B₃ (without biofertiligation).

The foliar application of crude edible oils had significantly influenced the number of leaves in papaya. The number of leaves was gradually increased from 90 to 360 DAT. The number of leaves was highest (16.28, 22.24, 28.25 and 33.69 at 90, 180, 270 and 360 DAT respectively) in C₄ (ground nut oil + castor oil + mustard oil @0.5%) followed by C₁ (ground nut oil @0.5%) (15.81, 20.69, 26.58 and 33.11 at 90, 180, 270 and 360 DAT respectively). The number of leaves was minimum (14.60, 19.43, 25.12 and 30.95 at 90, 180, 270 and 360 DAT respectively) in C₅ (without crude edible oil spray).

The interaction between the biofertiligation and crude edible oils spray treatments was found significant for number of leaves at all the stages of observations recorded except at 90 DAT. The interaction of biofertiligation and crude edible oils spray (B₁C₄ - the combination of 5 ml each of biofertilizers NFB + PSB + KRB and ground nut oil + castor oil + mustard oil @0.5%) had recorded the maximum number of leaves (27.65, 32.26 and 37.35 at 180, 270 and 360 DAT respectively) followed by B₁C₃ treatment (the combination of 5 ml each of biofertilizers NFB + PSB + KRB and mustard oil @0.5%) at 180 DAT (25.78) and B₁C₁ treatment (the combination of 5 ml each of biofertilizers NFB + PSB + KRB and ground nut oil @0.5%) at 270 DAT (31.01) and B₁C₃ treatment (the combination of 5 ml each of biofertilizers NFB + PSB + KRB and mustard oil @0.5%) at 360 DAT (36.34). The number of leaves was minimum in B₃C₅ (without biofertiligation and crude edible oils spray) at all stages of observations recorded (15.78, 21.06 and 26.33 at 180, 270 and 360 DAT respectively).

Number of leaves is having direct correlation with photosynthetic efficiency of plants. The combined application of biofertilizers and crude edible oils had significantly increased the number of leaves per plant by continuous supply of water, hormones and nutrients to the meristematic tissue and consequently produced more number of leaves per plant as earlier reported by Sumithra *et al.* (2013) and Jain *et al.* (2014). The results are in close conformity with the findings of Shivakumar, 2010, Suresh, *et al.* (2010), Yadav, *et al.* (2011) and Tandel *et al.* (2014) in papaya, Sharma *et al.* (2009), Godage *et al.* (2013) in guava cv. Allahabad safeda, Kumar *et al.* (2017) in guava and Maskar (2018) in sapota.

CONCLUSION:

On the basis of results obtained in the present investigation, it could be concluded that, the biofertiligation with 5 ml each of biofertilizers NFB + PSB + KRB (B₁), spraying of

crude edible oils *viz.*, with ground nut oil + castor oil + mustard oil @0.5% (C₄) and their interaction *i.e.* plants applied with 5 ml each of biofertilizers NFB+ PSB + KRB along with ground nut oil + castor oil + mustard oil @0.5% (B₁C₄) performed best with respect to growth characteristics of papaya. Hence it could be concluded that biofertilizers and crude edible oil application were found more effective in maintaining the overall development of the papaya plant growth.

References :

- Amar Singh and Tripathi, V.K. 2020. Influence of INM on vegetative growth, fruiting, yield and soil physical characters in papaya (*Carica papaya* L.). *International Journal of Current Microbiology and Applied Sciences*. 9(10): 3811-22.
- Amit, K, Manoj, K.K, Rajesh, L, Mandloi, D.S. and Ashok, D. 2019. Impact of inorganic, organic and biofertilizers on growth and yield of guava (*Psidium guajava* L.) var. G-27 under gwalior agro-climatic condition of M.P. *Journal of Pharmacognosy and Phytochemistry*. 8(6): 420-24.
- Awasthi, R.P, Godara, R.K. and Kaith, N.S. 1998. Interaction effect of VA-mycorrhizae and Azotobacter inoculation on micronutrient uptake by peach seedlings. *Journal of Horticulture*. 11: 1-5.
- Azcon, R. and Barea, J.M. 1975. Synthesis of auxins, gibberellins and cytokinins by *Azotobacter vinlandi* and *Azotobacter beijerinckii* related to effects produced on tomato plants. *Plant and Soil*. 43(3): 609-19.
- Bharathi Nirujogi, Madhavi, M, Naram Naidu, L, Vinaya Kumara Reddy, P, Salomi Sunnetha and Rama Devi, P. 2021. Influence of liquid and carrier based biofertilizers on growth characters of guava (*Psidium guajava* L.) cv. Taiwan White. *The Pharma Innovation Journal*. 10(11): 604-06.
- Bhattacharya, P. and Jain, R.K. 2000. Phosphorous solubilizing biofertilizers in the whirl pool of rock phosphate-challenges and opportunities. *Fertilizer News*. 45: 45-52.
- Bhatnagar, P. and Singh, J. 2015. Response of custard apple cv. Arka Sahan plants to integrated nutrient management. *Horticulture Flora Research Spectrum*. 4(3): 204-08.
- Candolle, D.A. 1884. *Origin of Cultivated Plants*. John Wiley and Sons, New York, 281.
- Dayeswari, D. 2019. Influence of pruning systems and application of crude edible oils on growth, yield and quality of ber (*Ziziphus mauritiana* Lamk) cv. Apple ber. *Ph.D. Thesis*. Dr YSRHU, Andhra Pradesh, India.
- Divya Hari and Bindu, B. 2021. Effect of organic manures and biofertilizers on growth and yield attributes of papaya (*Carica papaya* L.). *Journal of Tropical Agriculture*. 59(1): 102-106.
- Godage, S.S, Parekh, N.S. and Nehete, D.S. 2013. Influence of biofertilizers and chemical fertilizers on growth, flowering and fruit characters of guava (*Psidium guajava* L.) cv. Allahabad safeda. *International Journal of Agricultural Sciences*. 9(1): 309-13.

- Jain, S.A, Jagtap, M.S. and Patel, K.P. 2014. Physico-Chemical Characterization of farmland Soil used in some villages of Lunawada Taluka, Dist: Mahisagar (Gujarat) India. *International Journal of Science and Research Publications*. 4(3): 1-5.
- Kumar, R.K, Jaganath, S. and Guruprasad, T.R. 2017. Impact of organic, inorganic and biofertilizers with different spacing on vegetative growth and yield of guava (cv. Lalit) during summer season. *International Journal of Pure and Applied Biological Sciences*. 5(1): 310-19.
- Kundu, S, Datta, P, Mishra, Rashmi, K. and Ghosh, B. 2011. Influence of biofertilizer and inorganic fertilizer in pruned mango orchard cv. Amrapali. *Journal of Crop and Weed*. 7(2): 100-03.
- Marwaha, B.C. 1995. Biofertilizers - A supplementary source of plant nutrient. *Fertilizer News*. 40: 39-50.
- Maskar, S.L. 2018. Studies on effect of bio-fertilizers and inorganic fertilizers on growth, yield and quality of sapota (*Manilkara Achras* (Mill.) Forseberg). cv. Kalipatti. *Ph.D. Thesis*. Vasantao Naik Marathwada Krishi Vidyapeeth, Parbhani, India.
- Shankar, T, Sivakumar, T, Asha, G, Sankaralingam S. and Sundaram, V. 2013. Effect of PSB on growth and development of chilli and maize plants. *World Applied Sciences Journal*. 26(5): 610-17.
- Sharma, A, Kher, R, Wali, V.K. and Baksh, P. 2009. Effect of biofertilizers and organic manures on physico-chemical characteristics and soil nutrient composition of guava (*Psidium guajava* L.) cv. Sardar. *Journal of Research*. 8(2): 150-156.
- Sharma, R, Jain, P.K. and Sharma, T.R. 2016. Effect of inorganic and organic sources of nutrients on physico-chemical composition of Mango (*Mangifera indica* L.) cv. Amrapali. *Economic Affairs*. 61(4): 677-82.
- Shivakumar, B.S. 2010. Integrated nutrient management studies in papaya (*Carica papaya* L.) cv. Surya. *Ph.D. Thesis*. University of Agricultural Science, Dharwad, Karnataka, India.
- Sourabh, J.R.S, Baloda, S, Kumar, R, Sheoran, V. and Saini, V.H. 2018. Response of organic amendments and biofertilizers on growth and yield of guava during rainy season. *Journal of Pharmacognosy and Phytochemistry*. 7(6): 2692-95.
- Sparks, D.L. and Huang, P.M. 1985. Physical chemistry of soil potassium. *Potassium in Agriculture*: 201-76.
- Sumithra, S, Ankalaiah, C, Rao, D. and Yamuna, R.T. 2013. A case study on physico-chemical characteristics of soil around industrial and agricultural area of yerraguntla, kadapa district, A.P, India. *International Journal of Geology, Earth and Environmental Sciences*. 3(2): 28-34.
- Suneetha V. and Ramachandrudu. 2010. Effect of biofertilizers on growth and vigour of Oil Palm seedlings. *International Journal of Oil Palm*. 7(2): 29-31.

- Suresh, C.P, Nath, S, Poduval, M. and Sen, S.K. 2010. Studies on the efficacy of phosphate solubilizing microbes and VAM fungi with graded levels of phosphorus on growth, yield and nutrient uptake of papaya (*Carica papaya* L.). *Acta Horticulturae*. 851: 401-06.
- Tandel, B.M, Patel, B.N. and Patel, B.B. 2014. Effect of Integrated Nutrient Management on Growth and Physiological Parameters on Papaya cv. Taiwan Red Lady. *Biosciences*. 2175.
- Ullaman, W.J, Kirchman, D.L. and Welch, W.A. 1996. Laboratory evidence by microbially mediated silicate mineral dissolution in nature. *Chemical Geology*. 132: 11-17.
- Vankineni, T. and Muthusami, R. 2016. Turn-key solutions for commercial agriculture projects. <https://www.v=zCcALjZm9do>.
- Verma, M, Sharma, S. and Prasad, R. 2011. Liquid Biofertilizers: Advantages over carrier based biofertilizers for sustainable crop production. *International Society of Environmental Botanists*. 17(2): 63-68.
- Yadav, P.K. Yadav, A.L. Yadav, A.S. and Yadav, H.C. 2011. Effect of integrated nutrient nourishment on vegetative growth and physico-chemical attributes of papaya (*Carica papaya* Linn.) fruit cv. Pusa Dwarf. *Plant Archives*. 11(1): 327-329.

Table 1. Effect of biofertiligation and foliar application of crude edible oils on plant height (cm) of papaya cv. Red Lady.

Crude edible oils (C)	Biofertilizers (B)															
	90 DAT			Mean of (C)	180 DAT			Mean of (C)	270 DAT			Mean of (C)	360 DAT			Mean of (C)
	B ₁	B ₂	B ₃		B ₁	B ₂	B ₃		B ₁	B ₂	B ₃		B ₁	B ₂	B ₃	
C ₁	108.49	88.14	74.03	90.22	160.93	146.80	122.34	143.36	206.85	195.64	165.61	189.37	255.63	241.28	227.00	241.30
C ₂	104.78	85.52	72.81	87.70	158.43	145.21	119.76	141.13	204.21	192.49	161.62	186.11	255.08	244.03	224.70	241.27
C ₃	106.04	86.22	72.86	88.37	158.62	146.16	120.60	141.79	210.54	193.33	167.54	190.47	257.58	244.81	226.07	242.82
C ₄	113.16	92.06	75.31	93.51	163.13	148.93	124.08	145.38	215.43	197.47	170.58	194.49	268.53	252.51	227.35	249.46
C ₅	100.51	83.36	70.65	84.84	156.54	142.55	109.37	136.15	200.05	190.26	145.52	178.61	252.65	229.31	209.48	230.48
Mean of (B)	106.60	87.06	73.13		159.53	145.93	119.23		207.42	193.84	162.17		257.89	242.39	222.92	
Factors	S Em±		CD at 5%		S Em±		CD at 5%		S Em±		CD at 5%		S Em±		CD at 5%	
B	0.48		1.39		0.12		0.35		1.19		3.44		0.11		0.32	
C	0.62		1.79		0.16		0.45		1.53		4.44		0.14		0.42	
B x C	1.07		NS		0.27		0.78		2.65		7.68		0.25		0.72	

B₁: 5 ml each of NFB + PSB + KRB

B₂: 2.5 ml each of NFB + PSB + KRB

B₃: Without biofertilizer

C₁: Ground nut oil@0.5%

C₂: Castor oil@0.5%

C₃: Mustard oil@0.5%

C₄: Ground nut oil + Castor oil + Mustard oil@0.5%

C₅: No

crude oil spray

Table 2. Effect of biofertiligation and foliar application of crude edible oils on stem girth (cm) of papaya cv. Red Lady.

Crude edible oils (C)	Biofertilizers (B)															
	90 DAT			Mean of (C)	180 DAT			Mean of (C)	270 DAT			Mean of (C)	360 DAT			Mean of (C)
	B ₁	B ₂	B ₃		B ₁	B ₂	B ₃		B ₁	B ₂	B ₃		B ₁	B ₂	B ₃	
C ₁	12.43	10.87	8.58	10.63	22.51	19.96	16.42	19.63	31.02	24.22	20.66	25.30	35.75	29.05	27.03	30.61
C ₂	12.10	10.86	8.44	10.47	22.39	18.80	16.70	19.30	30.26	23.84	20.25	24.78	36.63	29.86	26.18	30.89
C ₃	12.29	10.47	8.78	10.51	21.43	19.77	16.71	19.30	29.82	23.36	20.04	24.41	35.36	29.86	26.53	30.58
C ₄	12.90	11.59	9.25	11.24	23.06	20.02	17.68	20.25	31.36	25.26	21.03	25.88	37.62	30.52	27.66	31.93
C ₅	11.70	9.53	7.55	9.59	20.36	18.15	15.84	18.12	27.66	23.07	19.66	23.46	33.49	28.71	25.53	29.24
Mean of (B)	12.28	10.66	8.52		21.95	19.34	16.67		30.02	23.95	20.33		35.77	29.60	26.59	
Factors	S Em±		CD at 5%		S Em±		CD at 5%		S Em±		CD at 5%		S Em±		CD at 5%	
B	0.10		0.29		0.11		0.33		0.14		0.41		0.25		0.71	
C	0.13		0.38		0.15		0.42		0.18		0.53		0.32		0.92	
B x C	0.23		NS		0.25		0.73		0.32		0.92		0.55		NS	

B₁: 5 ml each of NFB + PSB + KRB

B₂: 2.5 ml each of NFB + PSB + KRB

B₃: Without biofertilizer

C₁: Ground nut oil@0.5%

C₂: Castor oil@0.5%

C₃: Mustard oil@0.5%

C₄: Ground nut oil + Castor oil + Mustard oil@0.5%

C₅: No

crude oil spray

Table 3. Effect of biofertiligation and foliar application of crude edible oils on number of leaves per plant of papaya cv. Red Lady.

Crude edible oils (C)	Biofertilizers (B)															
	90 DAT			Mean of (C)	180 DAT			Mean of (C)	270 DAT			Mean of (C)	360 DAT			Mean of (C)
	B ₁	B ₂	B ₃		B ₁	B ₂	B ₃		B ₁	B ₂	B ₃		B ₁	B ₂	B ₃	
C ₁	18.56	15.33	13.53	15.81	24.93	20.26	16.87	20.69	31.01	26.05	22.69	26.58	36.18	36.03	27.11	33.11
C ₂	17.75	15.27	12.72	15.25	23.84	19.04	16.04	19.64	30.43	25.22	22.48	26.04	36.22	33.42	27.02	32.22
C ₃	18.18	15.16	13.16	15.50	25.78	19.68	16.53	20.66	30.66	25.87	22.47	26.33	36.34	33.34	27.42	32.37
C ₄	19.06	15.90	13.88	16.28	27.65	21.74	17.33	22.24	32.26	28.53	23.95	28.25	37.35	34.89	28.82	33.69
C ₅	16.85	14.94	12.01	14.60	23.52	18.98	15.78	19.43	29.29	25.02	21.06	25.12	34.49	32.04	26.33	30.95
Mean of (B)	18.08	15.32	13.06		25.14	19.94	16.51		30.73	26.14	22.53		36.12	33.94	27.34	
Factors	S Em±		CD at 5%		S Em±		CD at 5%		S Em±		CD at 5%		S Em±		CD at 5%	
B	0.10		0.28		0.03		0.08		0.02		0.07		0.03		0.08	
C	0.12		0.36		0.03		0.10		0.03		0.09		0.04		0.10	
B x C	0.21		NS		0.06		0.17		0.06		0.16		0.06		0.18	

B₁: 5 ml each of NFB + PSB + KRB
biofertilizer

B₂: 2.5 ml each of NFB + PSB + KRB

B₃: Without

C₁: Ground nut oil@0.5% C₂: Castor oil@0.5% C₃: Mustard oil@0.5% C₄: Ground nut oil + Castor oil + Mustard oil@0.5% C₅: No crude oil spray

UNDER PEER REVIEW