

Genotypic Characterization and Virulence Features of Bacteria Associated with Poultry Meat

ABSTRACT

Microbial investigation of live poultry birds were carried out in three different locations (Aluu, Elioparanwo and RSU) in Ikwerre, Obio-Akpor and Port Harcourt Local Government Areas, Rivers State, Nigeria. This study was designed to identify bacteria from the muscle, breast, intestine, wing and leg of poultry birds using molecular techniques and evaluating their virulence features. The birds were processed and microbiological analysis was done using standard procedures, molecular identification of the isolates was carried out. The mean total heterotrophic bacterial count of the muscle, breast, intestine, wing and leg in the three different locations ranged from 16.6 ± 2.3 to 25.2 ± 3.5 , 1.6 ± 0.2 to 24.8 ± 8.6 , 0.11 ± 0.1 to 8.9 ± 2.7 , 1.1 ± 0.8 to 10.6 ± 1.8 and 3.4 ± 0.1 to 4.9 ± 0.9 CFU/g, respectively. The mean total coliform counts of the three locations ranged from 2.1 ± 0.3 to 2.1 ± 0.2 , 1.8 ± 0.2 to 0.9 ± 0.6 , and 0.32 ± 2.5 to 0.32 ± 0.25 , 2.6 ± 0.3 to 7.7 ± 0.9 and 2.3 ± 0.4 to 2.9 ± 0.3 CFU/g, respectively. Virulence features of *A. caviae* (CPO72326) shows proteolytic activity, DNase, motility and biofilm formation. *A. baumannii* (ON60630, OL455994) showed lipolytic activity, catalytic, starch hydrolysis and biofilm formation. *C. braakii* (ON557391) was positive for motility, catalytic and proteolytic. *C. freundii* (motile, catalytic, proteolytic and biofilm formation). *M. odoratimimus* (lipolytic, motile, catalytic, DNase, proteolytic, ureolytic, hemolytic and biofilm formation). The study revealed that microbial contamination of poultry meat could arise from microorganism present in the environment and on birds. These findings, thus underscored the urgent need for improved biosecurity measures, enhanced hygiene practices to safeguard both poultry health and public health. There should therefore be stringent implementation of food safety measure throughout the poultry value chain from farm to table.

Keywords: Poultry meat, Public health, Rivers State, Virulence.

1. INTRODUCTION

Poultry farming in Nigeria has now assumed a significant source of animal protein [1]. Household and commercial poultry production now complement imported poultry meat, although the former is more widespread due to cost [2]. Consumption of chicken overshadows other poultry meat in Nigeria[3]. Chicken provide good quality protein and lowfat poultry meat at lowercost; hence it is preferable. With high acceptance of chicken meat, due in part to it not having religious restriction, their demand is ever increasing [4]. Yet, poultry meat is seen as being expensive foran average Nigerian. In villages, eating of poultry meat is kept for important events, meat are gotten from household flocks. Those living in cities consume larger quality of poultry due to their relatively higher income and they have quick access to fresh or frozen products in markets and fastfood outlets. Poultry are used for all kinds of ceremonies/events such as social-cultural and religious events in Nigeria, causing demand for poultry meat to spike around Christmas, New year, and Easter [5].

As nutritious and healthy as poultry meat, its production and processing for consumption can introduce both pathogenic and spoilage microorganisms into them [6, 7]. Human illness may follow from handling of raw meat, undercooking or mishandling of the cooked product. Foodborne illnesses from poultry consumption are of serious public health concern [8]. These pathogens can come from farms, during transportation, slaughter and particularly when processing, although modern practices requires sterility of the final products [9]. The presence of few pathogens in uncooked meats raises no objection as they can be handled through cooking before ingestion.

Meat can be processed under hygienic conditions and properly stored yet not immune to contamination from natural microflora which maybe pathogenic. Pathogenic bacteria contaminate organic poultry as they do conventional [10], concluded whatever differences are observed between organic and conventional might come from choice of breed, which ultimately embroil meat quality such as appearance and nutritive value. Since chicken protein is a good media supporting microbial growth, any unsanitary condition during the rearing, processing and retailing of poultry meat which would have an effecton the bacterial load of the poultry meat. The presence of pathogens of any kind in food is undesirable, hence the setting up of standards. It is in the public interest that food sold in the open market be regularly monitored for presence of pathogens like *Myroidesspp*, *Enterobacter spp*, *Citrobacter spp*, *Acinetobacter spp* and *Aeromonas spp*.

In recent years, the health benefits of meat as a regular part of the human diet is offset by risks. Meat, like any food, transmit certain diseases [11], but complete cooking and avoiding recontamination reduces this possibility. According to [12], food-borne disease is any illness that results from eating contaminated food or meats. Food- borne diseases associated with the consumption of poultry meat and its processed products are of public health significance worldwide [13]. It is caused by pathogenic bacteria, viruses, parasites that contaminate food, as well as chemical or toxins. Ensuring safe food supply has been one of the major challenges and concerns for producers, consumers and public health officials in both developing and developed countries. This is because foods excessively contaminated with pathogenic and spoilage microorganisms are undesirable and can cause food borne illnesses [14, 15]. Such

illnesses cost billions of dollars in medical care and sometimes even result in death. Several epidemiological reports have implicated foods from animal origin as major vehicles associated with illnesses caused by food borne pathogens [14, 15]. Therefore, this study was aimed to determine virulence features and molecularly characterize bacterial isolates from poultry meat.

2. MATERIALS AND METHODS

2.1. Description of Study Area

The study was conducted in three Local Government Areas (ObioAkpok, Port Harcourt and Ikwerre Local Government Area) in Rivers State Nigeria. Three poultry farms were chosen for this study Aluu, Elioparanwo and Rivers State University demonstration farm. Samples were collected trimonthly (Quarterly) for a period of one year from each of the farms.

2.2. Collection of Samples

A total of sixty-five (125) samples of Muscle, Breast, Wing, Intestine and Leg, were obtained from the three different farms namely; Aluu, Elioparanwo and RSU Farm and transported aseptically to the Department of Microbiology Laboratory, Rivers State University for bacteriological analysis.

2.3. Preparation of Samples

The meat samples were prepared for bacteriological analysis as described by [16]. The muscle, breast, wing, intestine and leg were collected with the aid of a sterile knife.

2.4. Enumeration of Bacterial Isolates

Collected samples were analysed using standard microbiological method. Samples were streaked on Nutrient, MacConkey, and Eosin methylene blue agar plate. Ten grams (10g) of each chicken parts were weighed and transferred into different conical flask containing 90ml of sterilized normal saline. The flasks were swirled to dislodge the microbial contents under study [17]. Subsequent 10-fold serial dilution were carried out by transferring 1 ml into test tubes containing 9ml sterile normal saline until 10^{-6} dilution. An aliquot (0.1ml) was collected from the test tube with a dilution factor of 10^{-3} and 10^{-6} using a pipette and inoculated on dried medium. Using a spreader, the aliquot was evenly spread across the medium.

The inoculated plates were inverted and incubated at 37°C for 24 hours after which plates were counted and recorded. Pure bacteria isolates were suspended in 10% (v/v) glycerol at -4°C [18].

2.5. Characterisation of Bacterial Isolates by Conventional and Molecular methods

The bacterial isolates were characterised based on their colonial/morphological and biochemical characteristics [19].

Isolates were also subjected to analytical processes using simple molecular tools such as polymerase chain reaction (PCR) and genome sequencing approach to classify and identify the isolates from their Deoxyribonucleic acid (DNA) [20].

2.6. Test for Virulence

2.6.1. Biofilm Formation Assay

This test was done to ascertain the ability of the known bacteria to produce biofilms. The possibility of the biofilm development from the recovered bacteria was determined using the Congo red agar method as adopted by [21], was employed in this study as a simple qualitative and quantitative analysis that involved the use of Brain Heart Infusion agar supplemented with agar-agar, sucrose and Congo red stain.

2.6.2. DNase Assay

The DNase test was done to determine the ability of the bacterial isolates to hydrolyze Deoxyribonucleic acid and utilize it as a source of carbon and energy for growth. DNase is an extracellular enzyme that is secreted outside a bacterial cell membrane. When the enzyme is expressed, it is an indicative of a virulent property of bacteria which enables it to cause disease. The test procedure was strictly carried out as described by [22]. In sorting this feature, the bacterial isolates were streaked on a freshly prepared (DNase) medium and incubated at 37°C for 24 hours. After incubation, growth colonies were observed, and the plates were then flooded with diluted hydrochloric acid. DNase positive cultures showed a distinct clear zone around the streak while negative cultures showed no clear zone.

2.6.3. Lipase Test

The investigation was done to know if the bacterial isolates can breakdown the lipids of its host. This breakdown is made possible by lipase which the bacterium possesses, and from the breakdown process, the bacterium is able to derive carbon as its source of energy for growth and survival. Thus, when the bacterium survives, its harmfulness is expressed. This assay was carried out according to [23]. The bacterial isolates were

streaked onto a freshly prepared lipid agar medium and for 24 hours at 37°C. After incubation, growth was observed and a blue spirit indicator chemical was used to flood the medium, and observed within seconds. The blue spirit indicator, on observation, was deep blue in appearance which later changed to a light blue opaque medium, signifying a positive result, while in the absence of a change to a light blue opaque medium it indicated a negative result.

2.6.4. Catalase Activity

The assay was to investigate the production of catalase (an enzymes that speeds the release of oxygen from hydrogen peroxide) in the test organism. Catalase production is a virulent determinant property, which a bacterium elaborates to survive the effect of hydrogen peroxide released by phagocytes. The procedure employed to express this activity in bacteria was according to [24], and involves transferring a small fraction of bacterial isolate onto a clean microscope slide and adding a loopful of hydrogen peroxide. After a few seconds, where gas bubble ensued it indicated positive result while absence of gas bubble indicated negative result.

2.6.5. Motility Test

This test was done to spot the presence of flagella in bacteria. More recently, it has become evident that flagellum, a structure for movement in some bacteria also have other biological functions [25]. Flagella has been reported to function as adhesions during bacterial invasion of a host cell; an action which signifies its virulence. The procedure for motility test as done by [25], required the preparation of emulsified nutrient medium, followed by pouring into a test tube and then with the aid of a sterile

needle, the isolate is inoculated in the medium by piercing. The tube was then incubated for 24 hours at 37°C. A diffused growth spread within the tube and not along the stab line indicated motility/virulence of the bacterial isolate.

2.6.6. Capsule Staining

The presence of a capsule in a bacterium is implicative of its virulence because of the capsule can impede phagocytotic response of the host cell. Capsule staining test was done to determine the presence of capsule in the bacterium. The procedure was carried by adding a few drops of crystal violet onto the test bacteria on a clean microscope slide, then stirred and viewed under a light microscope. A light blue appearance signified encapsulated cell, while the reverse signified that the cell was not capsulated [26].

2.6.7. Hemolytic Activity

Hemolytic activity of the isolates was investigated. This was to figure out if the bacterial isolates have the ability to digest red blood cells. Hemolytic activity was determined by streaking the isolates onto freshly prepared blood agar medium as described by [24], and incubated at 37°C for 24-48 hours. Positive cultures showed a distinct clear zone around the streak showing the ability of the bacteria to produce toxin while negative result revealed no clear zone around the streak.

2.6.8. Ureolytic Activity

Urease is a virulence property found in some bacteria, and its presence in a bacterium aid it in colonizing its host organism with a toxic effect on the host. The ureolytic test is aimed at determining the ability of the bacteria to produce urease. The urease so

produced, reduces urea to ammonia that is toxic to the host. The procedures for the test according to [24] involved inoculation of the test bacteria into urease broth medium, followed by incubation at 35°C for 18-24 hours. A positive result indicated by change of the medium colour from red to pink while a negative result was indicated by retention of the original colour of red.

2.6.9. Proteinase Activity

Protease activity of the bacterial isolate was screened using casein agar. In screening, the isolates were streaked onto casein agar medium, followed by incubation at 37°C for 48 hours. Positive plates were identified with zones of proteolysis, in which the casein is hydrolysed leaving a clear zone around the growth area of the isolates and negative plates observed without clear zone on the plate [27].

2.6.10. Starch Hydrolysis

The assay was aimed at observing if the bacterial isolate can breakdown starch in the host by the production of amylase enzyme. The procedure for the test required preparing a starch agar medium and streaking of the isolate onto it, followed by inoculation at 35°C for 48 hours. The process was then followed by flooding the agar plate with Gram's iodine. A purple-blue colouration around the organism signified the presence of amylase while an absence signified no amylase [28].

2.7. Statistical Analysis

The enumerated colonies were calculated using descriptive statistics. One-way and Two-way ANOVA was used to check for significant differences. A p-value of less than 0.05 was considered to indicate a statistically significant difference.

3. RESULTS

3.1. Total Heterotrophic Bacterial Load of Chicken parts for the three Sampled farms

Results of the total heterotrophic bacterial load of chicken parts for the three farms are presented in Table 1. Results showed that the mean range of the total heterotrophic bacterial counts for Breast, Wing, Leg, Intestine and Muscle (thigh) of the farms were 16.6 ± 2.3 to 25.2 ± 3.5 , 1.6 ± 0.2 to 24.8 ± 8.6 , 0.11 ± 0.1 to 8.9 ± 2.7 , 1.1 ± 0.8 to 10.6 ± 1.8 and 3.4 ± 0.1 to 4.9 ± 0.9 CFU/g, respectively.

More so, results showed that there were significant differences across the poultry parts surfaces in the respective farms and across the farms. Poultry parts such as the intestines, breast and wings of samples obtained from the RSU were significantly higher than those recorded in Aluu and Elioparanwo poultry parts.

Table 1: Mean Total Heterotrophic Bacterial Counts from Chicken Parts from the three Sampled Farms

Samples	Aluu ($\times 10^6$)	Elioparanwo ($\times 10^6$)	RSU ($\times 10^6$)	P-value
Breast (CFU/g)	16.6 \pm 2.3 ^b	9.9 \pm 7.0 ^a	25.2 \pm 3.5 ^b	<0.01
Wing (CFU/g)	1.6 \pm 0.2 ^a	20.1 \pm 1.5 ^b	24.8 \pm 8.6 ^b	0.004
Leg (CFU/g)	0.11 \pm 0.1 ^a	23.3 \pm 1.8 ^c	8.9 \pm 2.7 ^b	0.02
Intestine (CFU/g)	1.1 \pm 0.8 ^a	0.59 \pm 0.8 ^a	10.6 \pm 1.8 ^b	<0.001
Muscle (thigh) (CFU/g)	3.4 \pm 0.1 ^a	3.2 \pm 0.1 ^a	4.9 \pm 0.9 ^a	0.182

3.2. The Mean Coliform Counts for Chicken Parts from the three Sampled Farms

Results of the mean coliform counts for chicken parts from the three sampled farms is presented in Table 2. Results showed that the mean coliform counts of the chicken parts in Aluu, Elioparanwo and RSU farms ranged from 2.1 \pm 0.3 to 2.1 \pm 0.2, 1.8 \pm 0.2 to 0.9 \pm 0.6, 0.32 \pm 2.5 to 0.32 \pm 0.25, 2.6 \pm 0.3 to 7.7 \pm 0.9 and 2.3 \pm 0.4 to 2.9 \pm 0.3 CFU/g, respectively.

Table 2: Mean Total Coliform Counts for Chicken Parts from the three Sampled Farms

Samples	Aluu ($\times 10^3$)	Elioparanwo ($\times 10^3$)	RSU ($\times 10^3$)	P-value
Breast (CFU/g)	2.1 \pm 0.3 ^a	1.8 \pm 0.2 ^a	2.1 \pm 0.2 ^a	0.331
Wing (CFU/g)	1.8 \pm 0.2 ^a	1.8 \pm 0.2 ^a	0.97 \pm 0.6 ^a	0.331
Leg (CFU/g)	0.32 \pm 2.5 ^a	1.8 \pm 0.2 ^a	0.32 \pm 0.25 ^a	0.611
Intestine (CFU/g)	2.6 \pm 0.3 ^a	7.7 \pm 0.9 ^b	7.7 \pm 0.9 ^b	0.022
Muscle (thigh) (CFU/g)	2.3 \pm 0.4 ^a	2.1 \pm 0.9 ^a	2.9 \pm 0.3 ^a	0.441

UNDER PEER REVIEW




Fig. 1: Phylogenetic Tree showing the Evolutionary Distance between the Bacterial Isolates

3.3. Phylogenetic Analysis

The obtained 16s rRNA sequence from the isolate produced an exact match during the megablast search for highly similar sequences from the NCBI non-redundant nucleotide

(nr/nt) database. The 16S rRNA of the isolates showed a percentage similarity to other species at 100%. The evolutionary distances computed using the Jukes-Cantor method were in agreement with the phylogenetic placement of the 16S rRNA of the isolates and revealed close relatedness to *Citrobacter braaki*, *Citrobacter freundii* and *Enterobacter hormaechei* (Fig.1).

3.4. Agarose Gel Electrophoresis

The Agarose gel electrophoresis showing the plasmid bands. Lane 1-10 showing the 16S rRNA bands at 1500bp while lane L represents the 500bp molecular ladder in Plate 1

1 2 3 4 5 L 6 7 8 9 10




Plate 1: Amplified 16S rRNA gene bands at 1500bp on Agarose gel after electrophoresis

3.4. Results of the virulence features from Aluu Farm

Results of the virulence features of *A. cavie*, *A. baumanii*, *C. braakii*, *C. freundii*, *E. hormaechei* and *M. odoratiminus* from Aluu farm is presented in Table 3. Results showed

that the percentage of *A. baumannii* and *M. odoratimimus* to lipolytic activity was 100% while other isolates showed no lipolytic activity. *A. caviae*, *C. braakii*, *C. freundii*, *E. hormaechei* and *M. odoratimimus* presented in Table 3. Results showed that the percentage of *A. caviae*, *C. braakii*, *C. freundii*, *E. hormaechei* and *M. odoratimimus* were 100% motile. The possession of DNase by bacterial isolates showed that *A. caviae* and *M. odoratimimus* were 100% positive for DNase enzyme while *A. baumannii*, *C. braakii*, *C. freundii*, and *E. hormaechei* lack the DNase enzyme. The ability to hydrolyze starch by the bacterial isolates showed that only *A. caviae*, *A. baumannii* and *E. hormaechei* isolates were able to hydrolyze starch while for proteolytic activity, isolates of *A. caviae*, *C. braakii*, *C. freundii* and *M. odoratimimus* were the only isolates with proteolase enzyme. In the virulence test for ureolytic and haemolysin activity, only the isolates of *M. odoratimimus* possessed the enzymes. The ability to produce biofilm was also investigated and only isolates of *A. caviae*, *A. baumannii*, *C. freundii*, *E. hormaechei* and *M. odoratimimus* formed biofilm while *C. braakii* and *M. odoratimimus* are the only isolates that possessed capsule.

3.5. Results of Virulence Features from Elioparanwo Farm

Results of the virulence features of *A. caviae*, *A. baumannii*, *C. braakii*, *C. freundii*, *E. hormaechei* and *M. odoratimimus* from Elioparanwo farm is presented in Table 4. Results was similar to the virulence attributes of similar isolates reported in the Aluu farm. The exception was just the number of isolates which was higher than those reported in the Aluu farm. For instance, *A. baumannii* and *M. odoratimimus* to lipolytic activity was 100% while other isolates showed no lipolytic activity. *A. caviae*, *C. braakii*, *C. freundii*, *E. hormaechei* and *M. odoratimimus* presented in Table 4. Results showed that the

percentage of *A. caviae*, *C. braakii*, *C. freundii*, *E. hormaechei* and *M. odoratimimus* were 100% motile. The possession of DNase by bacterial isolates showed that *A. caviae* and *M. odoratimimus* were 100% positive for DNase enzyme while *A. baumannii*, *C. braakii*, *C. freundii*, and *E. hormaechei* lack the DNase enzyme. The ability to hydrolyze starch by the bacterial isolates showed that only *A. caviae*, *A. baumannii* and *E. hormaechei* isolates were able to hydrolyze starch while for proteolytic activity, isolates of *A. caviae*, *C. braakii*, *C. freundii* and *M. odoratimimus* were the only isolates with proteolase enzyme. In the virulence test for ureolytic and haemolysin activity, only the isolates of *M. odoratimimus* possessed the enzymes.

3.6. Results of Virulence Features from RSU Farm

Results of the virulence features of *A. caviae*, *A. baumannii*, *C. braakii*, *C. freundii*, *E. hormaechei* and *M. odoratimimus* from RSU farm is presented in Table 5. Similar to the results in Table 3 and 4, *A. caviae*, *A. baumannii*, *C. braakii*, *C. freundii*, *E. hormaechei* and *M. odoratimimus* produced virulent attributes including the possession of biofilm and capsules.

Table 3: Virulence Features of Bacterial Isolated from Aluu Farm

Bacteria	No. of Isolates	Lipolytic Activity n (%)	Motility n (%)	Catalytic Activity n (%)	DNase n (%)	Starch hydrolysis n (%)	Proteolytic n (%)	Ureolytic n (%)	Hemol n (%)	Biofilm n (%)	Capsule formation n (%)
----------	-----------------	--------------------------	----------------	--------------------------	-------------	-------------------------	-------------------	-----------------	-------------	---------------	-------------------------

Key: n (%) =Number of bacteria positive (percentage positive), *A. caviae*=*Aeromonas caviae*, *A. baumannii*=*Acinetobacter baumannii*, *C. braakii*=*Citrobacter braakii*, *C. freundii*=*Citrobacter freundii*, *E. hormaechei*=*Enterobacter hormaechei*, *M. odoratimimus*=*Myroides odoratimimus*, Hemol=Hemolysin.

<i>A. cavie</i>	41	0 (0)	41 (100)	30 (73.2)	41 (100)	41 (100)	33 (80.5)	0 (0)	0 (0)	41 (100)	0 (0)
<i>A. baumannii</i>	40	40 (100)	0 (0)	40 (100)	0 (0)	34(85)	0 (0)	0 (0)	0 (0)	32 (80)	40 (100)
<i>C.braakii</i>	15	0 (0)	15 (100)	10 (66.7)	0 (0)	0 (0)	15 (100)	0 (0)	0 (0)	0 (0)	0 (0)
<i>C. freundii</i>	22	0 (0)	22 (100)	18 (81.8)	0 (0)	0 (0)	17 (77.3)	0 (0)	0 (0)	22 (100)	0 (0)
<i>E. hormaechei</i>	44	0 (0)	44 (100)	0 (0)	0 (0)	40 (90.9)	0 (0)	0 (0)	0 (0)	38 (86.4)	0 (0)
<i>M. odoratimimus</i>	21	21 (100)	21 (100)	15(71.4)	21 (100)	0 (0)	21 (100)	21 (100)	21 (100)	21 (100)	21 (100)
Total	183										

Table 4: Virulence Features of Bacterial Isolated from Elioparanwo Farm

Bacteria	No. of Isolates	Lipolytic Activity n (%)	Motility n (%)	Catalytic Activity n (%)	DNase n (%)	Starch hydrolysis n (%)	Proteolytic n (%)	Ureolytic n (%)	Hemol n (%)	Biofilm n (%)	Capsule formation n (%)
<i>A.cavie</i>	53	0 (0)	53 (100)	40(75.5)	53 (100)	43 (81.1)	40 (75.5)	0 (0)	0 (0)	53 (100)	0 (0)
<i>A.baumannii</i>	42	42 (100)	0 (0)	42 (100)	0 (0)	42 (100)	0 (0)	0 (0)	0 (0)	21 (50)	42 (100)
<i>C.braakii</i>	28	0 (0)	28 (100)	20 (71.4)	0 (0)	0 (0)	22 (78.6)	0 (0)	0 (0)	0 (0)	0 (0)
<i>C. freundii</i>	33	0 (0)	33 (100)	33 (100)	0 (0)	0 (0)	25 (75.8)	0 (0)	0 (0)	28 (84.8)	0 (0)

<i>E. hormaechei</i>	71	0 (0)	71 (100)	0 (0)	0 (0)	62 (87.3)	0 (0)	0 (0)	0 (0)	60(84.5)	0 (0)
<i>M. odoratimimus</i>	20	20 (100)	20 (100)	20 (100)	20 (100)	0 (0)	20 (100)	20 (100)	10 (50)	20 (100)	20 (100)
Total	247				(100)						

Key: n (%) =Number of bacteria positive (percentage positive), *A. caviae*=*Aeromonascaviae*, *A. baumanii*=*Acinetobacter baumanii*, *C. braakii*=*Citrobacter braakii*, *C. freundii*=*Citrobacter freundii*, *E. hormaechei*=*Enterobacter hormaechei*, *M. odoratimimus*=*Myroidesodoratimumus*, Hemol=Homolysin.

Table 5: Virulence Features of Bacterial Isolated from RSU Farm

Bacteria	No. of Isolates	Lipolytic Activity n (%)	Motility n (%)	Catalytic Activity n (%)	DNase n (%)	Starch hydrolysis n (%)	Proteolytic n (%)	Ureolytic n (%)	Hemol n (%)	Biofilm n (%)	Capsule formation n (%)
<i>A. caviae</i>	29	0 (0)	29 (100)	29 (100)	29 (100)	17(58.6)	20 (68.9)	0 (0)	0 (0)	29 (100)	0 (0)
<i>A. baumanii</i>	29	29 (100)	0 (0)	22(75.9)	0 (0)	27 (93.1)	0 (0)	0 (0)	0 (0)	17 (58.6)	29 (100)
<i>C. braakii</i>	17	0 (0)	17 (100)	17 (100)	0 (0)	0 (0)	17 (100)	0 (0)	0 (0)	0 (0)	0 (0)
<i>C. freundii</i>	23	0 (0)	23 (100)	20 (86.9)	0 (0)	0 (0)	21 (91.3)	0 (0)	0 (0)	18	0 (0)

										(78.3)	
<i>E. hormaechei</i>	42	0 (0)	42 (100)	0 (0)	0 (0)	21 (50)	0 (0)	0 (0)	0 (0)	21 (50)	0 (0)
<i>M. odoratimimus</i>	29	29 (100)	29 (100)	27 (93.1)	29 (100)	0 (0)	29 (100)	29 (100)	29 (100)	29 (100)	29 (100)
Total	169										

Key: n (%) =Number of bacteria positive (percentage positive), *A. caviae*=*Aeromonascaviae*, *A. baumannii*=*Acinetobacter baumannii*, *C. braakii*=*Citrobacter braakii*, *C. freundii*=*Citrobacter freundii*, *E. hormaechei*=*Enterobacter hormaechei*, *M. odoratimimus*=*Myroidesodoratimumus*, Hemol=Hemolysin.

UNDER PEER REVIEW

4. DISCUSSION

4.1. Bacterial Isolates Associated with Poultry

Bacterial associated with poultry meat in Rivers State was investigated. The world's consumption of poultry meat is rising significantly; according to the most recent figures, it reached 14.2 kg per person per year [29]. Therefore, in light of the rising consumption and production of poultry meat, assuring the microbiological safety of poultry meat products is a crucial concern. In actuality, poultry parts like the legs, intestines, muscles, wings and breasts including processed meat products are contaminated during and after slaughter by the germs from animal microbiota, the slaughterhouse environment, and the equipment employed [29]. Also, microbial contamination of poultry meat could arise from the type of microorganisms present in that environment as well as from the microorganisms on the bird (which could be the normal or transient flora) that gets into the poultry meat via openings or cuts either during killing or processing [30]. This agreed with the present study which showed uneven distribution of enteric bacterial isolates across the various parts of the poultry meat. More so, during the preparation and storage of food, several of these bacterial contamination might proliferate or persist [29]. Cross contamination of poultry meat is also very possible especially during scalding and defeathering of the poultry [31]. The muscles of healthy live birds are considered to be sterile while the digestive tract, the lungs, feathers and skin are considered to accommodate microbial diversities [30]. Thus, the presence of bacterial isolates in the muscles could be attributed to cross contamination during processing of the poultry meats. It has been noted that microorganisms from the environment as well as the microbiota of the poultry could contaminate the poultry meat [32, 29]. More so, it has been noted that while bacteria primarily come from the exterior of the animal and its

intestinal tract during slaughter, dressing, and cutting, more are introduced via knives, clothes, the air, workers, carts, crates, and equipment in general [33].

4.2. Virulence of Isolates

The virulence of the bacterial isolates which were evaluated showed that all bacterial isolate possessed certain degree of virulence. In particular, DNases may play a role in bacterial proliferation, biofilm formation, and the capacity of bacteria to evade the immune system [34]. DNases are enzymes that hydrolyze nucleic acids to produce oligonucleotides [35]. Dnase could therefore suggest these isolates' capacity to evade the immune system's activity, spreading and producing disease within the host. DNases are frequently cited as contributing to the pathogenicity of *Staphylococci* or *Streptococci* [21]. Indeed, studies have revealed that DNase can aid bacteria in eluding the structures that neutrophils create called extracellular traps (NETs) in order to capture and destroy pathogens [36]. The pathogenicity of most diseases varies, and virulence is a parameter that effectively separates pathogenic from non-pathogenic strains [37].

The presence of pili, capsular polysaccharide, DNase, coagulase, fibrinolysin, proteolytic, haemolysin, bacteriocin production, haemagglutination, serum sensitivity, attachment of epithelial cells, hydrophobicity, lipase, antiphagocytic factor (coagulase), biofilm, extracellular enzyme production, presence of surface layer, lysine decarboxy[38]. Accordingly, the existence of these virulent characteristics (Dnase, lecithin, motility, amylase, haemolysin and capsule) in the bacterial isolates of the present study could imply that they are pathogenic and could cause infections or disorders when given the proper infective dose. According to[39], pathogenic bacteria use many strategies to infect human hosts and cause sickness. They also release a

variety of compounds that bind to host cell targets and enable a range of host responses. According to [40], the formation of biofilms is a crucial component of virulence because it promotes bacterial colonisation by facilitating cell adhesion to epithelial cells and intestinal villi, decreasing bacterial susceptibility to antibiotics, and lessening immune system recognition of the bacteria.

5. CONCLUSION

The survey of microorganism associated with poultry in three local government areas in Rivers State has provided valuable insights into the potential risks associated with these bacteria in the poultry farming ecosystem. The following conclusions can be drawn from the study:

The presence of pathogens in poultry raises concerns about food safety and potential zoonotic transmission to humans. Proper cooking and handling of poultry products, as well as strict adherence to hygiene measures in processing facilities, are essential to reduce the risk of foodborne illnesses.

The study underscores the importance of implementing improved management practices in poultry farming, including enhanced biosecurity, waste management, and water source protection. These measures can help reduce the prevalence and transmission of bacteria, ultimately improving the health and productivity of poultry.

REFERENCES

1. Folorunsho OR. and Onibi GE. Assessment of the nutritional quality of eviscerated waste from selected chicken types.2005; Pp 300.
2. Amadi EN, Kiin KD, Kpormon LB and Robinson VKK. Microbial flora and nutritional composition of adult palm-winebeetle. International Journal of Current Microbiology and Applied Science,2014; 3(11):189-192.
3. Olabode HOK, Eghatona NO and Iyoha H. A retrospective (2004-2006) study of poultry diseases diagnosed at Benin, Edo State. Nigerian Veterinary Journal, 2008; 29(1):76-80.
4. Geidam YA, Bukar MM. and Ambali AG. Chick quality control: a key to sustainable poultry production in Nigeria. Nigerian Veterinary journal,2006; 27(2):1-6.
5. Omorodion NJP. and Odu NN. Occurrence of *Listeria* and *Salmonella* spp in chicken meats from different poultry production system. Research Journal of Pure Science and Technology.2016b; 4
6. Mead GC. Poultry meat processing and quality. Cambridge, UK, Woodhead, p.283-303. Microbiological quality of poultry meat; a review. Brazilian Journal of Poultry Science, 2004; 6, 3.
7. Kabour GA. Evaluation of microbial contamination of chicken carcasses during processing in Khartoum State M.V.SC thesis Sudan.2011; University of Science and Technology Sudan.
8. Myint MS. Epidemiology of *Salmonella* contamination of poultry meat product: Knowledge gaps in the farm to store products. University of Maryland,2014; pp14-21.
9. Jay JM. Modern food microbiology.4th edition CBS publishers and distributors, New Delhi, India,2005; ISBN81-239-0475-4.
10. Galgano F, Roberta T, Maria AC, Teresa S. and Marisa CC. Conventional and organic foods: A comparison focused on animal products. Cogent Food and Agriculture, 2016: 2, 1142918.
11. Iman M, Selma O, Ahmed and Sabiel YA. Microbial quality of frozen chicken meat. Journal of Applied and Industrial Sciences, 2015; 2,107-173.
12. Gayathri T. and Anu SA. Isolation and Characterisation of Microorganisms from raw meat obtained from different market places in and around Chennai, Journal of Pharmaceutical, Chemical and Biological Sciences, 2015;3(2): 295-301.
13. Keener KM, Bashor MP, Curtis PA, Sheldon BW. and Kathariou S. Comprehensive review of *Campylobacter* and poultry processing. Comprehensive Revision. *Food Science Safety*.2004; 3,105-116.
14. Odetunde SK, Lawal AK, Akolade MA. and Bak'ry SB. Microbial flora of frozen chicken part varieties. International Research Journal of Microbiology (IRJM)2011; 2(11):423-427.

15. Scorum H. and Sunde M. Resistance to antibiotics in the normal flora of animals, *Veterinary Research*, 2001; 32 (3-4): 227-41.
16. Danba EP, Bichi AH, Ishaku S, Ahmad MK, Buba U, Bingari MS, Barau BW. and Fidelis UF. Occurrence of pathogenic bacteria associated with *Clarias gariepinus* in selected fish farms of Kumbotso Local Government Area of Kano State, Nigeria. *BayeroJournal of Pure and Applied Sciences*, 2014; 7 (2): 145-149.
17. Clarence SY, Obinna CN. and Shallom NC. Assessment of bacteriological quality of ready to eat food (meat pie) in Benin city metropolis, Nigeria. *African Journal of Microbiology Research*, 2009; 3 (6):390-395.
18. Amadi EN, Kiin KD, Kpormon LB and Robinson VKK. Microbial flora and nutritional composition of adult palm-winebeetle. *International Journal of Current Microbiology and Applied Science*, 2014; 3(11):189-192.
19. Hale LP, Greer PK, Trinh CT and Jame CL. Proteinase activity and stability of natural bromelain preparations. *International Journal of Immunopharmacology*, 2004; 5,783-793.
20. Robinson VK. and Wemedo SA. Molecular characterization of indoor air microorganisms of a model primary health care in Port Harcourt, Rivers State, Nigeria. *Journal of Advances in Microbiology*, 2019; 1-7.
21. Sumbly P, Barbian KD, Gardner DJ, Whitney AR, Welty DM, Long RD, Bailey JR, Parnell MJ, Hoe NP, Adams GG, Deleo RF. and Musser JM. Extracellular deoxyribonuclease made by group A *Streptococcus* assists pathogenesis by enacting evasion of innate immune response. *Proceeding of the NatationalAcademy Science of the USA*, 2005; 102 (5):1679-1684
22. Buchanan JT. DNASE Expression Allows the pathogen Group A *Streptococcus* to Escape Killing in Neutrophil Extracellular Traps. *Science Direct*, 2006; 34, 440-456.
23. Okwesili FC, Paul NA, Parker EJ. and Ikechukwu, U. R. Phytochemical, antihyperglycaemic and lip profile effects of methanom extract fraction of *Ricinus communis* seeds in alloxan induced diabstic male wistar albino. *Asian Journal of Biochemistry*, 2016; 11, 24-33.
24. Sagar A. Basic Microbiology/Culture Media. Xavier's College, Kathmandu, Nepal 2015; p.25.
25. Chakraborty P and Nishith KP. Manual of Practical Microbiology and Parasitology (First Edition). Kolkata, India: New Central Book Agency. 2008; 1-23.
26. Breakwell DP, Moyes RB and Reynold S. Differential staining of bacteria: capsule stain. *Current protocol in microbiology*, 2009; 3-6.
27. Hale LP, Greer PK, Trinh CT and Jame CL. Proteinase activity and stability of natural bromelain preparations. *International Journal of Immunopharmacology*, 2004; 5,783-793.

28. General Laboratory Technology. Percentage Concentration Calculation, Volume per volume, Weight Per Volume. *Journal of Biological Methods*, 2007; 6, 22-80.
29. Rouger A, Odile T. and Monique Z. Bacterial contaminants of poultry meat: source, species and dynamics *Pub med central*, 2017; 5(3): 50.
30. Amelie R, Odile T and Monique Z. Review Bacterial contaminants of poultry meat; Sources, species, and dynamics, *Microorganism*, 2017; 46-48.
31. Golden C, Koehn JZ, Shepon A. and Passarelli S. Aquatic foods to nourish nations. *Nature*, 2021; 598 (7880): 1-6.
32. Darshana BB, Thyagarajan D, Richard RC. and Punniamurthy N. Bacterial pathogens in chicken meat: Review. *International Journal of Life Sciences Research*, 2014; 2(3): 1-7.
33. Bhandare SG, Sherikarv AT, Paturkar AM, Waskar VS. and Zende RJ. A comparison of microbial contamination on sheep/goat carcasses in a modern Indian abattoir and traditional meat shops. *Food control*, 2007; 18, 854-868.
34. Prescott LM, Harley JP. and Klein DA. *Microbiology*, (9th Edition), London: WMC Brown Publishers. 2011; 133.
35. Wang F, Lu C. and Wilner I. From cascaded catalytic nucleic acids to enzymes-DNA nanostructures: controlling reactivity, sensing, logic operations and assembly of complex structures. *Chemical Reviews*, 2014; 114(5): 2881-2941.
36. Kabour GA. Evaluation of microbial contamination of chicken carcasses during processing in Khartoun State M.V.SC thesis Sudan. 2011; University of Science and Technology Sudan.
37. Niu C, Yu D, Yuelan W, Ren H, Jin Y, Zhou W, Li B, Cheng Y, Yue J, Gao Z and Liang L. Common and pathogen-specific virulence factors are different in function and structure. *Virulence*, 2013; 4 (6): 473-482.
38. Behera PC, Tripathy DP and Parija SC. Shatavari: potential for galactagogue in dairy cows. *Indian Journal of Traditional Knowledge*, 2013; 12 (1): 7-9.
39. Wilson JW, Schurr MJ, LeBlanc CL, Ramamurthy R. and Nickerson CA. Mechanism of bacterial pathogenicity. *Postgraduate Medical Journal*, 2002; 78 (918): 216-224.
40. Zanella R, Prestes O, Friggi CA, Martins ML. and Adaime M. An overview about recent advances in sample preparation techniques for pesticides residues analysis in cereal and feed stuff. *International Journal of Pure and Applied Sciences*, 2012; 8, 151-170.

UNDER PEER REVIEW