

# Single stage reconstruction of a post traumatic complete penoscrotal degloving injury: A case report and Review of Literature

## ABSTRACT

Penoscrotal skin degloving injuries are uncommon in Plastic surgery practice which are mostly due to industrial and agricultural related accidents. Although this type of injury is not dangerous to life, it can be psychosocially bothersome to patients if not treated properly. Here, we present the case of a 34-years patient with complete penoscrotal degloving injury due to agricultural machine accident. He came to our trauma centre 4 hours after the accident and treated by partial thickness skin grafting in a single stage. Postsurgical period was normal and he was discharged 9 days after the operation. Follow up at 3 months, patient showed an excellent outcome in terms of aesthesis, a well settled penoscrotal grafts, normal voiding and regained sexual functions.

**Keywords** – *Degloving injury, penoscrotal, genitalia, perineum, split thickness skin graft.*

## INTRODUCTION

Degloving injury of penoscrotal region resulting in incomplete avulsion of the scrotum with normal spermatic cords and testes are usually rare. Mostly skin degloving injuries of penile region are due to accidents at agricultural fields. In this, penoscrotal skin is injured by the entrapment of clothes which are caught by agricultural machines. However, corpora spongiosum, corpora cavernosa and testes are relatively spared. This is probably due to mobile nature of genitalia and special protective covering around the testes by tunica albuginea. Cremasteric reflex also plays a role.<sup>1-3</sup> In this injury mechanism, the penoscrotal skin avulsion injury initiates from the level of pubic symphysis and goes up to corona of penis. However, this type of injury is often difficult to treat due to mobile nature of genitalia, chances of hematoma formation in pelvis, infection and subsequent graft loss.<sup>4</sup> To have an ideal reconstruction of the scrotum, it should be a unistage procedure which gives a non-bulky tissue, maintains thermoregulation, provide a natural looking scrotal ptosis and colour match, resistant to shearing forces from the thighs with minimal donor site morbidity.<sup>5</sup> Here, we present a case of traumatic penoscrotal injury which was treated with a single stage procedure at our hospital.

## CASE PRESENTATION:

A 34 years old patient came to our advanced trauma centre 4 hours after sustaining degloving injury at the penoscrotal region. His loose skin at the penoscrotal level was accidentally pulled off by the rotating blades of a machine which entrapped his clothes while doing agricultural work at his fields. On

arrival, patient was conscious and vitals were stable. Physical examination showed a complete circumferential degloving of shaft of penis and scrotum upto the level of perineum. Both testis and spermatic cord were found to be intact (**Figure 1**). He was managed with pain killers, antibiotics and anti-tetanus injections. A Foley's urinary catheter was inserted. Ultrasound showed that there were no testicular or urethral injuries. The laboratory parameters were normal. Patient was taken to operation theatre and debridement of the necrotic tissues, thorough saline and betadine wash was given under spinal anaesthesia. All foreign bodies were removed. Foley's urinary catheter was changed to silicon catheter. (**Figure 2**).



**Figure 1:-** Complete circumferential penoscrotal degloving injury at presentation exposing testicles and spermatic cords.



**Figure 2:-** Intraoperative pictures following through washing and debridement.

From the right thigh, harvesting of two sheets of partial thickness skin grafts was done. Penile shaft was covered with unmeshed sheet of graft and the graft was fixed with vicryl rapide 4-0 suture. The scrotum defect was covered with minimally meshed partial thickness skin graft. Skin graft dressing was done by a tie over bolster. Around the penile shaft, a sponge was kept to make erect position for good take of the graft (**Figure 3**). These described techniques help to immobilise skin grafting that will help in optimal take of graft.



**Figure 3:-** Single stage reconstruction using split thickness skin grafting. Both testes were sutured together to prevent torsion before grafting.

Clinically, patient was stable and afebrile throughout the hospital stay. Tablet diazepam 10mg HS was given to prevent penile erections and subsequent graft mobility for around 10 days. On post operative days 5, 7 and 9, dressing was changed (**Figure 4**).



**Figure 4:-** At primary dressing on post operative day 5, showing good graft uptake. Machine belt mark can be seen on left thigh.

The patient was discharged on 9<sup>th</sup> post operative day and was kept on oral antibiotics for five days. Patient normally voided urine after the silicon urinary catheter was removed at 2 weeks. At the 3 months follow-up visit, the penoscrotal grafts were settled well. There were no infection signs and no graft contracture related to skin grafting (**Figure 5**). As soon as the graft take was good and wound healed, he was advised to start scar massage with moisturizers like coconut oil. This helps to prevent contractures and make scar supple. He was psychologically satisfied with the excellent cosmetic result with normal sensation, shape and size of his penis.



**Figure 5:-** At 3 month follow up, showing well settled skin graft, good contouring and excellent colour matching.

## DISCUSSION

Generally, only the skin is damaged in penoscrotal degloving injuries which bleeds minimally. There is usually no further damage to the testes, corpora spongiosa and corpora cavernosa. This is due to the fact that this injury occurs in an avascular plane. Scrotal defect can be closed primarily with the available surrounding tissue if the defect size is less than 50%. With significant loss, there are several reconstructive options: partial thickness skin grafts or a different type of musculocutaneous or fasciocutaneous flaps<sup>6-8</sup>. Burial of testes in medial thigh pockets can be avoided due to the associated pain, testicular atrophy, poor cosmetic and psychosocial effects, harmful effects on spermatogenesis, risk of pouch infection and endocrine dysfunction.<sup>9</sup> Medial thigh pedicle flaps were considered by some authors but these can lead to deformed contour of the scrotum which can be unsightly in appearance.<sup>10</sup> A proximal thigh flap cannot cover both the penoscrotal and perineo-scrotal junction in a single stage. Also, it can lead to obliteration of the perineal cleft which leads to scrotal contour deformity. In addition, these pedicled flaps may need multi-staged operations like shown by Zanettini et al<sup>11</sup> and Conley et al<sup>12</sup> which can extend up to several months that may have negative psychological effects. It was shown that better spermatogenesis can be achieved with thinner tissues over scrotal region.<sup>13</sup>

Many authors have reported excellent results of penoscrotal degloving injury managed with split skin grafting.<sup>14-16</sup> As the present case had complete circumferential loss of penoscrotal skin, the

reconstruction was done by using partial thickness skin grafting in a single stage. Skin defects of penoscrotal region can be best managed with this technique as it is simple and tolerated well by the patients. Other advantages of this method are that it is easy to execute by the surgeons who are inexperienced, it gives almost natural temperature for spermatogenesis, provides a natural looking scrotum, less post operative morbidity, good colour match, great scrotal contouring and can be finished in single stage if executed appropriately. Patient returns early to normal lifestyle without any psychosocial problems. Sanchez et al<sup>17</sup> stated that there are no differences between gracilis muscle flap and skin grafting in terms of testicular hormonal functions for scrotal reconstruction. Wang et al.<sup>18</sup> advised skin grafting over the scrotum of younger patients who wished to remain fertile. Demir et al<sup>19</sup> studied on rodents and found that as long as dartos thermoregulatory function remains intact, there will not be any effect on testicular hormonal functions. Sun et al<sup>20</sup> showed that the physiological function of the testis in terms of volume of sperms, motility, time taken to liquefy and their overall numbers at the 1 year follow up was within normal limits in patients with skin grafting and this was maintained for at least 2 years. However, skin grafting may lead to painful erections due to penoscrotal or perineo-scrotal junctional contractures which can be relieved by daily corticosteroid cream massage. The reconstructed scrotum in our patient was mobile, cosmetically appealing, had excellent colour match and didn't alter the sexual functions. The grafted scar had a score of 2 out of 13 as per the Vancouver scar scale which indicates great aesthesis. Similar cases reported by various authors.<sup>21</sup>

## CONCLUSION

Penoscrotal degloving injuries are rare surgical emergencies which can be difficult to manage by the surgeons as well as can be distressing to the patients. Reconstruction of the penoscrotal skin defect is recommended as early as possible. There is no perfect treatment for all cases. Instead, the ideal methodology is determined by the patient's expectations and defect characteristics. The present case has been managed with a single staged split thickness skin grafting which gave good cosmetic and functional results, also maintain the sexual and testicular function.

## CONSENT

The patient provided written informed consent for the use of photographs for the article.

## REFERENCES

1. Gencosmanoğlu R, Bilkay U, Alper M, et al. Late results of split-grafted penoscrotal avulsion injuries. *J Trauma*. 1995;39:1201–3.
2. Selikowitz SM. Penetrating high-velocity genitourinary injuries. Part I. Statistics mechanisms, and renal wounds. *Urology*. 1977;9:371–6.

3. Finical SJ, Arnold PG. Care of the degloved penis and scrotum: a 25-year experience. *PlastReconstr Surg*. 1999;104:2074–8.
4. Shetty BSK, Rao PJ, Menezes RG. Traumatic degloving lesion of male external genitalia. *J Forensic Legal Med*. 2008;15:535.
5. Iwuagwu FC, Mopuri N, Fitzgerald O'Connor E, Scrotal reconstruction with modified pudendal thigh flaps, *British Journal of Plastic Surgery*. (2015)
6. Hsu H, Lin CM, Sun TB, Cheng LF, Chien SH. Unilateral gracilismyofasciocutaneous advancement flap for single stage reconstruction of scrotal and perineal defects. *J PlastReconstrAesthet Surg*. 2007;60(9);1055-9.
7. Bickell, M., Beilan, J., Wallen, J., and Carrion, R. 2016. "Advances in Surgical Reconstructive Techniques in the Management of Penile, Urethral, and Scrotal Cancer." *UrolClin North Am* 43 (4): 545-59.
8. Mopuri, N., O'Connor, E. F., and Iwuagwu, F. C. 2016. "Scrotal Reconstruction with Modified Pudendal Thigh Flaps." *J PlastReconstrAesthetSurg* (69): 278-83.
9. Maharaj, D., Naraynsingh, V., Perry, A., and Ramdasset, M. J. 2002. "The Scrotal Reconstruction Using the 'Singapore Sling'." *PlastReconstrSurg* 110 (1): 203-5.
10. Jeffrey D. Friedman reconstruction of the perinium in Grabb and Smith's plastic surgery. 6th ed. p. 713.
11. Zanettini, L. A., Fachinelli, A., and Fonseca, G. P. 2005. "Traumatic Degloving Lesion of Penile and Scrotal Skin." *IntBraz J Urol* 31: 262-3.
12. Conley, J. J. 1956. "A One-Stage Operation for the Repair of the Denuded Penis and Testicles." *N Y State J Med* 56 (19): 3014-6.
13. Wang DL, Wang YM, Zheng H, Feng J, Fu SN, Gao ZY, et al. An experiment study and clinical observation of the testicle spermatogenesis after scrotum reconstruction. *Zhonghua Zheng Xing Wai Ke Za Zhi* 2004;20:203-5.
14. Aineskog, H., and Huss, F. 2016. "A Case Report of a Complete Degloving Injury of the Penile Skin." *Int J Surg Case Rep* (29): 1-3.
15. Campbell, R. M. 1957. "Dermatome Grafting of the Totally Denuded Testes." *PlastReconstrSurg* (19): 509-13.
16. Alton, J. D. 1963. "Complete Avulsion of the Scrotum." In *Transactions of the Third International Congress of Plastic Surgery*. Edited by T. R. Broadbent. Amsterdam: Excerpta Medica Foundation.
17. Sanchez G, Ardavin JP, Munoz C . Scrotal reconstruction with pedicledgracilis muscle flap versus split thickness skin graft. 2018. *CirurgiaPlastica libero Latinoamericana* 44(3) : 297-301.

18. Wang, D., Luo, Z., Sun, G., and Wei, Z. 2009. "Long-Term Prognosis of Free Skin-Grafted Penoscrotal Avulsion Injuries in Two Patients." *J Plast Reconstr Aesthet Surg* (62): 385-7.

19. Demir, Y., Aktepe, F., Kandal, S., Sancaktar, N., and Turhan-Haktanir, N. 2011. "The Effect of Scrotal Reconstruction with Skin Flaps and Skin Grafts on Testicular Function." *Ann Plast Surg* 68 (3): 308-13.

20. Sun, G., Wang, D., Wei, Z., Jin, W., and Deng, C. 2011. "Effect of Scrotal Reconstruction with Free Skin Graft on Spermatogenesis." *ZhongguoXiu Fu Chong Jian Wai Ke Za Zhi* 25 (11): 1357-9.

21. Alkahtani et al. 2020 Traumatic Degloving Injury of Penile and Scrotal Skin: A Case Report. *Plast Reconstr Surg Glob Open*. 2020 Aug; 8(8): e3024.

UNDER PEER REVIEW