

Impact of irrigation water on the bacteriological quality of lettuce (*Lactuca sativa*) produced in the market garden zone of Niamey, identification of enterobacteria

Abstract:

The urban community of Niamey is characterized by a prodigious development of urban and peri-urban agriculture. Lettuce (*Lactuca sativa*) is one of the main crops grown in these urban and peri-urban areas. The aim of this study is to assess the impact of irrigation water on the bacteriological quality of lettuce produced on market garden sites in the urban community of Niamey. Three (3) market garden sites were studied (*Harobanda* (site 1), *Gamkallé* (site 2) and *Gounti yéna* (site 3)). Eighteen (18) lettuce samples and twenty-nine (29) irrigation water samples were taken from the market garden sites and subjected to microbiological analysis using conventional methods appropriate to each indicator. Results showed that irrigation water samples from site 1 were heavily contaminated with Total Coliforms (TC) and Faecal Coliforms (FC) ($3.65 \pm 7.88 \cdot 10^6$ and $1.41 \pm 3.04 \cdot 10^6$ CFU/100ml respectively), those from site 2 by Faecal Streptococci (FS) ($1.19 \pm 2.03 \cdot 10^5$ CFU/100ml) and from site 3 by Enterobacteria (Ent) ($1.61 \pm 2.37 \cdot 10^8$ CFU/100ml) and *Escherichia coli* ($8.22 \pm 11.86 \cdot 10^5$ CFU/100ml). Thus, lettuce samples from site 1 are more contaminated with CT ($5.32 \pm 8.32 \cdot 10^6$ CFU/g) and Ent ($3.10 \pm 0.95 \cdot 10^7$ CFU/g). Then on site 2, CF ($1.35 \pm 1.13 \cdot 10^6$ CFU/g) and SF ($5.30 \pm 8.21 \cdot 10^5$ CFU/g) predominate, and finally site 3 is more contaminated with *E. coli* ($2.43 \pm 0.38 \cdot 10^5$ CFU/g). However, enterobacteria species such as *Citrobacter freundii*, *Enterobacter cloacae*, *Enterobacter sakazakii*, *Escherichia coli*1, *Klebsiella pneumoniae ssp ozaenae*, *Pantoea spp 1*, *Proteus mirabilis*, *Raoultella terrigena*, *Salmonella cholerae suis ssp arizonae*, *Salmonella spp*, *Serratia liquefaciens*, *Serratia marcescens* and *Serratia odorifera*1 were identified in irrigation water samples and lettuce at these sites. In addition, the quality of the irrigation water is not suitable for irrigation, and strongly influences the contamination of the lettuce produced by the bacteria. Consumption of this lettuce without the minimum of precautions could prove dangerous to consumers.

Keywords: lettuce, watering water, contamination, enterobacteria, Niamey/Niger

• Introduction

Over the past two decades, as a result of rapid urbanization and economic concentration, agriculture has expanded considerably in urban and peri-urban areas of West Africa [1, 2]. In difficult contexts marked in particular by a lack of financial resources for the supply of drinking water, market gardeners are driven to use wastewater for irrigation and animal dung as fertilizer for the soil [2, 3]. In most developing countries, crops are irrigated using treated, poorly treated, diluted or even raw wastewater [4]. The reuse of

wastewater in market gardening and horticulture has thus become a common practice, responding among other things to demographic growth in cities and youth unemployment [5]. In fact, their reuse without prior treatment has negative impacts on both the environment and people's health [6]. From a health point of view, the greatest threat from wastewater reuse is the presence of pathogenic microorganisms. Viruses, bacteria and parasites can be present in wastewater in high concentrations and survive for long periods. Spreading this water on vegetables can cause microbiological contamination of these products and encourage the introduction of pathogenic microorganisms into the food chain [7, 8].

As in other African cities, Niamey has seen the development of local agriculture. The urban community of Niamey is characterized by inadequate drainage systems and the almost total absence of wastewater treatment plants. As a result, wastewater is discharged directly into the lowlands and other receiving areas, notably the River Niger, without prior treatment. Before flowing into the river, this water passes through several market-gardening sites, notably through streams, the most important of which in Niamey is the Gounti yéna outfall [9]. Three (3) sources of irrigation water have been identified in Niamey: river water, well water and wastewater according to market garden sites [10]. Market gardeners on the *Gamkallé* corniche use river water, those on *Saga* use well water and those on *Gounti yéna*, a non-permanent tributary, use wastewater from toilets. Thanks to this water, market gardening is practicable throughout the 12 months of the year [10].

This study aims to assess the impact of irrigation water on the bacteriological quality of lettuce produced in the market gardening zone of the city of Niamey.

- **Material and method**

- **Study area**

The study was carried out in the urban community of Niamey. The city of Niamey is located in the south-western part of Niger between 13° 24' and 13°35'N latitude and, 2°00' and 2°15'E longitude with an altitude between 160 and 250 m. Its administrative boundaries extend over 552.27km², including around 297.46km² of urbanized area [11]. Niamey's population is estimated at around 1,407,635. The city of Niamey is subdivided into five (5) communal arrondissements, whose population distribution by communal arrondissement is as follows: Niamey I: 287,902 inhabitants; Niamey II: 338,455 inhabitants; Niamey III: 223,685 inhabitants; Niamey IV: 376,271 inhabitants; Niamey V: 181,321 inhabitants [12]. The climate is Sahelo-Sudanian, with a long dry season from October to May and a short rainy season from June to September. The cold dry season is the most favorable for vegetable production, during which most crops are grown. This study has been focused on three (3) major vegetable-growing sites in the urban community of Niamey.

- Site1, located in Niamey's commune V (*Harobanda*), stretches along the river over a length of around 2km and a width varying from 200m to 700m depending on the extent of the riverbed.

- Site2, located right in the center of town, the *Gamkalé* corniche forms part of the left bank of the River Niger. It is bounded on one side by the Kennedy Bridge and on the other by the Niamey abattoir.
- Site3 is the *Gounti yéna* valley, covering an area of around 38.1 ha. The environment around this valley is complex: less than 2 km from the valley and often within it, there are roads with high traffic density; a wastewater treatment plant collecting urban water and open dumps [11] (Djibo et al., 2016).




Figure 1 : Geological map of the urban community of Niamey presenting the market garden sites of study

- **Sampling**

Lettuce and irrigation water samples were taken from three (3) production sites in the urban community of Niamey. Samples were taken under sterile conditions, using single-use sterile gloves and alcohol to avoid external contamination. A lettuce sample corresponds to three (3) heads of lettuce, weighing around 150g, taken at random from different corners of the field. A sample of irrigation water corresponds to 100ml of water taken from the source by dipping the bottle directly into the pool. An information sheet was attached to each sample. A total of eighteen (18) lettuce samples and twenty-nine (29) irrigation water samples were taken.



a



b



c



d

Figure 2: some sources of irrigation water (a;b) and lettuce dishes (c; d) at the Harobanda site

- **Transport conditions**

Each sample collected was well packaged in a polyethylene bag, then carefully labeled. The samples were then transferred to the microbiology laboratory of the Faculty of Science and Technology (FAST), after being packaged and placed in a cooler containing carboglass to keep the temperature down to around 4°C.

- **Microbiological analysis of samples**

Preparation of stock solutions: The stock solution was prepared by grinding the lettuce sample in a sterile polyethylene bag around the flame of a Bunsen burner. Then, 25 grams of the crushed material were taken and introduced directly into 225 mL of previously prepared and sterilized buffered peptone water. Next, 1 mL of each stock solution was taken and introduced into a test tube containing 9 mL of

buffered peptone water to perform the various decimal dilutions in accordance with ISO 6887-V08-010-6 (2013). All water samples were homogenized by vigorously shaking the bottles in a vertical motion. In fact, 1 mL of each water sample was taken and used directly for the decimal dilutions.

Bacterial culture: Each indicator was tested and enumerated using the appropriate methods. Total Aerobic Mesophilic Flora (FAMT) was counted according to AFNOR NF VOS-051 on Plate Count Agar (PCA); coliforms according to NF ISO 4832 on Violet Red Bile Lactose (VRBL) medium; fecal streptococci to ISO 7899/1 on Kanamycin Aesculin Azide (KAA) medium; *Clostridium perfringens* to ISO 7937 on Tryptone Sulfite Neomycin (TSN) agar; *E. coli* according to ISO 18140 on Eosine Methylene Blue (EMB) medium, and enterobacteria according to ISO 21528-1 on Mac conkey agar.

Enterobacteria identification: Enterobacteria from water and lettuce samples were randomly selected and streaked onto nutrient agar. Biochemical characterization was carried out on Api 20 E strips. Strains were identified by comparing their characteristics with those of known taxa as described in the bio Mérieux SA manual.

- **Interpretation**

Bacterial load was calculated in accordance with ISO 7218 (2007) using the following formula:

$$N = [13]$$

- Σc = Total number of colonies counted in plates with colony counts between 15 and 300.
- n_1 = number of plates counted in the first dilution;
- n_2 = number of plates counted in the second dilution;
- d = dilution factor at which 1st counts were made;
- v = inoculum volume.

- **Statistical analysis**

IBM SPSS statistics software version 23.0.0.0 was used to calculate means and standard deviations.. The document was drafted using Microsoft Word. Arc Gis software was used to design the geographical map of the study area. Differences were considered significant for values of $P < .05$.

- **Results and discussion**

- **Results**

Water contamination

The microbiological characteristics of the irrigation water are shown in Table (1). Average total coliform loads ranged from $1.53 \pm 3.15.106$ to $3.65 \pm 7.88.106$ CFU/100ml of water for sites 2 and 1 respectively.

Fecal coliform loads varied from $4.29 \pm 7.88.10^5$ CFU/100ml (site 2) to $1.41 \pm 3.04.10^6$ (site 1). Enterobacteria have the highest values of all indicators. These range from $4.34 \pm 6.84.10^7$ (site1) to $1.61 \pm 2.37.10^8$ CFU/100ml (site3). Average E. coli loads ranged from $3.77 \pm 5.68.10^5$ (site1) to $8.22 \pm 11.86.10^5$ CFU/100ml (site3). Fecal streptococci present the lowest loads. They range from $2.18 \pm 5.25.10^4$ (site1) to $1.19 \pm 2.03.10^5$ CFU/100ml (site2) of irrigation water. Indeed, site1 is more loaded with total and faecal coliforms, site2 with faecal streptococci and site3 with enterobacteria and E. coli. The values obtained for all irrigation water samples are above the threshold values.

Table 1 : level of contamination of irrigation water by production site

Sites	Means loads \pm standard deviation of irrigation water samples (UFC/100ml)				
	TC	FC	Ent	E. coli	FS
Site 1	$3,65 \pm 7,88.10^{6a}$	$1,41 \pm 3,04.10^{6b}$	$4,34 \pm 6,84.10^{7c}$	$3,77 \pm 5,68.10^{5b}$	$2,18 \pm 5,25.10^{4a}$
Site 2	$1,53 \pm 3,15.10^{6a}$	$4,29 \pm 7,88.10^{5b}$	$8,15 \pm 12,32.10^{7c}$	$5,29 \pm 8,96.10^{5b}$	$1,19 \pm 2,03.10^{5a}$
Site 3	$1,47 \pm 4,33.10^{6a}$	$2,33 \pm 4,64.10^{5b}$	$1,61 \pm 2,37.10^{8c}$	$8,22 \pm 11,86.10^{5b}$	$3,70 \pm 8,23.10^{4a}$
Total	$2,37 \pm 5,75.10^6$	$7,64 \pm 20,15.10^5$	$9,24 \pm 15,91.10^7$	$5,64 \pm 8,83.10^5$	$5,20 \pm 1,19.10^4$

TC : Total coliform ; FC : Faecal Coliform ; Ent : Enterobacteria ; E. coli : Escherichia coli ; FS: Faecal Streptococcus. Site1 : Harobanda ; site2 : Gamkalé ; Site3 : Gounti yéna. Values with the same letter in the same column are not significantly different ($P > .05$).

Contamination of lettuce samples

Table 2 shows the bacteriological characteristics of lettuce samples according to production site. The level of contamination ranged from $1.60 \pm 0.52.10^5$ (site3) to $5.32 \pm 8.32.10^6$ CFU/g (site1) of lettuce for total coliforms and $1.65 \pm 0.92.10^5$ (site3) to $1.35 \pm 1.13.10^6$ CFU/g (site2) of lettuce for faecal coliforms. Enterobacteria recorded the highest level of contamination. Loads ranged from $1.13 \pm 0.09.10^7$ (site3) to $3.10 \pm 0.95.10^7$ CFU/g (site1) of lettuce. In addition, a predominance of TC ($5.32 \pm 8.32.10^6$ CFU/g) and Enterobacteria ($3.10 \pm 0.95.10^7$ CFU/g) was observed at site1. Site2 was heavily contaminated with CF ($1.35 \pm 1.13.10^6$ CFU/g) and SF ($5.30 \pm 8.21.10^5$ CFU/g), while site3 was more heavily loaded with E. coli ($2.43 \pm 0.38.10^5$ CFU/g lettuce). The level of contamination of all lettuce samples, whatever the site, was higher than the French standards that vegetables for human consumption must meet.

Table 2 : Microbial contamination levels of lettuce samples by production site

Means loads \pm standard deviation of lettuce samples (UFC/100ml)	
---------------------------------------------------------------------	--

Site s	FAMT	CT	CF	Ent	SF	CP	<i>E. coli</i>
Site 1	2,49±2,50. 10 ^{7a}	5,33±8,47. 10 ^{6a}	2,14±1,19.1 0 ^{6b}	9,05±13,54. 10 ^{6a}	4,56±7,23. 10 ^{5a}	3,80±6,13. 10 ^{4a}	1,83±1,10. 10 ^{5a}
Site 2	5,31±1,63. 10 ^{7b}	8,30±8,36. 10 ^{5a}	1,35±1,21.1 0 ^{6ab}	3,20±1,22.1 0 ^{6a}	6,57±9,40. 10 ^{5a}	1,85±1,94. 10 ^{4a}	2,00±2,10. 10 ^{3b}
Site 3	3,03±1,76. 10 ^{6a}	1,60±1,26. 10 ^{5a}	1,65±0,99.1 0 ^{5a}	4,40±2,17.1 0 ^{5a}	-	-	1,95±1,17. 10 ^{5b}
Total	2,94±2,66. 10 ⁷	2,68±5,98. 10 ⁶	1,44±1,28.1 0 ⁶	5,19±9,47.1 0 ⁶	4,22±7,33. 10 ⁵	2,30±4,35. 10 ⁴	1,26±1,24. 10 ⁵

TC : Total coliform ; FC : Faecal coliform ; Ent : Enterobacteria ; E. coli : Escherichia coli ; FS : Faecal Streptococcus. Site1 : Harobanda ; site2 : Gamkalé ; Site3 : Gounti yéna. Values with the same letter in the same column are not significantly different ($P > .05$).

Comparative analysis of irrigation water and lettuce contamination levels

Figure 3 shows the variation curves in contamination levels for irrigation water and lettuce. It can be seen that all loads are above standards. For all indicators, the irrigation water samples are more contaminated than the lettuce samples. A certain similarity is observed in the variation of loads according to indicators between water and lettuce samples.

Figure 3: level of contamination of irrigation water and lettuce samples

Origin of faecal contamination

TC/FS ratios are shown in table (3). Fecal contamination of irrigation water samples was of human origin at all sites (CT/SF >4). Fecal contamination of human origin was also observed in lettuce samples from sites 1 and 3. Contamination of lettuce samples from site 2 was mixed but predominantly human (4 >CT/SF > 2).

Table 3 : faecal contamination origin

	Ratio faecal coliforms /faecal streptococci		
	Site 1	Site 2	Site 3
Lettuce	69,68	2,55	9,43
Origin of faecal contamination	Human	Mixed predominantly human	Human
Water	59,80	12,50	20,56

Origin of faecal contamination	Human	Human	Human
---------------------------------------	--------------	--------------	--------------

Site1 : Harobanda ; site2 : Gamkalé ; Site3 : Gounti yena.

Diversity of enterobacteria isolated at production sites

Table (4) shows the diversity of *Enterobacteriaceae* isolated at the production sites. A total of thirteen (13) species belonging to nine (9) genera were identified. The species identified were *Citrobacter freundii*, *Enterobacter cloacae*, *Enterobacter sakazakii*, *Escherichia coli*1, *Klebsiella pneumoniae* ssp ozaenae, *Pantoea* spp 1, *Proteus mirabilis*, *Raoultella terrigena*, *Salmonella arizonae*, *Salmonella* spp, *Serratia liquefaciens*, *Serratia marcescens* and *Serratia odorifera* 1. On all sites, *Salmonella* spp, *Escherichia coli*1, *Citrobacter freundii*, *Enterobacter cloacae*, *Enterobacter sakazakii* and *Serratia liquefaciens* predominate (with 26.1%, 13.0% and 8.70% respectively for each of the last 4 species).

Table 4: Diversity of enterobacteria species isolated at production sites

<i>Species identified</i>	Distribution by production site (%)			Total
	site 1	site 2	site 3	
<i>Citrobacter freundii</i>	-	-	(2)8,70 ^a	(2)8,70
<i>Enterobacter cloacae</i>	(1)4,30 ^a	(1)4,30 ^a	-	(2)8,70
<i>Enterobacter sakazakii</i>	(1)4,30 ^a	-	-	(1)4,30
<i>Escherichia coli</i> 1	(2)8,70 ^a	-	(1)4,30 ^a	(3)13,0
<i>Klebsiella pneumoniae</i> ssp ozaenae	-	(1)4,30 ^a	-	(1)4,30
<i>Pantoea</i> spp 1	-	-	(1)4,30 ^a	(1)4,30
<i>Proteus mirabilis</i>	-	(1)4,30 ^a	-	(1)4,30
<i>Raoultella terrigena</i>	(1)4,30 ^a	-	-	(1)4,30
<i>Salmonella arizonae</i>	-	-	(1)4,30 ^a	(1)4,30
<i>Salmonella</i> spp	(1)4,30 ^a	(2)8,70 ^a	(3)13,0 ^a	(6)26,1
<i>Serratia liquefaciens</i>	(1)4,30 ^a	-	(1)4,30 ^a	(2)8,70
<i>Serratia marcescens</i>	-	(1)4,30 ^a	-	(1)4,30
<i>Serratia odorifera</i> 1	-	(1)4,30 ^a	-	(1)4,30
Total	(7)30,4	(7)30,4	(9)39,1	(23)100

Site1 : Harobanda ; site2 : Gamkalé ; Site3 : Gounti yena. Values with the same letter in the same column are not significantly different ($P > .05$).

Frequency of enterobacteria species isolated from irrigation water

Enterobacteriaceae species isolated from irrigation water and their frequencies are presented in **Table (5)**. A total of eight (08) species belonging to seven (7) genera were identified from irrigation water. *Escherichia coli* 1, *Salmonella spp* and *Serratia liquefaciens* were isolated with a frequency of 21.4% (three (3) species each). Of all the species, only one was identified at site 2 (*Salmonella spp*).

Table 5: Diversity of species isolated from irrigation water

Species of enterobacteria identified	Distribution (n)%			Total
	Site1	Site2	Site3	
<i>Citrobacter freundii</i>	-	-	(2)14,30 ^a	(2)14,3
<i>Enterobacter cloacae</i>	(1)7,10 ^a	-	-	(1)7,1
<i>Enterobacter sakazakii</i>	(1)7,10 ^a	-	-	(1)7,1
<i>Escherichia coli</i> 1	(2)14,30 ^a	-	(1)7,10 ^a	(3)21,4
<i>Raoultella terrigena</i>	(1)7,10 ^a	-	-	(1)7,1
<i>Pantoea spp</i> 1	-	-	(1)7,10 ^a	(1)7,1
<i>Salmonella spp</i>	(1)7,10 ^a	(1)7,10 ^a	(1)7,10 ^a	(3)21,4
<i>Serratia liquefaciens</i>	(1)7,10 ^a	-	(2)14,30 ^a	(3)21,4
TOTAL	(7)50,0	(1)7,1	(6)42,9	(14)100,0

Site1 : Harobanda ; site2 : Gamkalé ; Site3 : Gounti yéna. Values with the same letter in the same column are not significantly different (P>.05).

Frequency of species isolated from lettuce samples by site

Figure (3) shows the species identified from lettuce samples according to production site. A total of seven (7) species belonging to five (5) genera were identified in the lettuce samples. Of these, six (6) were identified at site 2 (11.1% each) and three (3) at site 3 (11.1% for *Salmonella ssp arizonae* and 22.2% *Salmonella spp*). No enterobacteria species were identified at site1.

Figure 4: frequency of enterobacteria species isolated from lettuce depending on market garden site

- **Discussion**

This study assessed the impact of irrigation water on the bacteriological quality of lettuce produced in the market-garden zone of the urban community of Niamey. Three (3) market garden sites in the urban community of Niamey were studied. The production sites concerned are: *Harobanda* (site1), *Gamkallé* (site2) and *Gounti Yéna* (site3). Total and faecal coliforms, faecal streptococci, enterobacteria and *Escherichia coli* were tested in samples of irrigation water and lettuce. The study continues with the characterization of some enterobacteria species isolated from water and lettuce.

The loads of all the indicators tested in the irrigation water samples exceeded the values set in the WHO guide (10^5 CFU/100 ml) for non-restrictive irrigation [14]. A similar finding was made in Senegal by [8]. Total coliform loads ranged from $1.47 \pm 4.33 \cdot 10^6$ to $3.65 \pm 7.88 \cdot 10^6$ CFU/100ml of irrigation water. In Côte d'Ivoire, [15] reported results slightly higher than these. The values obtained by these authors range from $2.19 \cdot 10^7$ CFU/100ml to $2.74 \cdot 10^7$ CFU/100ml. [16], on the other hand, reported results lower than these, ranging from 0.95 ± 0.10 to 3.38 ± 0.00 Log₁₀ CFU/100ml in irrigation water of lettuce in Maroua (Cameroon). Faecal coliform loads were in order of $1.41 \pm 3.04 \cdot 10^6$ CFU/100ml of irrigation water. These results corroborate those reported by [4]. Faecal streptococci loads ranged from $2.18 \pm 5.25 \cdot 10^4$ to $1.19 \pm 2.03 \cdot 10^5$ CFU/100ml. [2] found almost similar results ($2.1 \cdot 10^4 \pm 2.7 \cdot 10^3$ CFU/100ml). Heavy contamination of Gourou basin water [15].

What is the main implication of this finding??

The irrigation water of Site1 is more loaded with total and faecal coliforms (respectively $3.65 \pm 7.88 \cdot 10^6$ and $1.41 \pm 3.04 \cdot 10^6$), and site3 with enterobacteria and *Escherichia coli* (respectively $1.61 \pm 2.37 \cdot 10^8$ and $8.22 \pm 11.86 \cdot 10^5$ CFU/100ml). Site2, on the other hand, was more loaded with faecal streptococci ($1.19 \pm 2.03 \cdot 10^5$ CFU/100 ml). This higher level of contamination in site3 can be explained by the nature of the water used for watering, which is wastewater from the city. This water is discharged directly into the lowlands and other receiving areas, notably the River Niger, without any prior treatment. Before flowing into the river, this water passes through a number of market garden sites, notably via watercourses, the most important of which in Niamey is the *Gounti Yena* outfall [9]. It should be noted that the loads of these indicators are higher than the values set (10^3 CFU/ 100 ml) by WHO guidelines for the safe discharge of wastewater into the environment [17]. A similar finding has been made by other authors in Africa [15, 18, 19, 20, 21].

In addition, the microbiological quality of lettuce samples, regardless of the production site considered, is not fit for human consumption. All loads are well above the values recommended by the French Standardization Association [22, 23]. This level of contamination could be explained not only by poor production practices, but also by the use of wastewater as irrigation water. Indeed, the level of FAMT contamination varies from $3.03 \pm 1.24 \cdot 10^6$ CFU/g (site3) to $5.31 \pm 1.27 \cdot 10^7$ CFU/g of lettuce (site2). The variation in load is significant between these sites ($P = .000$). The higher level of contamination in lettuce samples from site2 could be explained by the proximity of this site to industrial zones, which discharge their wastewater directly into the lowland through this site which contribute to water pollution. Thus, [10] reported the use of poor condition water at this site. These results are similar to those obtained by [23] in Côte d'Ivoire. Poor exposure and storage conditions for vegetables (dust, insects, rodents, flies, etc.) are also responsible for the high levels of microorganisms on these plant products [23, 24, 25].

The highest level of contamination in total coliforms is around $5.33 \pm 8.47 \cdot 10^6$ CFU/g and in faecal coliforms around $2.14 \pm 1.19 \cdot 10^6$ CFU/g obtained for site1. There was no significant difference between these sites and their contamination levels ($P > .05$). Lettuce samples are highly contaminated with

coliforms (total and faecal). The latter are good indicators of hygiene. This can be explained by the use of organic fertilizers (manure), which are made up of animal faeces. Some market gardeners use watering cans to water their vegetables, and in order to collect water, they step into the pool, sometimes barefoot. In Côte d'Ivoire, for example, [26] reports that the use of excreta contributes to the contamination of lettuce produced on Abidjan's market garden sites. In addition, the presence of animals such as reptiles, migratory birds and domestic pets (dogs, etc.) can contribute to the faecal contamination of production sites. The excreta of these animals are carried by run-off water during the rainy season to sources of water for irrigating vegetables [27, 28]. This may explain the presence of these indicators in lettuce samples. According to [29], irrigation techniques play an important role in the contamination of lettuce. The sprinkling of polluted water on lettuce with a watering can thought to be the root cause of *Salmonella* contamination. A similar finding has been made by other authors [23, 26]. The presence of enteric bacteria in lettuces suggests a lack of good hygiene practices, and faecal contamination could be due to the inappropriate processing undergone by these raw edible vegetables. Indeed, enterobacteria are normal hosts of the digestive tract of humans and animals, capable of proliferating in abundance in the environment (soil and water) and thus participating in the major cycles of organic matter degradation [28, 30]. These are indicators of faecal contamination that provide a more complete picture of potentially pathogenic germs [23, 26, 29, 31].

The highest level of faecal Streptococcus contamination is observed at site2 ($6.57 \pm 9.40 \cdot 10^5$ CFU/g). A high level of faecal streptococcus contamination has been reported by other authors in other countries [26, 27, 32]. Also noteworthy is the high *E. coli* load of $1.95 \pm 1.17 \cdot 10^5$ CFU/g at site3. This can be explained by the proximity of the *Gounti yéna* valley (site3) to metallurgical activity centers, main roads and uncontrolled dumps, but also by the use of untreated raw wastewater for irrigation of market garden crops [33]. Indeed, these results corroborate those obtained in Nigeria [34, 35], Egypt [36], Ghana [37] and Côte d'Ivoire [38].

On the other hand, faecal contamination of irrigation water and lettuce samples is almost entirely of human origin at almost all sites. Only faecal contamination of lettuce samples from site 2 was of mixed origin, predominantly human. This situation may be due to the state of the market garden environment in the urban community of Niamey. The results of the survey carried out on these sites showed a lack of sanitary facilities both in the fields and in the vicinity of the fields. Some 85% to 90% of fields have no latrines, and 80% to 90% have none in their immediate vicinity. The lack of latrines in or near fields leads market gardeners to defecate in the open air in or near fields [39]. This defecation in the vicinity of fields, constitutes another additional source of contamination for vegetables [38]. In addition, sites2 and 3 receive wastewater, which is most often used for watering (generally on site 3). A similar situation has been observed in Côte d'Ivoire [2]. They reported that the source of faecal contamination of well water at site 2 was its location on the edge of the Ebrié Lagoon, the main receptacle for wastewater from the city of Abidjan. It is also a place of defecation for local populations [2, 40, 41].

However, a wide diversity of *Enterobacteriaceae* species has been isolated from irrigation water and lettuce samples. This diversity is represented by twenty-three (23) species, belonging to nine (9) genera of *Enterobacteriaceae*. It generally consists of *Salmonella spp*, *Escherichia coli*1, *Citrobacter freundii*, *Enterobacter cloacae*, *Enterobacter sakazakii*, *Serratia liquefaciens*. The presence of *Enterobacteriaceae* in lettuce samples could be due to the precarious hygienic conditions under which lettuce is grown, as previously reported by [2, 42]. The presence of these germs indicates the presence of enteropathogenic bacteria in lettuce. Thus, [16] report the presence of enterobacteria in irrigation water and lettuce samples. Many of these enteric bacteria are incriminated in a growing number of collective food poisoning outbreaks [16, 43, 44]. *Salmonella enterica* serovars are among the pathogenic bacteria causing serious health problems in humans [2, 45].

- **Conclusion**

The aim of this study was to assess the impact of irrigation water on the bacteriological quality of lettuce produced in the market-garden areas of the urban community of Niamey, and to identify a number of enterobacteria species. The results show that irrigation water has a negative impact on the bacteriological quality of lettuce produced in these areas. Microbiological analysis results show that all lettuce contamination indicators are above French standards for fresh vegetables. In addition to the numerous species of *Enterobacteria* isolated from the irrigation water, these were also found in the lettuce samples. The worrying potential for the spread of food-borne infectious diseases therefore exists in the consumption of lettuce. If consumers fail to observe hygiene rules, eating raw lettuce may increase the risk of carrying infectious diseases such as cholera, typhoid fever, gastroenteritis, etc. It would be important to establish the antibiotic resistance profile of these *Enterobacteria* in order to better control infectious diseases.

References

1. Cissé G. (1997). Health impact of the use of polluted water in urban agriculture: case of market gardening in Ouagadougou (Burkina Faso). Thesis, Ecole Polytechnique Fédérale de Lausanne, 31p.
2. Koffi-Nevry R., Assi-Clair BJ., Assemmand EF., Wognin AS., Koussémon M. (2012). Origin of evidence of fecal contamination of water used to water lettuce (*Lactuca sativa*) grown in the peri-urban area of Abidjan. *Journal of Applied Biosciences*, 52:3669–3675.
3. Petterson SR., Ashbolt NJ., Sharma A. (2010). Microbial risks from wastewater irrigation of salad crops: a screening-level risk assessment. *J Food Sci.*, 75(5): 283-290.
4. Djegbe I., Tamou-Tabé Ts., Topanou N., Soglo Fm., Paraiso A., Djouaka R., Kelome Cn., (2018). Seasonal variation in the physicochemical and microbiological quality of irrigation water and vegetables from the Bawéra market gardening site and associated health risks. *International Journal of Biology and Chemistry Sciences* 12(2): 781-795.

5. Thiaw, (2006). The health quality of market garden products in the city of Ouagadougou: Impact of the irrigation water source on human health. Engineering dissertation, University of Burkina Faso, 68p.
6. Barour AA., Benslama M., Chefrour A., Barour C. (2012). Contribution to the study of the waters of the Medjerda wadi in the far east of Algeria: Souk Ahras. Rev. Sci. Technol. Synthesis, 25: 88-96.
7. Melloul AA., Hassani L., (1999). Salmonella infection in children from the wastewater-spreading zone of Marrakesh city (Morocco). Journal of applied Microbiology, 87(4): 536-539
8. Ndiaye ML., Pfeifer HR., Niang S., Dieng Y., Tonolla M., Peduzzi R. (2010). Impacts of the use of polluted water in urban agriculture on the quality of the Dakar aquifer (Senegal). The Environmental Sciences Review, 10(2): 1-14
9. Tankari Dan-Badjo A., Yadji G., Nomaou DL., Moussa TO. (2012). Risk of exposure of the population of Niamey (Niger) to heavy metals through the consumption of market garden products, 2: 88-99.
10. Djibo H. (2016). Comparative study on three market gardening sites: case of Niamey. Revue Sciences, Langage et Communication, 1(3): 1-8
11. INS (2018). Niamey regional statistical yearbook 2013 – 2017; 93p.
12. INS (2022). General information on the region: administrative division of the Niamey region, projection based on RGP/H_2012 data.
13. Joseph-pierre, Giraud (2003). Food microbiology. Paris: Dunod, 651p (ISBN 2 10 007259 5)
14. WHO (2006). Guide lines for the safe use of wastewater, excreta and greywater: wastewater use in agriculture, vol. II. World Health Organization, France. Available at:
15. Coulibaly-Kalpy J., Koffi Kouadio S., Yéo A., Yah Kouadio L., Bamba A., Coulibaly E., Kacou-N'douba A., Dosso M. (2017). Study of the quality of wastewater from the two reservoirs of the Gourou basin at the Indénié crossroads in Abidjan. Journal of Applied Biosciences 113:11138-11144.
16. Maïwore J., Adamou M., Djaouda M., Baane MP., Amale Y., Youssouf D., Ngoune LT. (2020). Influence of some bacteriological sources of contamination on the quality of lettuce consumed in Maroua (Cameroon), identification of enterobacteria. Journal of Applied Biosciences, 154:15926–15939
17. WHO (1989). The use of wastewater in agriculture and aquaculture: Recommendations for health advice. Report of a WHO scientific group. World Health Organization, Technical Report Series 778p.
18. Abouelouafa1 M., Berrichi A., El Halouani H., Kharboua M. (2002). Effects of the reuse of raw wastewater from the city of Oujda on some agronomic and bacteriological parameters. Acts Inst. Agron. Vet. (Morocco), 22 (3):151-156.
19. El Addouil J., Chahlaoui A., Chafi A., Berrahou A. (2008). Monitoring and analysis of the risk linked to the use of wastewater in agriculture in the Meknes region of Morocco. 2IE- Sud Sciences et Technologie, 16: 29-35.

20. Adjahouinou DC., Yehouenou B., Liady MND., Fiogbe ED. (2014). Bacteriological characterization of raw wastewater from the city of Cotonou (Benin). *Journal of Applied Biosciences*. 78: 6705 – 6713
21. El Ouali LA., Zanibou A., Bekhti K., Zerrouq F., Merzouki M. (2014). Control of the microbiological quality of domestic and industrial wastewater from the city of Fez in Morocco. *J. Mater. Approximately. Sci.* 5 (1): 2325-2332.
22. AFNOR (1996). *Microbiological analyses. Volume II. Quality control of food products*, AFNOR Editions, 545p.
23. Wognin As., Ouattara Mb., Assi-Clair Bj., Koffi-Nevry R. (2022). Evaluation of the levels of bacteriological contamination of lettuce according to the production and sales sites in market gardening sites in Abidjan and peri-urban areas. *Int. J. Biol. Chem. Sci.* 16(4): 1580-1592.
24. De Roever, C. (1998). Microbiological safety evaluations and recommendations on fresh produce. *Food Control*, 6:321–347.
25. FAO (2007). *Food safety risk analysis: Guide for authorities s responsible for food safety*. FAO Food and Nutrition Study 87. Rome, ISBN 978-92-5-205604-1.
26. Wognin A.S. (2014). Contamination risk factors and virulence genes associated with *Escherichia coli* in the market gardening environment: case of lettuce (*Lactuca sativa*) in the peri-urban area of Abidjan. Doctoral thesis from Nangui Abrogoua University: microbiology and molecular biology. 122p.
27. Amoah P., Drechsel P., Abaidoo C. (2005). Irrigated urban vegetable production in Ghana: Sources of pathogen contamination and health risk elimination. *Irrig. Drainage*, 54:49-61.
28. Adjrah Y., Soncy K., Anani K., Blewussi K., Karou DS., Ameyapoh, Y., De Souza C., Gbeassor M. (2013). Socio-Economic Profile of Street Food Vendors and Microbiological Quality of Ready-to-Eat Salads in Lomé. *International Food Research Journal*, 20: 65-70.
29. Alio Sanda A., Inoussa MM., Samna Soumana O., Bakasso Y. (2017). Diversity and dynamics of *Salmonella* isolated from lettuce (*Lactuca sativa* L.) in market garden crops in Niger (West Africa). *Journal of Applied Biosciences* 119:11917-11928.
30. Geser N., Stephan R., Korczak M., Beutin L., Hachler H. (2012). Molecular identification of extended-spectrum-beta-lactamase genes from *Enterobacteriaceae* Isolated from healthy human carriers in Switzerland. *Ant. Ag. Chem.*, 56: 1609-1612.
31. Blaak H., Lynch G., Italiaander R., Hamidjaja RA., De Roda HAM. (2015). Multidrug-resistant and extended spectrum beta-lactamase-producing *Escherichia coli* in Dutch surface water and wastewater. *PLoS One*, 10(6): 0127752.
32. Obiri-Danso K., Paul N. & Jones K., 2001. The effect of UVB and temperature on the survival of natural populations and pure cultures of *Campylobacter jejuni*, *Camp. Coli* *Camplari* and urease-positive thermophilic campylobacters (UPTC) in surface waters. *Journal Applied Microbiology*, 90: 256-267.

33. Tankari Dan-Badjo A., Yajji G., Nomaou DL., Moussa TO. (2013). Risk of exposure of the population of Niamey (Niger) to heavy metals through the consumption of market garden products, 2: 88-99
34. Lennox JA., Efiuvwevwere BJO. (2012). Microbiology of freshly harvested and market retailed cucumber. *Asian Journal of Science and Technology*, 4(11): 071-073.
35. Shenge KC., Whong CMA., Yakubu LL., Omolehin RA., Erbaugh JM., Miller SA., Lejeune JT. (2015). Contamination of Tomatoes with Coliforms and *Escherichia coli* on Farms and in Markets of Northwest Nigeria. *Journal of Food Protection*, 78(1): 57–64.
36. Uyttendaele M., Jaykus LA., Amoah P., Chiodini A., Cunliffe D., Jacxsens L., Holvoet K., Lise Korsten L., Lau M., McClure P., Medema G., Sampers I. , Jasti PR. (2015). Microbial Hazards in Irrigation Water: Standards, Norms, and Testing to Manage Use of Water in Fresh Produce Primary Production. *Comprehensive Reviews in Food Science and Food Safety*, 14: 336-356.
37. Arnponsah-Doku F., Obiri-Danso K., Abaidoo R. C., Andoh L. A., Drechsel P., Kondrasen F. (2010). Bacterial contamination of lettuce and associated risk factors at production sites, markets and street food restaurants in urban and peri-urban Kumasi, Ghana. *Science Research and Essay*, 5 (2): 217-223.
38. Toe E., Dadié A., Dako E., Loukou G. Djé, K., (2017). Bacteriological quality and risk factors for contamination of raw mixed vegetable salads served in collective catering in Abidjan (Ivory Coast). *Advances in Microbiology*, 7:405-419.
39. Almou AA., Yaou C., Rabiou MM., Alio SA., Sabo HS., Sadou H. (2024). Analysis of Risk Factors for Microbial Contamination of Vegetables in Market Gardening Environments in Niamey, Niger. *Journal of Advances in Microbiology*, 24 (1): 39-51.
40. Sackou KJ., Claon JS., Oga AS., Aguessi KT., Lorougnon D., Diby Y., Kouadio K. (2006). Health quality of lettuce grown in Abidjan. *Microbiol. hyg. Power supply* 18(52): 48-50.
41. Koffi-Nevry R., Manizan NP., Wognin AS., Koussemon M., Koffi OS., Kablan T., Kakou C. (2008). Characterization of the Spatio-Temporal Distribution of Bacteria at the Water-Sediment Interface of a Tropical Lagoon: Case of Banco Bay, Abidjan, Ivory Coast. *European Journal of Scientific Research*, 21: 164-174.
42. Amoah P., Drechsel P., Henseler M., Abaidoo RC. (2007). Irrigated urban vegetable production in Ghana: microbiological contamination in farms and markets and associated consumer risk groups. *J. Water Health*, 5(3): 455-466.
43. Martínez-Vaz BM., Fink RC., Diez-Gonzalez F., Sadowsky MJ. (2014). Enteric Pathogen-Plant Interactions: Molecular Connections Leading to Colonization and Growth and Implications for Food Safety. *Microbes and Environments*, 29(2).
44. Jang H., Matthews KR. (2018). Survival and interaction of *Escherichia coli* O104:H4 on *Arabidopsis thaliana* and lettuce (*Lactuca sativa*) in comparison to *E. coli* O157:H7: Influence of plant defense response and bacterial capsular polysaccharide. *Food Research International* 108:35–41.

45. Berger CN., Sodha SV., Shaw RK., Griffin PM., Pink D., Hand P., Frankel G. (2010). Fresh fruit and vegetables as vehicles for the transmission of human pathogens. *Environmental Microbiology*, 12(9): 2385–2397.