

## Biochemical and Haematological Effects of Aqueous Leaf Extract of *Sida cordifolia* in *Plasmodium berghei* infected rat wistar albino rats

**Comment [MD1]:** italics

**Comment [MD2]:** title needs changes as given below

Biochemical and Hematological Effects of *Sida cordifolia* Leaf aqueous Extract on *Plasmodium berghei* infected wistar albino rats.

**Comment [MD3]:** ?

### Abstract

Assessment of haematological and biochemical parameters can be predictive of the adverse effects resulting from the ingestion of a foreign substance. Freshly, harvested leaves of *Sida cordifolia* were washed and dried at room temperature, after which they were ground to powder and subsequently extracted. Twenty-five adult wistar rats were divided into five groups of five rats each. Group I was the normal control and was administered with 2 ml of distilled water. Group II was infected without treatment, while Group III and IV were infected and afterwards administered with 200 and 400 mg/kg of aqueous extract of *Sida acuta* respectively. Group V was administered with the standard drug. The Packed Cell Volume reported for group II (negative) control was significantly ( $P < 0.05$ ) lower than that reported for the normal control. However, oral administration of 200 and 400 mg/kg of *Sida cordifolia* leaf extract significantly ( $P < 0.05$ ) increased it though to a level which was significantly ( $P < 0.05$ ) lower than that reported for the normal control. Similar observation was made on Haemoglobin Concentration (Hb) and Red Blood Cell (RBC). However, a contrary observation was made on the white blood cell. Urea and creatinine reported for the negative control were significantly ( $P < 0.05$ ) higher than those reported for the normal control. However, the aforementioned parameters were significantly ( $P < 0.05$ ) reduced following oral administration of the said extract. In conclusion, it can be deduced from this study that *Sida cordifolia* leaf extract has the ability to restore a distorted haematological and biochemical states resulting from *P. berghei* infection.

**Keywords:** *Sida cordifolia*, Blood, Cell, Haemoglobin, Leaf

**Comment [MD4]:** *Sida cordifolia* Leaves

### Introduction

*Sida cordifolia* a notorious specie in the genus (Malvaceae) is commonly recognized for its therapeutic potentials [1]. It is commonly known in India as Bala and mallow country one of the ingredients used in making Ayurvedic formulations [2]. *Sida cordifolia* is reportedly used as antioxidant, anticancer, and antidiabetic [3]. Other activities of *Sida cordifolia* include analgesic and anti-inflammatory activities among others [3].

**Comment [MD5]:** Which country?

**Comment [MD6]:** Which part of the plant?

**Comment [MD7]:** What are the major active constituents in *Sida cordifolia* leaves and roots?

The wrongly held impression that plant based medicinal preparations wield minimal or no side effects has undermined approach to cautious application of this category of therapy in the

treatment of diverse human diseases and its attendant consequences [4]. Medicinal plants being endowed with diverse arrays of biological compounds with great complexity tend to have several broad and actions on the physiological and biochemical systems [5].

Assessing the haematological parameters can be relied upon to diagnose unpleasant effects resulting from the usage of foreign compounds on the blood constituents of an animal [6]. Furthermore, ingestion of chemical compounds at toxic doses can orchestrate alterations in blood parameters that are suggestive of hematological disorders [7]. The role of biochemical markers in performing accurate diagnosis as well as in the assessment of risk and adoption of therapy that improve clinical outcomes [8] cannot be overemphasized thereby informing the imperativeness of this study.

## Materials and Methods

### Collection of plant sample

Mature leaves of *Sidacordifolia* were harvested from a bush in Uturu, Isikwuato Local Government of Abia State. The leaves were conveyed in a dark polythene bag to the herbarium unit of the Department of Forestry, Michael Okpara University of Agriculture Umudike, Southeastern Nigeria.

**Comment [MD8]:** Herbarium accession number , authentication report ?

### Experimental Animals

Twenty-five (25) adult mice were gotten from the animal House of the Department of Science Laboratory Technology, Akanulbiam Federal Polytechnic, UwanaAfikpo. The animals were handled in accordance with ethics governing the handling and use of laboratory animals for

**Comment [MD9]:** procured

**Comment [MD10]:** animal ethical committee sanction letter no. to be included

research purposes. They were kept in transparent plastic cages and were fed guinea grower feed for two weeks to acclimatize them before the experiment.

### Malaria parasite

Four mice passaged with the parasite *Plasmodium berghei* used for this study were bought from the Department of parasitology, University of Nigeria Nsukka.

**Comment [MD11]:** include strain identification number

### Extraction

**Comment [MD12]:** a phytochemical examination of the leaf extract is to be included .

Clean tap water was used to wash leaves of *Sidacordifolia* to get rid of dirt. The leaves were subsequently dried at room temperature after which they were ground into fine powder. Exactly 500 g of powdered plant sample was subsequently soaked in 3 L of distilled water for 24 h. The mixture was filtered with a clean sieve and was concentrated to dryness in a water bath for 3 days at 50°C.

**Comment [MD13]:** why the leaves of the plant are selected specifically?

### Acute Oral Toxicity Study

The method of the Organization for Economic Cooperation and Development (OECD 425, 2008) was used to determine the acute toxicity of extract using limit test dose of 2 g/kg. Five apparently healthy mice were which had been denied food for 4 h were dosed and subsequently weighed to determine the dose. No mortality was recorded on the first animal administered a limit dose of 2000 mg/kg. The remaining four male mice were dosed and observed for signs of toxicity such as absence of tremor, diarrhea, lethargy, and paralysis periodically for the first four and were later followed for 14 days for any lethality [9].

**Comment [MD14]:** Reference number

**Comment [MD15]:** Sentence construction need revision

### Parasite passaging

The approach of by Peter and Anatoli (1998) was relied upon to inoculate *Plasmodium berghei* (NK 65) into the mice by intraperitoneal route (Anatoli, 1998). Red blood cells infected

**Comment [MD16]:** Reference number ?

**Comment [MD17]:** Referencenumber ?

with *Plasmodium berghei* was obtained from the tail vein of the infected mice and was subsequently diluted in 5 mL of phosphate buffered saline (PBS), so that 1 mL of parasitized blood contains  $5 \times 10^9$  RBC  $m^{-1}$  infected erythrocytes, each 0.2 mL of the blood that was subsequently injected into an animal contained  $1 \times 10^6$  *Plasmodium berghei* parasitized red cells (Huang et al., 2015). Administration of extract began three (3) days after inoculation.

### Biochemical analysis

**Sample preparation:** Kidney function test was performed by introducing exactly 2 mL of blood into the EDTA tube prior to centrifugation at 4,000 rpm for 15 min and the plasma obtained was stored for biochemical analysis.

### Serum Urea Determination

Exactly 10  $\mu$ L of sample was placed into a tube containing 1000  $\mu$ L of the working reagent. The contents of the tube were thoroughly mixed, incubated for 5 minutes at 37°C (Kaplan, 1982).

Blood urea concentration was determined using the formula below:

$$Urea\ conc\ (mg/dl) = (A\ Sample) / (Acal / STD) \times conc.\ cal / STD\ (mg/dl)$$

### Haematological Evaluation

Hematological parameters (Red Blood Cells, Haemoglobin concentration and packed cell volume and white blood cell) were determined with the aid of an automatic hematological analyzer (Coulter STKS, Beckman) (Yang et al., 2019).

Comment [MD18]: Reference number?

Comment [MD19]: Formula  
Check spelling

Comment [MD20]: Reference number ?

## Data Analysis

Data obtained were expressed as Mean  $\pm$  Standard Deviation using SPSS (Ver. 23). Data were analysed using one way Analysis of Variance (ANOVA). Differences in mean were compared using Turkey Test. *p-values* less than 0.05 was considered statistically significant.

Table 1: Effect of Oral administration of Aqueous Extract of *Sidacordifolia* on the Haematological Indices of *Plasmodium berghei* Infected Rats

TREATMENT	PCV	Hb (g/dl)	WBC ( $\times 10^9/l$ )	RBC ( $\times 10^6/l$ )
Group I (Normal Ctrl)	48.05 $\pm$ 3.73 <sup>d</sup>	18.09 $\pm$ 2.42 <sup>c</sup>	6.24 $\pm$ 0.24 <sup>b</sup>	7.20 $\pm$ 2.59 <sup>d</sup>
Group II (Negative Ctrl)	28.09 $\pm$ 1.05 <sup>a</sup>	8.40 $\pm$ 1.98 <sup>a</sup>	9.82 $\pm$ 0.82 <sup>e</sup>	3.70 $\pm$ 3.05 <sup>a</sup>
Group III: INF+200 mg/kg Extract	35.02 $\pm$ 4.54 <sup>b</sup>	12.09 $\pm$ 2.33 <sup>b</sup>	8.21 $\pm$ 0.20 <sup>d</sup>	4.20 $\pm$ 3.22 <sup>ab</sup>
Group IV: INF+400 mg/kg Extract	39.09 $\pm$ 5.08 <sup>c</sup>	13.07 $\pm$ 1.82 <sup>bc</sup>	7.08 $\pm$ 2.31 <sup>c</sup>	5.20 $\pm$ 2.30 <sup>c</sup>
Group V: INF+STD	47.08 $\pm$ 3.42 <sup>d</sup>	15.07 $\pm$ 0.92 <sup>d</sup>	4.70 $\pm$ 5.30 <sup>a</sup>	7.30 $\pm$ 3.52 <sup>d</sup>

Values are expressed as mean  $\pm$  standard deviation of three determinations. Values with different superscript in a column are significantly ( $p < 0.05$ ) different

**Comment [MD21]:** Include the designation explanations for a, b, c, d along with data

Table 2: Effect of Oral Administration of Aqueous Extract of *Sidacordifolia* on the Renal Health

Treatment	Urea (mg/dl)	Creatinine (mg/dl)
Group I (Normal CTRL)	14.40 $\pm$ 5.87 <sup>a</sup>	0.81 $\pm$ 0.98 <sup>a</sup>
Group II (Negative CTRL)	31.50 $\pm$ 4.32 <sup>c</sup>	3.10 $\pm$ 0.19 <sup>c</sup>
Group III: INF+200 mg/kg Extract	20.12 $\pm$ 2.02 <sup>b</sup>	1.02 $\pm$ 0.32 <sup>b</sup>
Group IV: INF+400 mg/kg Extract	19.14 $\pm$ 0.34 <sup>b</sup>	0.96 $\pm$ 0.23 <sup>c</sup>
Group V: INF+STD (chloroquine)	14.98 $\pm$ 0.98 <sup>a</sup>	0.89 $\pm$ 0.23 <sup>a</sup>

Values are expressed as mean  $\pm$  standard deviation of three determinations. Values with different superscript in a column are significantly ( $p < 0.05$ ) different

**Comment [MD22]:** Explain all probability values with respect the data designations a, b, c, d

## RESULT AND DISCUSSION

Assessment of haematological parameters can be diagnostic of adverse effects of foreign compounds on blood constituents of animals[6]. Table 1 shows the effect of oral administration of aqueous extract of *Sidacordifolia* on the haematological indices of *Plasmodium berghei* infected rats indicating that the packed cell volume (PCV) reported for the negative control was significantly ( $p<0.05$ ) was lower than that reported for the normal control. However, oral administration of aqueous extract of *S.cordifolia* significantly ( $p<0.05$ ) increased it in a dose dependent manner. It was also observed that there was no significant ( $P>0.05$ ) difference in the PCV reported for the normal control and the standard control. The hemoglobin concentration (Hb) reported for the negative control was significantly ( $p<0.05$ ) lower than that reported for other groups. However, following oral administration of extract, there was a significant increase in the Hb reported for groups II and III which though were not significantly ( $p>0.05$ ) different from each other but lower than that reported for the normal control. The red blood cell (RBC) reported for the negative control was significantly ( $p<0.05$ ) lower than those reported for other groups. The RBC reported for group III was not significantly ( $p>0.05$ ) different from that reported for the negative control. Similar observation was made between the normal control and the standard control. The anti-anemic effect of *S.cordifolia* extract could be attributed to antioxidant activity of the plant extract which had been reported in previous study. This is consistent with the finding of Ukpanukpong et al. [12] which showed that ethanolic leaf extract of *S.rhobofolia*, a member of the *malvaceae* family to which *S. cordifolia* belongs orally administered on rats induced with artificial infertility significantly ( $P<0.05$ ) increased the red blood cell. The White Blood Cell (WBC) reported for the negative control was significantly ( $p<0.05$ ) higher than that reported for the normal control which in turn was significantly

**Comment [MD23]:** What is present in the aqueous extract of the leaves ?

**Comment [MD24]:** Sentence needs grammar revision

**Comment [MD25]:** ???

( $p < 0.05$ ) lower than those reported for groups III and IV which were infected and treated with extract. This is contrary to the finding made by Ukpanukpong et al. [12] which showed that oral administration of 200 and 400 mg/kg of *S. rhombifolia* extract significantly increased the white blood cell in rats induced with artificial infertility. Biochemical markers play an important role in accurate diagnosis and also for assessing risk and adopting therapy that improve clinical outcome [8]. Furthermore, creatinine is commonly used as a measure of kidney function [13]. In addition, the most frequently determined clinical indices for estimating renal function depends upon the concentration of urea in the serum [14]. The kidney is one of the organs that receive the bulk of the toxic impacts of toxic substances. Table 2 shows the effect of oral administration of aqueous extract of *Sida cordifolia* on the renal health of rats infected with *Plasmodium berghei* indicating that serum creatinine and urea were significantly ( $p < 0.05$ ) higher in the negative control group which were infected without treatment than those reported for the normal control group. These were significantly ( $p < 0.05$ ) reduced in a dose dependent manner. This is consistent with the finding of Ukpanukpong et al. (2019) which showed that ethanolic leaf extract of *Sida cordifolia* was not toxic.

#### Conclusions

It has been revealed through this study that aqueous leaf extract of *Sida cordifolia* has the potential to reverse distorted haematological and biochemical status of rats previously infected with *P. berghei*.

## References

Dinda B, Das N, Dinda S, Dinda M, SilSarma I. The genus *Sida* L.–A traditional medicine: It is ethnopharmacological, phytochemical, and pharmacological data for commercial exploitation in the herbal drugs industry. *Journal of Ethnopharmacology*. 2015;176:135-76

Ravi KV and Diwan P. Analgesic, antiinflammatory and hypoglycaemic activities of *Sida cordifolia*. *Phytotherapy Research*. 1999;13(1):75-7.

Rodrigues FC, de Oliveira AFM. The genus *Sida* L.(Malvaceae): An update of its ethnomedicinal use, pharmacology, and phytochemistry. *South African Journal Botany*. 2020;132:432-62.

Philomena G. Concerns regarding the safety and toxicity of medicinal plants - An overview. *J Applied Pharmaceutical Science*. 2011; 01(6): 40-44.

Kazemipoor M, Radzi CW, Cordell GA, Yaze I. Safety, efficacy and metabolism of traditional medicinal plants in the management of obesity: a review. *International journal of chemical engineering*. 2012;3(4):288-292.

Ashafa, A.O., Orekoya, L.O., Yakubu, M.T. Toxicity profile of ethanolic extract of *Azadirachta indica* stem bark in male Wistar rats. *Asian Pacific Journal of Tropical Biomedicine*, 2012;2:811-817.

Price EA, Schrier SL Anemia in the elderly: introduction. *Seminar Hematology*, 2008;45: 207-209

Organisation for Economic Co-operation and Development. OECD guidelines for the testing of chemicals. Acute oral toxicity

Kaplan, A and Teng, L.L. Selected Methods of Clinical Chemistry. In W.R. Faulkner & Mietes, S (Eds.), AACC, Washington. 1982; 357-363.

Yang M, Wu Z, Wang Y, Kai G, S, Njateng, G. S, Cai, S., et al. Acute and Subacute Toxicity Evaluation of Ethanol Extract from Aerial Parts of *Epigynum Auritum* in Mice. *Food Chemical Toxicology*, 131;2019;(1), 110534.

Ukpanukpong RU, Uyabeme RN, Adebisi, F.D. D.I. Basiru, B.S. Akinfeso and P.O. Aigbadumah. Hematological Studies on *Sida Rhombifolia* Ethanolic Leaf Extract of Micronor Induced Infertility in Female Rats. *International Journal of Current Microbiology and Applied Sciences*. 2019; 8(3): 1751-1759,

Corbett JV. (2008). Laboratory tests and diagnostic procedures with nursing diagnoses. 7th Ed. pp.90-107.

**Comment [MD26]:** Referencing pattern does not follow any style uniformly, it should be revised

**Comment [MD27]:** Following alphabetical order or numbers ?? clarify with text

**Comment [MD28]:** Year, document number?

Branten AJ, Vervoort G, Wetzels JF. Serum creatinine is a poor marker of GFR in nephrotic syndrome. *Nephrol Dial Transplant* 2005; 20: 707–711.

UNDER PEER REVIEW

