

Effect of NPK through chemical fertilizers combination with FYM and bio-fertilizers on quality attributes of onion in Jaipur region of Rajasthan

Abstract

This field experiment was conducted in loamy sand soils of the Horticulture farm, Vivekanand Global University Jaipur, during *rabi* season of 2020-21 and 2021-22. The experiment, comprising of 32 treatment combinations replicated three times, was laid out in split-plot design with four fertility levels of NPK (0,50, 75 and 100% of recommended dose of NPK) and two levels of FYM (without FYM and with FYM @ 25 t ha⁻¹ were added in main plots. Four bio-fertilizer levels (No inoculation N₂ fixer *Azotobacter*, PSB inoculation and N₂ fixer *Azotobacter* + PSB inoculation) were added in sub plots. The results of the study have clearly shown that application of chemical fertilizer up to 100% RDF increased all the quality attributes (pungency, chlorophyll content and TSS), N,P,K and S content and uptake and availability of N, P and K in soil after harvest the crop. With application of FYM @ 25 t ha⁻¹ significantly improved all quality attributes (Allyl propyl disulphide, chlorophyll and TSS), N, P, K and S content and uptake and availability of N, P and S in soil after harvest the crops. Use of biofertilizers (N₂ fixers and PSB) alone or in combination increased all the quality attributes (Allyl propyl disulphide, chlorophyll and TSS content), NPK and S content and uptake and availability of N, P and K in soil after harvest the crop.

Comment [Vk1]: SHOULD BE IN ITALICS and R should be in capitals

Comment [Vk2]: Put it in Italics and check throughout the MS check as per journal format

Keywords: FYM, Bio-fertilizer, Allyl propyl disulphide, chlorophyll and TSS content

Introduction

Onion (*Allium cepa* L.) is one of the most important commercial vegetable crops cultivated extensively in India and it is native of the Central Asia and Mediterranean region and commercially grown in China, India, U.S.A., erstwhile U.S.S.R., Japan, Spain, Turkey, Brazil, Egypt etc it belongs to family Alliaceae. Onion is considered to be the second most important vegetable crop grown in the world after tomato [1]. Onion is liked for its flavor and pungency in onion is due to presence of a volatile oil 'allyl propyl disulphide'- organic compound rich in Sulphur [2]. The beneficial compound called 'quercetin' present in onion is a powerful antioxidant [3].

In Rajasthan, onion occupies an area 64.76 thousand ha with the production 996.73 thousand ton and productivity 15.39 MT/ha [4]. In Rajasthan, major onion producing districts are Jodhpur, Sikar, Nagaur, Alwar, Jaipur etc. It is a unique vegetable that is used throughout the year in the form of the salad or condiment or for cooking with other vegetables. Onion is also used in preparing soups, sauces, curries, pickles and for flavouring or seasoning food [5].

Further, knowing the deleterious effect of using only chemical fertilizers on soil health, use of chemical fertilizers supplemented with organic waste and biofertilizers will be environmentally benign. Therefore, biofertilizers are widely accepted as low-cost supplements to chemical fertilizers with no deleterious effect either on soil health or environment Bhagyaraj and Suvarna [6]. Availability of nitrogen is important for growing plants as it is major indispensable constituent of protein and nucleic acid. The application of nitrogen with different doses increased plant growth and yield of onion [7]. Phosphorus has its beneficial effect on early root development, plant growth, yield and quality. Potassium plays an important role in crop productivity [8]. Keeping in view the significance of above aspects, the main objectives of study is to work out the possibility of reducing the use of chemical fertilizers by application of organic manures and bio-fertilizers.

FYM is the commonly used organic manure but its supply is limited. It contains low and widely varying nutrient concentration. Combined use of FYM and inorganic fertilizers is of special significance under intensive cropping system as these are complementary and supplementary to each other in sustaining crop yields and soil productivity [9]. Biofertilizers are the inoculation of microorganism, which are capable of mobilizing nutrient element from unavailable to available form through biological processes [10]. These are biologically active strains or products containing active form of microorganisms. These may be helpful for increasing the crop production by way of enhancing the soil fertility. Use of biofertilizers not only supplement the nutrient but also improve the efficiency of applied nutrients [11]. Studies have also shown that integrated use of chemical fertilizers, organic residues such as FYM, compost etc. and biofertilizers resulted in reduced losses of nutrients and environmental pollution [12]. Balanced fertilization has to be made for different crops based on soil testing for attaining maximum yield and profit. There is meager information on the balanced use of chemical fertilizers with FYM and biofertilizers for onion crop grown in Rajasthan. So in last few years, a greater concern regarding use of biofertilizers and organic sources as an alternative/supplement to chemical fertilization has been derived to reduce the high cost that inorganic fertilizers represent in agricultural production.

Application of fertilizers is imperative to maintain the desired pace of crop production. It needs to be used with caution by alleviating consequent hazards in soil with the help of supplemental organic manuring. Continuous use of inorganic fertilizers has depleted soil organic matter, resulting into inherent loss of native soil N, available P, available K and ultimately lowered the productions [13]. The repeated and excessive application of chemical fertilizers is harmful to the plants. Microflora of the soil, which is extremely essential for maintaining biological health, texture and structure of soil, also gets affected [14].

Method and Materials

The present experiment was conducted at Horticulture farm, Vivekanand Global University, Jaipur, during *Rabi* season of 2020-21 and 2021-22 was laid out in split-plot design in both the years with 32 treatment combinations replicated thrice. Jobner is situated at 26.5° North latitude, 75.20° East longitude and an altitude of 427 meters above mean sea level, in Jaipur district of Rajasthan. This region falls under Agro-climatic zone IIIA (Semi-Arid Eastern Plain) of the state. The climate of Jobner is typically semi-arid characterized by extremes of temperature in both summer and winter with low rainfall and moderate relative humidity. Maximum temperature in summers is as high as 45 °C and minimum temperature in winters fall around 0 °C. To find out the effect of conjoint use of NPK through chemical fertilizers with FYM and bio-fertilizers on growth and yield of onion. The soil of experimental field was loamy sand in texture, slightly alkaline in reaction, poor in organic carbon with low available nitrogen, phosphorus and sulphur and medium in potassium status. The treatments comprised of Chemical fertilizers, FYM and biofertilizers with ten treatments *Viz.*, F₀- Control, F₁-50% of recommended dose of NPK, F₂-75% of recommended dose of NPK, F₃-100% of recommended dose of NPK, M₀- Without FYM, M₁- With FYM @ 25 t ha⁻¹, B₀- No inoculation, B₁- N₂ fixer *Azotobacter*, B₂- PSB inoculation and B₃- N₂ Fixer *Azotobacter* + PSB inoculation, respectively.

The treatments of chemical fertilizers, FYM, and bio-fertilizers were applied as per treatment in respective plot. The spacing 15 cm row to row and 10 cm plant to plant was maintained. The seedlings were transplanted in cool evening according to the layout plan. A light irrigation was applied just after the transplanting and subsequent irrigation was given at an interval of 10-12 days depending upon the soil condition. Harvesting of onion was done on last week of May, 2020 and 2021.

Comment [Vk3]: Give numbering as per journal format

Comment [Vk4]: In heading you have mentioned only about quality

Result and Discussion

Comment [Vk5]: numbering

Quality attributes

Data pertaining to pungency for both study years and their pooled mean presented in table 1 that different level of fertility significantly influenced the pungency in bulb. Progressive increase in level of fertility from control to 100 % RDF brought about significant improvement in pungency in onion bulb over preceding levels. Application of 100% RDF attained the maximum mean pungency (6.34 mg/100 g) among all the treatment and it registered 16.13, 8.94 and 3.85 per cent increase in pungency over control, 50 and 75 % RDF, respectively. Data given in same table also indicating that application of FYM @ 25 t ha⁻¹ significantly increased the pungency in onion bulb to the extent of 5.58 per cent over no FYM application in pooled analysis. It is also evident from the data (Table 1.) that use of biofertilizers viz., *Azotobacter* and PSB or in combination significantly increased the pungency. Use of *Azotobacter* and PSB increased the pungency to the tune of 4.40 and 3.61 per cent over no inoculation, respectively. Combined application of *Azotobacter* and PSB represented the maximum increase of 8.97 per cent over control.

Comment [Vk6]: T should be in capital letter

Table 1: Effect of NPK, FYM and bio-fertilizers on Pungency (Allyl Propyl Disulphide mg/100 g).

Comment [Vk7]: Better to mention crop name

Treatment	Pungency (Allyl Propyl Disulphide mg/100g)		
	2021-22	2022-23	Pooled mean
Fertilizers			
F ₀ = Control	5.50	5.41	5.46
F ₁ = 50 % RDF	5.86	5.77	5.82
F ₂ = 75 % RDF	6.15	6.05	6.10
F ₃ = 100 % RDF	6.38	6.29	6.34
SEm _±	0.12	0.09	0.07
CD(p=0.05)	0.35	0.28	0.22
FYM			
M ₀ = Control	5.82	5.69	5.76
M ₁ = 25 t ha ⁻¹	6.12	6.07	6.10
SEm _±	0.08	0.07	0.05
CD(p=0.05)	0.24	0.20	0.15
Biofertilizers			
B ₀ = Control	5.73	5.64	5.69
B ₁ = <i>Azotobacter</i>	5.98	5.89	5.94
B ₂ = PSB	5.92	5.86	5.89

Comment [Vk8]: Check Alignment

B ₃ = Azotobacter + PSB	6.26	6.13	6.20
SEm±	0.10	0.09	0.07
CD (p=0.05)	0.29	0.26	0.19

It is evident from table 2 that different levels of fertility significantly affected chlorophyll content of leaves during both the years and in pooled analysis. The mean maximum chlorophyll content (0.967 mg/g fresh weight of leaves) was observed with 100% RDF followed by 75% RDF (0.965 mg/g fresh weight of leaves) but these were at par with each other. Minimum chlorophyll content was recorded in control (0.942 mg/g fresh weight of leaves), which was 2.65, 1.15 and 0.26 percent less as compared to 50, 75 and 100% RDF, respectively in pooled analysis. Application of FYM also had significant effect on chlorophyll content in both the study years as well as in pooled analysis. Application of FYM @ 25 t ha⁻¹ increased the chlorophyll content by 0.78 per cent during individual's years and pooled analysis respectively, over control. Data from table 2 further showed that use of biofertilizers alone or in combination significantly increased the chlorophyll content of leaves. Use of *Azotobacter* and PSB increased the chlorophyll content with the use of 1.37 and 1.11 percent respectively over control. Application of *Azotobacter* and PSB in combination represented the maximum increase of 2.38 per cent over no inoculation in pooled analysis.

Comment [Vk9]: Check font style

Table 2: Effect of NPK, FYM and bio-fertilizers on chlorophyll content (mg/g fresh weight of leaves)

Treatment	Chlorophyll content (mg/g fresh weight of leaves)		
	2021-22	2022-23	Pooled mean
Fertilizers			
F ₀ = Control	0.945	0.939	0.942
F ₁ = 50 % RDF	0.958	0.954	0.956
F ₂ = 75 % RDF	0.967	0.962	0.965
F ₃ = 100 % RDF	0.970	0.964	0.967
SEm±	0.001	0.001	0.001
CD (p=0.05)	0.004	0.004	0.003
FYM			
M ₀ = Control	0.957	0.950	0.954
M ₁ = 25 t ha ⁻¹	0.963	0.959	0.961
SEm±	0.001	0.001	0.001
CD (p=0.05)	0.003	0.003	0.002

Comment [Vk10]: Is it correct unit

Biofertilizers			
B ₀ = Control	0.949	0.943	0.946
B ₁ = Azotobacter	0.962	0.956	0.959
B ₂ = PSB	0.959	0.954	0.957
B ₃ = Azotobacter + PSB	0.971	0.966	0.969
SEm+	0.001	0.001	0.001
CD (p=0.05)	0.004	0.004	0.002

Data pertaining to the effect of integrated nutrient management or total soluble solids of bulb are presented in table 3. Application of different levels of fertility had significant influence on TSS% in both year of experiment as well as in the pooled analysis. The mean maximum TSS (12.88%) was recorded in 100% RDF treatment as followed by 75% RDF (12.71 %). However, it was found at par with each other.

Table 3: Effect of NPK, FYM and bio-fertilizers on total soluble solids of bulb

Treatment	Total soluble solids (%)		
	2021-22	2022-23	Pooledmean
Fertilizers			
F ₀ =Control	12.22	12.15	12.18
F ₁ =50 %RDF	12.55	12.47	12.51
F ₂ =75 %RDF	12.74	12.68	12.71
F ₃ =100 %RDF	12.90	12.86	12.88
SEm±	0.15	0.16	0.11
CD(p=0.05)	0.43	0.47	0.31
FYM			
M ₀ =Control	12.31	12.25	12.28
M ₁ =25 t ha ⁻¹	12.89	12.82	12.85
SEm±	0.10	0.11	0.08
CD(p=0.05)	0.31	0.33	0.22
Biofertilizers			
B ₀ = Control	12.10	12.06	12.08
B ₁ = Azotobacter	12.68	12.64	12.66
B ₂ = PSB	12.66	12.63	12.65
B ₃ = Azotobacter + PSB	12.96	12.82	12.89
SEm+	0.13	0.13	0.09
CD (p=0.05)	0.35	0.36	0.25

The application of 100 % RDF registered an increase of 5.71, 2.96 and 1.34 per cent over control, 50 and 75% RDF, respectively in pooled analysis. TSS % of bulb was also significantly affected by application of FYM @ 25 t ha⁻¹ in comparison to control. It

registered 4.44 percent more TSS per cent over no application of FYM. Data from table 3 further indicate that inoculation of biofertilizers had significant effect on TSS % of bulb in both the years as well as in pooled analysis. The maximum mean of TSS % in bulb (12.89%) was recorded with the combined use of *Azotobacter* and PSB which led to an increase of 6.71, 1.82 and 1.94 per cent over control, *Azotobacter* and PSB, respectively. However, the treatment with *Azotobacter* and PSB were at par with each other in pooled analysis. The similar results have also been reported by Sharma et al. [16], Soni [17], Pandey et al. [15], Vachhani and Patel [20], Sreenivas et al. [18], Sharma et al. [14], Thilakavathy and Ramaswamy [19].

Comment [Vk11]: Reasoning part has been missed please mention some spectacular reasons

Conclusion

Maximum TSS in bulb, allyl propyl disulphide content in bulb chlorophyll content in leaves, n content and uptake, P content and uptake, K content and uptake, S content and uptake in onion were recorded with 100% RDF levels of fertility. The maximum available N, P and K in soil after harvest were also recorded in same level of fertility. With the application of FYM @ 25 t ha⁻¹ has exhibited maximum TSS, chlorophyll content, pungency content, n content and uptake, p content and uptake, k content and uptake by onion crop. Available NPK content in soil after harvest was maximum in m1 (FYM @ 25 t ha⁻¹) treatment. Maximum TSS in bulb, allyl propyl disulphide content in bulb chlorophyll content in leaves, NPK and s content in onion bulb as well as in leaves, NPK and s uptake by onion crop was recorded under f3 combined application of *Azotobacter* and PSB.

References

1. Gulati A, Wardhan H, Sharma P. Tomato, Onion and Potato (TOP) value chains. Agricultural value chains in India. 2022:33.
2. Sharma K, Almulla S, Mahato N, Pradhan S, Mateen S, Pashikanti S. 16 Metabolomics of Allium. Edible Alliums: Botany, Production and Uses. 2022:270.
3. Črnivec IG, Skrt M, Šeremet D, Sterniša M, Farčnik D, Štrumbelj E, Poljanšek A, Cebin N, Pogačnik L, Možina SS, Humar M. Waste streams in onion production: Bioactive compounds, quercetin and use of antimicrobial and antioxidative properties. Waste management. 2021;126:476-86.
4. Anonymous 2018. Indian Horticulture database, Ministry of Agriculture, Government of India.

5. Ramkumar TR, Karuppusamy S. Plant diversity and ethnobotanical knowledge of spices and condiments. *Bioprospecting of Plant Biodiversity for Industrial Molecules*. 2021;19:231-60.
6. Bhagyaraj DJ, Suvarna CV. Biofertilizers and their application. *Agri east*, 1999; 1:16-21.
7. Patel KP, Patel JC, Patel BS, Sadaria SG. Yield and nutrient uptake by onion (*Allium cepa*) as influenced by irrigation, Nitrogen and phosphorus, *Indian Journal of Agronomy*. 1992;37:395-396.
8. Ali MM, Petropoulos SA, Selim DA, Elbagory M, Othman MM, Omara AE, Mohamed MH. Plant growth, yield and quality of potato crop in relation to potassium fertilization. *Agronomy*. 2021;11(4):675.
9. Abid M, Batool T, Siddique G, Ali S, Binyamin R, Shahid MJ, Rizwan M, Alsahli AA, Alyemini MN. Integrated nutrient management enhances soil quality and crop productivity in maize-based cropping system. *Sustainability*. 2020;12(23):10214.
10. Singh D, Thapa S, Geat N, Mehriya ML, Rajawat MV. Biofertilizers: Mechanisms and application. In *Biofertilizers 2021 Jan 1* (pp. 151-166). Woodhead Publishing.
11. Bayu T. Review on contribution of integrated soil fertility management for climate change mitigation and agricultural sustainability. *Cogent Environmental Science*. 2020;6(1):1823631.
12. Sharma T, Arya VM, Sharma V, Sharma J, Gulshan T, Bera A, Laskar YA. Integrated nutrient management: A long-term approach towards sustainability. *International Journal of Plant & Soil Science*. 2022;34(20):433-46.
13. Das D, Sahoo J, Raza MB, Barman M, Das R. Ongoing soil potassium depletion under intensive cropping in India and probable mitigation strategies. A review. *Agronomy for Sustainable Development*. 2022;42(1):4.
14. Sharma OL, Katole NS and Gautam KM. Effect of irrigation schedules and nitrogen levels on bulbs yield and water use by onion. *Agric. Sci. Digest-Karnal*. 1994;14: 15-8.
15. Pandey UB, Singh L, Singh RK, Singh L, Kumar R and Chaudhary SP. Response of different levels of N, P and K on the yield and quality of *kharif* onion. *Recent advances in medicinal aromatic and spice crops*. Vol. 1 International conference held on 28-31 Jan. 1989 at New Delhi 231-234.
16. Sharma RP, Datt N and Sharma P. Combined application of nitrogen, phosphorus, potassium and farm yard manure in onion under high hill, dry temperature conditions

of North Western Himalayan. *Indian J. Agril. Sci.*, 2003;73 (4) : 225-227.

17. Soni AK. Effect of crop geometry and fertility levels on growth, yield and quality of *kharif* onion (*Allium cepa* L.) cv. N- 53 in semiarid conditions. . Ph. D. Thesis, RAU, Bikaner. 2005.
18. Sreenivas CH, Muralidhar S and Singh, M. Yield and quality of ridge gourd fruits as influenced by different levels of inorganic fertilizers and vermicompost. *Ann. Agric Res.* 2000;2(1):262-266.
19. Thilakavathy S and Ramaswamy N. Effect of inorganic and bio fertilizers treatments on yield and quality parameters of multiplier onion (*Allium cepa* L. var *aggregatum*). News Letter. Nat. Hort. Res. Devt. Foundation. 1998;18 (2): 18-20.
20. Vachhani MU and Patel ZG. Effect of nitrogen, phosphorus and potash on bulb yield and quality of onion (*Allium cepa*). *Indian J. Agronomy.* 1993; 38 (2): 333-334.

Comment [Vk12]: Check as per journal format

UNDER PEER REVIEW