

## **Productivity and economic feasibility of pigeonpea base companion cropping under additive series planting system.**

Comment [1]: Title can be concised more

### **Abstract**

An experiment was carried out during two consecutive *Kharif* seasons of 2016-2017 and 2017-18 at Soil Conservation and Water Management Farm, C.S. Azad University of Agriculture and Technology, Kanpur. The main objective was to work out the yields of main crop of pigeonpea, black gram and sesame and their economic in term of net profit. The soil of the experimental field was sandy loam, having pH 7.8, organic carbon 0.32%, available P<sub>2</sub>O<sub>5</sub> 17.3 kg/ha and available K<sub>2</sub>O 181.2 kg/ha therefore, the fertility status was medium. The five treatments of companion cropping i.e. pigeonpea sole, black sole, sesame sole, pigeonpea + black gram (1+1) additive series and pigeonpea + sesame (1+1) additive series. The total productivity of pigeon pea + black gram was highest (23.53 q/ha) in comparison to pigeonpea + sesame (20.30 q/ha) and pigeonpea alone (17.34 q/ha). The pooled data display that maximum cost of cultivation Rs. 57876.00/ha observed under pigeonpea + black gram companion cropping followed by pigeonpea + sesame (Rs. 56191/ha). The highest gross return Rs. 134024/ha, net return Rs. 76148/ha and BCR 1:2.32 were also observed with pigeonpea + black gram (1+1) additive series followed by pigeonpea + sesame (1+1) additive series, while pigeonpea alone gave net return Rs. 50387/ha and BCR 1:2.01.

Comment [2]: Kg ha<sup>-1</sup>

Comment [3]: ?????

Comment [4]: Delete gap in both sides in all the intercropping system

Comment [5]: No gap

Comment [6]: Unit should be q ha<sup>-1</sup> in all cases of the texts

Comment [7]: In two years data pooled data can't be done

Comment [8]: Unit should be written properly

Comment [9]: ??????

**Keywords:** BCR, Companion cropping, Net income, Pigeonpea base, Pooled data.

Comment [10]: Not sound good

### **Introduction**

Pigeonpea (*Cajanus cajan* L.) is an important grain legume crop of the semi-arid tropics cropping, about the area and production with 35% and 29% world, respectively available in this tract. Further, pigeonpea occupied second position after chickpea and contributed 4.2 mt. from an area of 4.43 mt. with average productivity of 960 kg/ha. In Uttar Pradesh, pigeonpea covered 2.85 lakh hectares and total production is 2.72 lakh tones during 2020 (Anonymous, 2021). Pigeonpea grown as single crop display in efficient utilization of resources especially the space because of its slow initial growth rate, therefore, cultivation of pigeonpea as a sale crop is reported less profitable due to higher

Comment [11]: Unit??????? for area

Comment [12]: ????

duration and wider spacing (Sekhon *et al.* 2018). To make the cultivation of pigeonpea more viable, it is necessary to utilize the inter row space through companion cropping. Intercropping with short duration pulse like black gram, green gram, pearl millet like finger millet, foxtail millet in pigeonpea may enhance total productivity and may also provide early cash flow. Companion cropping also suppress the weed growth in between two rows of pigeonpea. Efficient of resource utilization can be increased by companion cropping and it also help in harnessing benefits of positive interaction in crop association. Growing a intercrop which produces the maximum cover, reduces soil loss, black gram is important cover crop for rainy season. The crop gives early and dense ground cover, which generally co-occurrence with peak rate of runoff.

Comment [13]: Why like?

In areas where the annual rainfall is in the range of 600-850 mm companion cropping is being recommended and practiced. The different intercropping systems including sesame or oilseeds are followed in different parts of country. Sesame and pulse crop like black gram which are drought resistant and have a capacity to low level management conditions may be ideal combination for sustainable cropping system (Yadav *et al.*, 2013). Companion cropping is practiced as an insurance of crop failure under rainfed condition. The main objective is higher productivity per unit area in addition to stability in production. Companion cropping system utilizes resource efficiency and their productivity is increased.

Comment [14]: No reference cited

The economic resources of rainfed area are the national treasure and need proper planning to make best use of them. Therefore, sustainable management practices are urgently needed all over the world to preserve the production potential of agricultural land. Efficient management and maintenance of soil health is the being to accomplish sustainable high productivity, food security and environment safety (Kumar *et al.*, 2017).

Therefore, productivity and profitability increases through companion cropping system is the subject matter of this manuscript.

Comment [15]: There is no mention about additive series

## Materials and Methods

An experiment was conducted under rainfed condition during two consecutive Kharif seasons of 2016-17 and 2017-18 at Soil Conservation and Water Management Farm, C.S. Azad University of Agriculture and Technology,

Kanpur. The five treatments were tested i.e., pigeonpea sole planted at 60 cm apart, black gram sole planted at 30 cm apart, sesame sole planted at 30 cm apart, pigeonpea + black gram (1+1) planted under additive series and pigeonpea + sesame (1+1) planted under additive series. The main crop was pigeonpea, while black gram and sesame were planted in the inter spaces of two rows of pigeonpea. The soil of experimental site was a typic eroded Gangetic alluvial representing Kanpur Type-1. The soil was sandy loam, having pH 7.7, organic carbon 0.33%, available N 172.00 kg/ha, available P<sub>2</sub>O<sub>5</sub> 17.50 kg/ha and available K<sub>2</sub>O 183.00 kg/ha, therefore fertility status of plant nutrients was medium. The pH was determined by Electrometric glass electrode method as discussed by Piper (1966). The organic matter was analyzed by Walkley and Black's rapid titration method (Walkley and Black, 1934). The available P<sub>2</sub>O<sub>5</sub> and available K<sub>2</sub>O were analyzed by Olsen's method and Flame photometric method, respectively, (Muhr *et al.*, 1963). The available N was analyzed by Kjeldahl's method as suggested by Subhiah and Asija (1956). The cultivar Amar (KA-32-1) of pigeonpea, cv. Shekhar-2 (KU-300) of black gram and cv. Shekhar (SH-446) of sesame were shown under intercropping. The sole crops of pigeonpea and black gram were fertilized with 20 kg N + 40 kg P<sub>2</sub>O<sub>5</sub>/ha, while sole and intercropped sesame were fertilized with 40 kg N + 20 kg P<sub>2</sub>O<sub>5</sub> + 20 kg K<sub>2</sub>O/ha. The recommended methods of fertilizer application were applied for obtaining better response. The recommended conservation agronomical practices were followed for raising of experimental crops during two experimental seasons. The harvesting of crops was made at complete maturity stage. The treatments were replicated thrice in a split plot design. The obtained data analyzed by standard method as suggested by Cochran and Cox (1963). The economics computed and pooled to draw valid conclusion from the study, is the subject matter of this study.

### Results and Discussion

The pooled data on seed yield of main and intercrops and economic studies are presented in Table-1 and discussed here under appropriate heads.

- (A) **Total productivity:** Purusal of data available in Table1 make it clear that the highest total productivity was weighed under pigeonpea + black gram intercropping system by 23.53 q/ha, followed by inter cropping of

**Comment [16]:** Correction of presentation of unit should be maintained

**Comment [17]:** ?????

pigeonpea + sesame 20.30 q/ha. The order of performance of alone cropping system was pigeonpea alone (17.34 q/ha) > black gram alone (9.44 q/ha) and > sesame alone (6.32 q/ha).

(B) **Economic Study:** The maximum cost of cultivation was observed under pigeonpea + black gram (1+1) additive intercropping by Rs. 57876/ha. It might be attributed to total population adjustment under both component crops of the system and their total input requirements. The highest gross return (Rs. 134024/ha), net return (Rs. 76148/ha) and BCR (1:2.32) were observed under pigeonpea + black gram intercropping followed by pigeonpea + sesame (1+1) additive intercropping. It may be due to higher total productivity of the planting system. These results are similar, with the findings of Reddy *et al.* (2007), Dudhade *et al.* (2009), Sharma *et al.* (2012) and Kumawat *et al.* (2013).

#### **Conclusion and recommendation**

Since, the cropping system of pigeonpea + black gram gave net return by Rs. 76148/ha and pigeonpea + sesame gave net profit by Rs. 54862/ha, therefore, farm families residing in the vicinity of rainfed area may be advocated for adoption of intercropping of pigeonpea + black gram and pigeonpea + sesame for higher total productivity and profitability and harvest the fruits of newly generated technology.

#### **References:**

- Anonymous, 2021.** Kharif keeSaghanPadhatiyen. Publication, Directorate of Agriculture, U.P. Lucknow : 229.
- Cochran, W.G. and Cox, G.M. 1963.** Experimental Design. Asia Publication House, New Delhi.
- Dudhade, D.D., Deshmukh, G.P., Harer, P.N. and Patil, J.V. 2009.** Studies on intercropping of pulse crops with pigeonpea under rainfed condition. *Legume Research*, 32(3):215-217.
- Kumar, S.N.A., Gowda, J., Venkate, Porama, V.R., Ramkrishna and Satish, A. 2017.** Qualifying and mapping of soil nutrients in Bantanahalli micro watershed using GIS and GPS. *Journal of Soil and Water Conservation*, 16(1): 18-24.

- Kumawat N., Singh, R.P., Kumar, R., Kumari, Anupma and Kumar, P. 2013.** Response of intercropping and integrated nutrition on production potential and profitability on rainfed pigeonpea. *Journal of Agricultural Science (Toronto)*, 4(7):154-162.
- Muhr, G.R., Datta, N.P., Sankarasubramoney, H, Diver, R.F., Leley, V.K. and Donadhue, R.L. 1963.** Soil Testing in India, U.S. Agency for International Development Mission to India, New Delhi.
- Piper, C.S. 1966.** Soil and Plant Analysis. Hans Publication, Bombay.
- Reddy, M.M., Padmaja, B. and Rao, L.J. 2007.** Performance of pigeonpea cultivars in intercropping system under rainfed conditions in vertisols. *Indian Journal of Dryland Agricultural Research and Development*, 22(2):208-212.
- Sharma, A.Rathod, P.S., Dharmraj, P.S. and Mohan Charan, 2012.** Response of pigeonpea to bio-fertilizers in pigeonpea based intercropping systems under rainfed conditions. *Karnataka Journal of Agricultural Sciences*. 25(3):322-325.
- Sekhon, F.S., Singh, T. and Singh, S. 2018.** Growth phenology and yield of pigeonpea (*Cajanus cajan*) as affected by intercropping systems and application of nutrients level to intercrop. *Indian Journal of Agricultural Science*, 88(3):509.
- Subhiah, B.V. and Asija, C.L. 1956.** A rapid procedure for estimation of available nitrogen in soil. *Current Science*, 25:259-260.
- Wakley, A.J. and Black, I.A. 1934.** Estimation of organic carbon by Chromic acid titration method. *Soil Science*, 37:29-38.

**Table-1:** Yield and Economy of pigeonpea and intercrops under different treatments.(Pooled data of two years)

S.N.	Cropping system	Yield (q/ha)			Cost of cultivation (Rs./ha)	Gross return (Rs./ha)	Net return (Rs./ha)	BCR
		Main crop	Inter crop	Total productivity				
1.	Pigeonpea alone	17.34	-	17.34	49861	100248	50387	2.01
2.	Black gram alone	09.44	-	09.44	32966	51025	18059	1.54
3.	Sesame alone	06.32	-	06.32	31561	34056	2495	1.07
4.	Pigeonpea + black gram (1+1) a.s.	19.34	4.19	23.53	57876	134024	76148	2.32
5.	Pigeonpea + sesame (1+1) a.s.	16.61	3.69	20.30	56191	116053	54862	2.07

UNDER PEER REVIEW

