

# ROLE OF SULPHUR NUTRITION IN OILSEED CROPS:A REVIEW

## Abstract

India is a significant importer of edible oils and an important producer of oilseed. India stands as fourth leading oilseeds producing countries after USA, China and Brazil. Oilseed has become one of the most valuable agricultural commodities. Sulphur (S) plays an important role in determining the quality of oil seed from these crops. It is referred to as the fourth major nutrient after N, P and K because crops need sulphur slightly less than they require. It is required for the formation of proteins, vitamins and chlorophyll. It is an integral part of amino acid such as cystine, cysteine and methionine which are essential components of protein. Plant can absorb sulphur in the form of sulphate ( $\text{SO}_4^{2-}$ ) which is decomposed into sulphur. It is a factor for the determination of higher yield seed and quality of seed that contain oil in oilseed crops. Since, sulphur is immobile in plants, it must be continuously supplied from crop emergence till maturity. So, reduced yield can come from sulphur shortage at any stage of growth. It can be overcome by supplying sulphur containing fertilizers to plants by soil application and foliar application. Among the sources of sulphur, elemental sulphur and gypsum are widely utilized in sulphur deficiency.

**Keywords: Oilseed, Sulphur, deficiency, quality, nutrient uptake.**

## Introduction

India is the 4<sup>th</sup> largest producer country of oilseed crops after USA, China, Brazil; contributing 10% to global production and 20% to the global area. Oilseed self-sufficiency achieved during the “Yellow Revolution” in the early 1990s could only last for a limited time (Roy *et al.*, 2022). India is also one of the major oil importing countries in the world. The largest oilseed production states in India include Andhra Pradesh, Haryana, Gujarat, Karnataka, Madhya Pradesh, Maharashtra, Rajasthan, Tamil Nadu, Uttar Pradesh and West Bengal. The major producing states are Rajasthan, Gujarat, Madhya Pradesh and Maharashtra with respective shares of the total production of about 20%, 20%, 19% and 16%. Oilseeds have a lot of fat in addition to having a lot of protein. Therefore, they are not only good sources of protein, also concentrated sources of energy (Sarwar *et al.*, 2013). Oilseeds are a vital component of our nutrition and the oil extracted from them is used in a number of ways such as paints, varnishes, hydrogenated oil, soaps, perfumes, lubricants and other. Growing oilseeds may significantly alter the diet of humans, improving food security and creating income generation.

Sulphur is one of the essential micronutrients that is need for higher yield and quality of oilseed crops. In accordance with its chemical nature, it plays significant role in a number of cellular metabolic process (Hawkesford *et al.*, 2005). With an average concentration of 0.06 percent, it is the 13<sup>th</sup> most prevalent element in the crust of the earth. Sulphur comprises amino acids are cysteine, cystine and methionine, which makes it an essential factor for plant growth (Parmaret *et al.*, 2018). These amino acids contain almost 90% of the sulphur present in plant (Tandon and Messick, 2002). Crop cannot reach their maximum potential in terms of protein content, oil quality and yield without satisfying sulphur requirement (D'Hooghe *et al.* 2013; Poisson *et al.* 2018a, b). Sulphur is required for the production of co-enzyme A, which is involved in the oxidation of fatty acids, their synthesis and the intermediates of citric acid cycle (Raghuvver M, 2017). Sulphur is utilized as a pesticide, a plant nutrient for increasing yield and quality of oilseed crops, an amendment for amelioration and a chemical agent (Kanwar and Mudahar, 1986). Sulphur fertilization in oilseed crops help to increase the productivity of oilseed crops. It also enhances the availability of other nutrients like phosphorus, potassium, zinc and suppress the uptake of sodium and chlorine, which are harmful to plant growth and development.

Oilseed productivity is low, due to inadequate and unbalanced fertilization is the most important one. One of the main causes of imbalanced fertilization is sulphur deficiency. The shortage of S causes a reduction in plants' photosynthetic efficiency, which in turn causes a decline in crop productivity particularly in oilseed crops, which have a high S demand (Aarabi *et al.*, 2021). Sulphur deficiency causes by decreasing the supply of sulphate, which delays the crop maturity and reduces the plant height and quality (Hawkesford, 2000; Dobermann, 2000). And also reduces oil content and quality of oilseeds (Scherer, 2001). Sulphur deficiency in coarse texture soils is due to low availability of organic matter (Takkar, 1988).

### **Sulphur nutrition requirements in oilseed crops**

Several researchers have determined the S need for oilseed crops in the field (Banu *et al.*, 2017; Tandon 1991a, 1991b; Aulakh and Pasricha, 1988; Pasricha *et al.*, 1988 and Kanwar *et al.*, 1983). The Sulphur requirement of oilseeds can be satisfied by using a number a S-containing substances including phosphogypsum, elemental S, gypsum and pyrite. In general, applying 30-40 kg S/ha to groundnut was more beneficial (Singh *et al.*, 1991b; Kale, 1993; Patra *et al.*, 1995). Gypsum added in the pegging zone at flowering at a rate of 250 kg/ha

improved pod yield by 20.5% in the demonstration plots (Singh *et al.*, 1991). In general, the ratio (P:S) of the S needs of crops in comparison to P is 1.3 for cereals, 0.8 for legume forages and 0.6 for crucifers. It appears that S uptake in oilseeds is almost two times as high as phosphorus uptake (Tiwari *et al.*, 2006). Among various crops, oilseeds have higher sulphur requirements than cereals and pulses due to the crucial function that sulphur plays in the synthesis of oil and the production of bold grains, which are necessary for the production of oil (Khalid *et al.*, 2011).

### **Role of sulphur in oilseeds**

Sulphur is mostly absorbed by plant roots as sulphate ( $\text{SO}_4^{2-}$ ), but it can also be taken up as thiosulphate ( $\text{S}_2\text{O}_3^{2-}$ ). In addition, leaves can absorb a little amount of  $\text{SO}_2$  (Havlin *et al.*, 2005). The amount of sulphur present in oilseed crops is 1.1-1.7 %. Reduced amount of sulphur applied to oilseed crops, which results in the production of low-quality seed (Haneklaus *et al.*, 2016). In rapeseed-mustard, sulphur is necessary to enhance the quantity of glucosides, glucosinolate and protein (Chandeleet *et al.*, 2003). Rapeseed-mustard is an important oilseed crop, has a high demand for sulphur. Due to its high sulphur requirements, application of 60 kg/ha significantly affected the amount of glucosinolate present and oil percentage increased by 44.6%.

When sulphur fertilizer levels increased from 0 to 60 kg S kg<sup>-1</sup>, the content of allyl isothiocyanate in mustard seeds increased (Chandeleet *et al.*, 2003). Sulphur has a crucial function in the production of protein, vital enzymes and redox processes, which makes it essential for the formation of chloroplasts and chlorophyll. Sulphur is also associated with nodulation, flowering and the quality of oilseeds, particularly groundnut (Rao *et al.*, 2013).

### **Sulphur and quality of crop**

Sulphur is involved in oil synthesis and oil storage organs which are rich in sulphur. Despite the composition of plant protein and oil, variations in the supply of S for plant growth change the quantity of various kind of chemicals and the composition overall. Sulphur improves the forage's digestibility for ruminants (Rendig, 1986). Zhao *et al.*, (1997) and enhanced the glucosinolate content of rapeseed. S deficient rapeseed crop seed contains lesser erucic acid and linolenic and higher oleic acid (Beaton and Fox, 1971). The increase in protein content as a result of S application have been reported to be 14% in soybeans (Aulakh *et al.*, 1990), 6%

in lentils (Tiwari, 1989), 19% in sunflower (Gangadhara *et al.*, 1990), 28-34% in rapeseed and mustard (Pasricha *et al.*, 1987). S is a component of oil compounds and raises the oil content of the majority of crops. With S fertilization, an average 5% increase in oil content has been observed (Pasricha *et al.*, 1987). However according to research, S application increased oil percentage on average by 11.3% in groundnut, 9.6% in rapeseed-mustard, 6% in linseed and 2-3.8% in safflower, sunflower and sesame (Tandon 1991a). Singh and Singh (1983) in a study examined different sources of S for seed and oil yields, sulphur and oil contents and S uptake by mustard and revealed that all of these increased with each source of S, but element S was the best. Sunflower seed weight and head diameter increased after S application (Gangwar and Parmeswaran, 1996a). In rapeseed-mustard, the S application increased oil yield by 16% and oil content by 3.4% (Sachdeva and Dev, 1990).

### **Source of sulphur for various oilseed crops**

Ammonium sulphate (24% S), single super phosphate (12% S), gypsum (13% S), pyrite (22-30% S) and element S (85- 100% S) are the most widely used S fertilizers. The other sources of S are potassium sulphate (18% S), magnesium sulphate (13% S), phosphogypsum (16% S), sulphur sludge (10-16% S), press mud (2-3% S). For most of the oilseed crops, the S fertilizers containing  $\text{SO}_4^{2-}$  were effective (Tandon, 1991a). Applying pyrites and elemental sulphur 20-25 days before planting is recommended since elemental sulphur requires oxidation, pyrites is a slow-release fertilizer (Tiwari *et al.*, 1984). Gypsum and phosphogypsum are suitable for both alkaline and acidic soils, however they are more suited for crops that need a lot of Calcium, such as groundnut (Rao *et al.*, 2013). The ammonium sulphate corrects both the N and S deficiency and is the best suited for oilseed crops that do not fix nitrogen. The single super phosphate is a good source of S and P necessary for integrated P and S application in most oilseeds.




Fig 1: Sulphur Cycle

Source: RaghuveerMunigela, Effect of sulphur on oilseed crops. Nov. 29, 2018.

### Deficiency symptoms of sulphur in oilseed crops

Since only nitrogen, phosphorus and potassium managed nutrients in India, there are significant sulphur deficiency problems (Tandon, 1991). There is a wide spread sulphur deficiency in the world's soil (Scherer, 2009). The main symptom of sulphur nutrient deficiency is yellowing of younger leaves, which may be caused by inadequate chlorophyll production. Sulphur is immobile in plant so its deficiency appears on young leaves. Deficiency of sulphur mostly observed in the shoots rather than the roots, which causes plants become stunted and shoot become slender (Karthika *et al.*, 2018). Sulphur deficiency leads to reduced root hydraulic conductivity, which is likely associated with indicating nutrient starvation from root to shoot. Additionally, sulphur deficiency causes an imbalance between the ratio of nitrogen and sulphur pool and increases the soluble nitrogen with amide and nitrate. When there is a sulphur deficiency in groundnuts, trifoliolate leaves grow and V-shape because they are upright from the petiole (Sing *et al.*, 1999). Young leaves of groundnut frequently develop chlorosis due to sulphur deficiency (Supakammerdet *et al.*, 1993; Singh and Chaudhari, 1995). In rapeseeds, the young leaves become cupped shape with various shape ranging from pale green to complete yellow, including the vein, which can occasionally turn into reddish or purple. In soyabeans, the newly emerging leaves still have a pale yellow-green, indicating a deficit of sulphur. In case of severe deficiency, the entire plant becomes pale yellow and the size of leaves and internodes are reduced. In sunflower, shorter internodes cause a reduction in plant height and the leaves and inflorescence both become pale.



Fig 2: Effect of sulphur in oilseed crops

Source: [agritech.tnau.ac.in/agriculture/plant\\_nutri/sul.html](http://agritech.tnau.ac.in/agriculture/plant_nutri/sul.html)

### **Assessment of sulphur in soil and plant sample**

#### **a) Assessment of sulphur in soil analysis:**

Different techniques can be used to determine the amount of sulphur in soil, however in India the most common way is to use a 0.15%  $\text{CaCl}_2$  solution (Kour and Jalali, 2008). This approach is mostly used for sulphur deficient soils or soils with less than 10ppm of sulphate. The method should be reliable and appropriate for the crop (Jackson, 2011).

#### **b) Assessment of sulphur in plant analysis:**

The level of sulphur in the soil can be determined using basic analytical techniques. There are two ways to assess sulphur in plants: wet digestion and dry digestion. In wet digestion,  $\text{HNO}_3$  and  $\text{HClO}_4$  can be used, while dry ashing can be used to extract sulphur from plant digest (Mahanta *et al.*, 2017).

### **Application of sulphur fertilizers**

The main ways for applying sulphur fertilizers in soils are broadcast or dribble banded, broadcast and subsequent incorporation, band placement, seed placement and banded near the seed. Broadcasting and band placement techniques are two of them that are frequently used. Sulphur use efficiency can be increased by managing sulphur fertilization in soil using the 4R nutrient stewardship approach (Right Source of nutrient applied at the Right Rate, Right Time and Right Place). To prevent the loss of nutritional element S, scientific management techniques can be used. For both a short-term and long-term source of plant nutrition, a mixture of soluble  $\text{SO}_4^{2-}$  and elemental S may be helpful.

### **Effect of Sulphur on nutrient uptake of oilseed**

In an experiment on groundnuts, Venkateshet *al.*, (2002) found that gypsum application at 45kg/ha of sulphur increased S uptake by straw (4.62 kg/ha) and kernel (10.37 kg/ha) over control (4.39 kg/ha, 1.91 kg/ha). Ammonium sulphate and the application of 20 kg S/ha during two consecutive years resulted in sulphur use efficiency of 9.7,10.1 kg/S kg<sup>-1</sup> and 15.9, 3.8 kg/S kg<sup>-1</sup>(Pandey and Ali,2012). The uptake of N (131.0 kg/ha), P(31.1 kg/ha), K(87.92 kg/ha) and S (27.47 kg/ha) by mustard was significantly impacted due to the application of 90 kg Sha<sup>-1</sup> over control (83.86, 19.8, 53.91,11.20 kg/ha) as noted by Pachauri and Trivedi, 2012. Application of S enhanced the concentration and uptake of N, P, S, Fe and Zn but reported effect on the concentration of K, Mg, Mn or Cu in ground nut leaves (Singh and Chaudhari, 1995). Sulphur uptake (20.7 kg/ha) in Indian mustard has been reported highest because of 45 kg/ha Sulphur application.40 kg S/ha application, mustard's uptake of total N (39.34 kg/ha), total P (10.67 kg/ha) and total S (17.73 kg/ha) was considerably impacted (Singh and Thenua, 2016).

## **Conclusion**

In addition to promoting yield and quality, sulphur enhances the oil content in oilseed crops, contributing to better nutritional value. It also plays an important role in growth and development of plant. Deficiency of sulphur caused growth retarded and yield. Gypsum, pyrites, element S and phosphogypsum are the sources that can provide sulphur. It can also be applied using fertilizers that contain primary nutrient like ammonium sulphate, single super phosphate and sulphate of potash. Oilseeds need to produce a good yield along with good quality and this can only be sustained when the plant receives enough sulphur. To achieve the best production results in oilseed crops, sulfur application must be balanced according to crop requirements and soil condition.

## **Reference**

- Aarabi F, Rakpenthai A, Barahimipour R, Gorka M, Alseekh S, Zhang Y, Hoefgen R. Sulphur deficiency Induced genes affect seed protein accumulation and composition under sulphate deprivation. *Plant Physiol.* 2021.
- Aulakh MS and Pasricha NS. Sulphur fertilizer of oilseeds for yield and quality. *Proc. TSI-FAI Symposium on Sulphur in Indian Agriculture, New Delhi, India.1988;SII/3: 1-14.*

- Aulakh MS et al. P-S interrelationships for soybean on P and S deficient soil. *Soil Science* 1990;150, 705-709.
- Banu R, Shroff JC, Shah SN. Effect of sources and levels of sulphur and biofertilizer on growth, yield and quality of summer groundnut. *International Journal of Agronomy Science*. 2017;13(1): 67-70.
- Beaton JD, RL Fox. Production, marketing and use of sulphur products. 1971; pp.335-379. In R.A. Olsen et al. (eds). *Fertilizer technology and use*, 2nd ed. Soil Science Society. America, Madison.
- Chandel RS, Sudhakar PC, Singh K. Response of sulphur nutrition in mustard—A review. *Agricultural Reviews*. 2003; 24(3),175-182.
- D'Hooghe P, Escamez S, Trouverie J, Avice JC. Sulphur limitation provokes physiological and leaf proteome changes in oilseed rape that lead to perturbation of sulphur, carbon and oxidative metabolisms. *BMC plant boil*. 2013; 13:1-5.1471-2229-13-23.
- Dobermann A. Rice: Nutrient disorder and nutrient management. International Rice research Institute 2000.
- Dr. Kumar V, Tiwari A. Sparking Yellow Revolution in India Again. Department of Economic Analysis and Research. 2020;1pp.
- Gangasaran, GGiri. Influence of N, P and S on mustard under semiarid rainfed conditions of north west India. *Indian Journal of Agronomy*. 1990; 35:131-136.
- Gangwar, MS, PM Parmeswaran. Phosphorus-sulphur relationship in sunflower. I Effect of P and S on yield. *Oilseed Journal*. 1976;6 (3): 28-32.
- Haneklaus S, Bloem E, Schnug E, de Kok LJ, Stulen I. Sulfur. *Plant Nutrition*. 2016; 7,183.
- Hawkesford MJ. Plant responses to sulphur deficiency and the genetic manipulation of sulphate transporters to improve S-utilization efficiency. *Journal of Experimental Botany*. 2000 Jan; 51(342):131-8. PMID: 10938804.
- Hawkesford, M.J. Sulfur. In *Nutritional Genomics*. Broadley MR., White P, Eds.; Blackwell Publishers: Oxford, UK, 2005; pp 87-11.
- Havin JL, Beaton JD, Tisdale SL, Nelson WL. *Soil fertility and Fertilizers: An introduction to nutrient management*. 7, ed. New Jersey, Pearson Prentice Hall, 2005; 528p.

- Jackson D. Soils, nutrients and water. Temperate and Subtropical Fruit Production. 2011; p.101.
- Kale MR. Effect of sulfur on growth and yield of groundnut. Indian Journal Agronomy 1993;38(3):504-505.
- Kanwar JS, Mudahar MS. Fertilizer Sulphur and Food Production. Martinus Nijhoff/ Dr W. Junk Publisher, Dordrecht. 1986;247p.
- Kanwar JS, Nijhawan HL, Raheja SK. Groundnut Nutrition and Fertilizer Responses in India. ICAR New Delhi. 1983;185p.
- Karthika KS, Rashmi I, Parvathi MS. Biological functions, uptake and transport of essential nutrients in relation to plant growth. In Plant nutrients and abiotic stress tolerance Springer, Singapore. 2018; pp. 1-49.
- Khalid R, Khan KS, Akram Z, Qureshi R, Gulfraz M. Relationship of plant available sulphur with soil characteristics, rainfall and yield levels of oilseed crops in Pothwar Pakistan. Journal of Botany. 2011;43(6):2929-2935.
- Kour S, Jalali VK. Forms of sulphur and their relationships in soils of different agro-climatic zones of Jammu region. Journal of the Indian Society of Soil Science. 2008; 56(3),309-312.
- Mahanta PL, Singh AK, Radhamani R, Rao DP. Determination of Total Sulfur and Sulfate Sulfur D in Geological Materials of Rocks, Soils, and Sediments by ICP-OES After Sample Digestion Using Alkali Flux. *Atomic Spectroscopy*. 2017;38, 99.
- Manoj Pandey, Javed Ali. Effect of sources and levels of sulphur on sulphur uptake, yield and quality of linseed (*Linum usitatissimum*). Annals of Agricultural Research New Series. 2012;33(1-2):32-35.
- Naphade PS, Wankhede SG. Response of groundnut to different levels and source of S. PKV Research Journal 1988;12: 71-73.
- Parmar NN, Patel AP, Chaudhary M. Effects of Sources and Levels of Sulphur on Growth, Yield and Quality of Summer Sesame South Gujarat Condition (*Sesamum indicum* L.). IJCMAS 2018; 7(2): 2600-2605.

- Pasricha NS, Aulakh MS, Bahl GS, Baddesh HS. Nutritional requirement of oilseed and pulse crops in Punjab. *Department of Soils*. PAU, Ludhiana. 1987.
- Pasricha NS, Aulakh MS, Sahota NS, Baddesh HS. Fertilizer use research in oilseed crops. *Fertilizer News*. 1988;33(9): 15-22.
- Patra AK, Tripathy SK, Samui RC. (1995). Response of groundnut (*Arachis hypogea* L.) to sulphur. *Indian Journal of Agricultural Science*. 1995;65 (80): 597-598.
- Poisson E, Brunel-Muguet S, Kauffmann F, Trouverie J, Avice JC, Mollier A. Sensitivity analyses for improving sulphur management strategies in winter oilseed rape. *PLoS ONE* 2018a;13: e0204376.
- Poisson E, Mollier A, Trouverie J, Avice JC, Brunel-Muguet S. SuMoToRI model simulations for optimizing sulphur fertilization in oilseed rape in the context of increased spring temperatures. *European Journal of Agronomy*. 2018b; 97:28-37.
- Raghuveer M. Effect of Sulphur on Oilseed Crops. *Biotect Articles*. 2017.
- Rending VV. Sulphur and crop quality. In *Sulphur in Agriculture* (M. A. Tabatabaie). Agronomy monograph no. 27, American Society of Agronomy, Madison, Wisconsin, USA. 1986; pp. 635-653.
- Raman Kumar Pachauri, Trivedi SK. Effect of sulphur levels on growth, yield and quality of Indian mustard (*Brassica juncea*) genotypes. *Annals of Agricultural Research New Series*. 2012;33(3):131-135.
- Rao KT, Rao AU, Sekhar D. Effect of sources and levels of sulphur on groundnut. *Journal of Academia and Industrial Research*. 2013; 2(5), 268-270.
- Roy A, Singh KM, Kumari T, Ahmad N, Singh RP, Sinha DK. Dynamics of Oilseeds Production and Decomposition of output Components of Oilseeds in Bihar. *Economic Affairs*. 2022;67(02): 23-29.
- Sachdev MS, Dev DL. Nitrogen and S uptake and efficiency in mustard-moong-maize cropping system. *Fertilizer News*. 1990;35(7): 49-55.
- Sarwar MF, Sarwar MH, Sarwar M, Qadri NA, Moghal S. The role of oilseeds nutrition in human health. *Journal of Cereals and Oilseeds*. 2013; 97pp.
- Scherer HW. Sulphur in crop production. *European Journal of Agronomy* 2001; 81-111.

- Scherer HW. Sulphur in soils. *Journal of Plant Nutrient and Soil Science*. 2009;172: 326-335.
- Singh AL, Chaudhari V. Source and mode of sulphur application on groundnut productivity. *Journal of Plant Nutrition*. 1995;18: 2739-2759.
- Singh AL. Sulphur Nutrition of Oilseed Crops. *Advances in Plant Physiology* (Ed. A. Hemantranjan). 1999;2,201-226.
- Singh BP, Singh HG. Comparative efficacy of S on production of green matter, mineral composition and uptake of mustard on Vertisols. *Forage Research*. 1983; 9:37-41.
- Singh MV. Proc. 18th workshop meeting and results of practical utility. All India Coordinated Scheme of Micro and secondary Nutrients and Pollutants Elements in soils and Crops. Indian Institute of Soil Science. Bhopal. 1991.
- Singh SB, Thenua OVS. Effects of phosphorus and sulphur fertilization on yield and NPS uptake by mustard (*Brassica juncea* L.). *Progressive Research-An International Journal* 2016;11(1):80-83.
- Supakamnerd N, Dell B, Bell RW. Diagnosis of sulphur deficiency in peanut (*Arachis hypogaea*) by plant analysis. In: M.L. Van Beusichem (Ed) *Plant Nutrition-Physiology and application*. XI<sup>th</sup> IPNC 30 July 4 August 1989, Wageningen, Netherland Kluwer Academic Publishers, Dordrecht. 1990; pp.791-795.
- Takkar PN. Sulphur status of Indian soils. Proc. TSI-FAI Symp. Sulphur in Indian Agriculture, New Delhi 1988. S/I/2/1-31.
- Tandon HLS. Sulphur Research and Agriculture Production in India 3<sup>rd</sup> edition FDCO, New Delhi, India. 1991a.
- Tandon HLS. Secondary and micronutrients in Agriculture-Guide Book-cum-Directory. 2<sup>nd</sup> Ed. FDCO, New Delhi, India. 1991b;122 pp.
- Tandon HLS, Messick DL. Practical Sulphur Guide. The Sulphur Institute, Washington, D.C. 2002; pp. 20.
- Tiwari KN, Dwivedi BS, Pathak AN. Evaluation of iron pyrites as sulphur fertilizer. *Fertilizer Research*. 1984; 5:235-243.

Tiwari KN. Pyrites. In: H.L.S. Tandon (ed.) Sulphur Fertilizers for Indian Agriculture-A guide book. Fertilizer Development and consultation organization, New Delhi, India. 1989;pp. 72-83.

Tiwari KN, GuptaBR. Sulphur for Sustainable High Yield Agriculture in Uttar Pradesh. Indian Journal of Fertility. 2006;1, 37-52.

TNAU, 2015. Mineral Nutrition: Oilseed: Sunflower. [agritech.tnau.ac.in/agriculture/plant\\_nutri/sul.html](http://agritech.tnau.ac.in/agriculture/plant_nutri/sul.html)

Venkatesh MS, Majumdar B, Lal B, Kailash Kumar. Relative performance of sulphur sources on sulphur nutrition of groundnut (*Arachis hypogea*) in acid Alfisol of Meghalaya. Indian Journal of Agricultural Sciences. 2002;72(4):216-9.

Zhao FJ, Withers PJA, Evans EJ, Monaghan J, Salmon SE, Shewry PR, Mcgrath SP. Sulphur nutrition: An important factor for the quality of wheat and rapeseed. In Plant Nutrition- For Sustainable Food Production and Environment (T. Ando et al Eds). Kluwer Academic Publishers, Printed in Japan. 1997;pp 917-922.