

# Farmer perceptions on the effects of termites in Kwa Vonza Location, Kitui County, Kenya

## ABSTRACT

**Aims:** The aim of this study was to assess: (i) the perception of Kwa Vonza farmers regarding the presence of termites in their land (ii) the perceived importance of termites (iii) how they control termites in their properties.

**Study design:** This study is based on responses to a questionnaire sent to farmers.

**Place and Duration of Study:** The study took place in Kwa Vonza Location, Yatta Sub County, Kitui County, Kenya between April and November 2017.

**Methodology:** 60 questionnaires each comprising 20 multiple choice questions were sent out to farmers. Out of these, 54 responses were received. Where the response did not require a Yes or No answer, it elicited a response from a standardized five-point scale to demonstrate agreement to the provided proposition. Data analysis was done using descriptive statistics in which percentages were tabulated and frequency tables generated using Microsoft Excel.

**Results:** 73% of farmers perceived their land as infertile with the causes of infertility ranging from soil erosion, drought, and mono-cropping. 87% of farmers acknowledged termite infestation in their farms but perceived them as destructive. Termite infestation was attributed to deforestation, drought and flooding. Chemical control was the method of choice to manage termites.

**Conclusion:** Kwa Vonza farmers do not perceive termites to be of any agronomic significance but view them as destructive agents. Further research to address the complex issue of soil management at the farm-scale level that involves farmers is necessary to fill gaps in scientific knowledge and producing advice for practical use.

*Keywords: termites; soil fertility; tropical ecosystems; soil biodiversity; farmer perceptions*

## 1. INTRODUCTION

Termites (*Isoptera*) are an order of insects consisting of 2,500 species out of which 300 are considered pests (Ndiaye et al., 2004). Depending on their family or sub-family, they build their nests underground, in wood or termite mounds (Heyde et al., 2021). Because termites display a high sensitivity to the biotic and abiotic environmental conditions they are exposed to, they can play a key role in tropical ecosystems (Jouquet et al., 2016) just like earthworms do in temperate ecosystems (Manono and Moller, 2015). These soil organisms play important roles in linking abiotic and biotic components of soil ecosystem by supplying 'soil services' such as nutrient cycling, decomposition, and plant growth (Apori et al., 2020; Jouquet et al., 2016; Manono 2016a).

24 Large amounts of essential nutrients in the soil are bound in organic form (Khasabulli et al.,  
25 2023; Manono, 2014). For these nutrients to be released and made available for plant  
26 absorption, they must undergo decomposition and mineralization (Marzi et al., 2021;  
27 Manono et al., 2019). Through their feeding and barrowing activities, termites and  
28 earthworms directly influence the breaking down, mixing, and transportation of organic  
29 matter and mineral nutrients. They create macrospores that enhance infiltration, water  
30 storage, and air regulation while providing channels for root growth and penetration (Manono  
31 2019; Jouquet et al., 2016). On the negative side, they both contribute to the emissions of  
32 greenhouse gases (Quevedo et al., 2021; Manono 2016b).

33 Despite playing these important roles, termites are one of the most damaging pests in the  
34 tropics and can cause considerable problems in agriculture (Kagezi et al., 2023;  
35 Govorushko, 2019). They feed on dead organic matter, but when not available, they will eat  
36 live plant material including crops. In arid and semi-arid lands, termites build underground  
37 nests and collect live green plant material such as living grass, crops, seedlings, and weak  
38 wilting plants (Lopez-Hernandez, 2023; Lepage et al., 1993). Because of the beneficial roles  
39 these organisms play in agroecosystems, they should be carefully managed considering  
40 their benefits against the rate of loss to the ecosystem (Manono, 2016c; Nyeko and Olubayo,  
41 2005).

42 Individual farmers are the stewards and decision makers about what happens on their land  
43 (Kalovoto et al., 2020; Manono, 2016c; Sileshi, et al., 2008). It therefore follows that  
44 sustaining agricultural productivity depends on maintaining and enhancing the abundance  
45 and functional activities of these soil organisms (Manono, 2016c; Sileshi, et al., 2008).  
46 Farmers could benefit from more quantitative evidence of their perception and knowledge on  
47 these organisms and their activities in soil fertility and crop production (Manono, 2016c). For  
48 example, improved farmer understanding of the importance of these organisms could enable  
49 development of suitable decision support tools that emphasize their management in  
50 contributing to agricultural sustainability. This is particularly important in systems that are  
51 susceptible to changing land use and management pressures associated with persistent  
52 drought like Kwa-Vonza location, Kitui County, Kenya. It is with this background that this  
53 study explores how small-scale farmers in Kwa-Vonza perceive termites in relation to soil  
54 fertility and crop production. Thus, the aim of this study was to assess: (i) the perception of  
55 Kwa vonza farmers regarding the presence of termites in their land (ii) the perceived  
56 importance of termites (iii) how they control termites in their properties.

## 57 **2. MATERIAL AND METHODS**

### 58 **2.1 Study area**

59  
60  
61  
62 This study is based on responses to a questionnaire sent to farmers in Kwa Vonza Location,  
63 Yatta Sub County, Kitui County, Kenya, between April and November 2017. The coordinates  
64 of the study area are 44°38' to 44°54' S and 170°59' to 171°08' E. Kwa Vonza is located at  
65 130 Kilometers South East of Nairobi on the Machakos – Kitui road. It is part of the Yatta  
66 Plateau, which stretches from the north to the south of the county and lies between Rivers  
67 Athi and Tiva. The area experiences a semi-arid climate with very erratic and unreliable  
68 rainfall. Annual temperatures range between a minimum of 14 to 22° centigrade and a  
69 maximum of 26 to 34° centigrade. There are two rainfall seasons: long rains between March  
70 and May and short rains between October and December. The area's soils are low in fertility  
71 and range from sedimentary rocks, red sandy soils, to clay black cotton soils.

### 72 **2.2 Farmer recruitment and analysis.**


73  
74

75 Respondents were farmers in Kwa Vonza Location with over 3 years of farming experience.  
 76 The questionnaire comprised 20 multiple choice questions designed after trials to take no  
 77 longer than 25 min to complete. Where the response did not require a Yes or No answer, it  
 78 elicited a response from a standardized five-point scale to demonstrate agreement to the  
 79 provided proposition. A blank space was left after each question for respondents to give an  
 80 open-ended response or clarification. Participating farmers were given informed written  
 81 consents, had their anonymity guaranteed, were reminded that they did not have to  
 82 participate and that they could stop participation at any stage or refuse to answer certain  
 83 questions. Sixty questionnaires were sent, out of which 54 responses were received. Some  
 84 respondents did not answer every question. Therefore, the percentages reported in the  
 85 analysis are for individual questions. Data analysis was done using descriptive statistics in  
 86 which percentages were tabulated and frequency tables generated using Microsoft Excel.  
 87


### 88 3. RESULTS AND DISCUSSION

#### 89 3.1 Farmer's perception of their land's fertility

90 Farmers perceive their land's fertility differently with a majority saying theirs is neither fertile  
 91 nor infertile followed by those saying theirs are infertile (Figure 1). The two categories  
 92 combined comprised of 73% of the respondents. When asked what they perceive to be the  
 93 cause of the infertility, the farmers gave a mix of results, ranging from soil erosion, drought,  
 94 and mono-cropping (Figure 2). This observation was consisted with (Okoba and De Graaff,  
 95 2005). Although the majority of farmers perceived termites to be destructive (Figure 4), only  
 96 14% of the farmers attributed soil infertility to termites (Figure 2). This was so, even when a  
 97 greater percentage (87.5%) of the farmers acknowledged of termite infestation in their lands,  
 98 with a majority of these farmers comprising of 65% not attribute termites to any soil benefits  
 99 (Figure 3).  
 100  
 101




102  
 103  
 104 **Fig.1. Farmers perception on soil fertility.**  
 105  
 106  
 107  
 108



109  
110  
111

**Fig. 2. The reasons for soil infertility.**




112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127

**Fig. 3. Termite infestation in the farms and perception of benefit.**

**3.2 Effect of termite infestation in farms.**

Majority of farmers comprising 40% associated termites with crop destruction while 33% associated their presence with lower yields (Figure 4). A further 16% associated termites with house destruction. Only 5% of the farmers’ perceived termites has playing a role in enhancing soil fertility (Figure 4). From this perspective, it would be noted that a majority of farmers comprising 89% of respondents perceived termites as destructive in consistent with other studies (Govorushko, 2019; Lepage et al., 1993). It should be noted that the arguments put forward by scientists on the beneficial roles of termite as a result of their bioturbation activities that lead to breaking up of surface crusts, reducing soil compaction, increasing soil porosity, improving water infiltration and enhancing water holding capacity, and litter degradation (Jouquet et al., 2016) may be invisible to farmers and therefore, were not able to be recognize these roles and associate them with termites.




128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140

**Fig. 4. Effects of termite infestation in the farms.**

### 3.3 Causes of termite infestation

A majority comprising of 72% of respondents attributed termite infestation to three factors, viz, deforestation, drought, and flooding (Figure 5). Out of these, deforestation and drought accounted for 52%. Only 2% of respondents considered termites to be food, a contradiction to the majority who consider termites as food in western Kenya (Kinyuru et al., 2013). Kwa-Vonza being a drought prone area, when it rains, it causes flooding in the termite mounds. This forces them out of their mounds to avoid drowning just like earthworms do (Manono, 2014). In such circumstances, people are able easily see them. However, during droughts and deforestation termites tend to be all over the place (Charles et al., 2021).




141  
142  
143  
144  
145  
146  
147  
148

**Fig. 5. Causes of termite infestation.**

### 3.4 Strategies of controlling termites

Because farmers perceived termites to be destructive, it was prudent to enquire how they controlled them. Majority of the respondents, comprising 45% controlled termites by chemicals while 26% physical destroyed termite mounds to remove the queen (Figure 6). Chemical control is used as a termite control strategy in many places (Ahmad et al., 2021;

149 Ejomah et al., 2020). Another 22% used wood ash from their kitchen. This method has been  
150 reported to be used in other studies (Oliver-Villanueva et al., 2013).  
151  
152



153  
154 **Fig. 6. Methods used to control termites.**

### 155 **3.4 Methodological constraints**

157 Caution should be taken when interpreting the outcome of this study because of the small  
158 sample size and homogeneous properties of the farmer respondents. Nevertheless, 90% of  
159 farmers contacted responded to questionnaires and this should be considered as a model  
160 study that should be expanded to other regions.

## 161 **4. CONCLUSION**

162  
163 This study revealed that farmers in Kwa Vonza do not perceive termites to be of any  
164 agronomic significance but view them as destructive agents. This calls for further social  
165 science research to address the complex issue of soil management at the farm-scale level.  
166 Farmer involvement may help in prioritizing options for filling gaps in scientific knowledge  
167 and producing advice for practical use.

## 168 **ACKNOWLEDGEMENTS**

169  
170 We thank all the individual farmers who participated in this study.

## 171 **COMPETING INTERESTS**

172  
173 "Authors have declared that no competing interests exist."

## 174 **AUTHORS' CONTRIBUTIONS**

175  
176 Authors contributed equally in the design, data collection, analysis and writing of this article.  
177 All authors read and approved the final manuscript.

183  
184  
185  
186  
187  
188  
189  
190  
191  
  
192  
193  
194  
  
195  
196  
197  
198  
199  
200  
  
201  
202  
203  
  
204  
205  
206  
  
207  
208  
209  
  
210  
211  
212  
213  
  
214  
215  
216  
217  
  
218  
219  
220  
  
221  
222  
223

## REFERENCES

1. Ahmad, F., Fouad, H., Liang, S. Y., Hu, Y., & Mo, J. C. (2021). Termites and Chinese agricultural system: applications and advances in integrated termite management and chemical control. *Insect science*, 28(1), 2-20.
2. Apori, S. O., Murongo, M. F., Hanyabui, E., Muli, G. K., & Wamuyu, B. (2020). Role of military termites (*Pseudocanthotermes militaris*) in improving soil productivity in tropical agroecosystems.
3. Charles, G. K., Riginos, C., Veblen, K. E., Kimuyu, D. M., & Young, T. P. (2021). Termite mound cover and abundance respond to herbivore-mediated biotic changes in a Kenyan savanna. *Ecology and Evolution*, 11(12), 7226-7238.
4. Ejomah, A. J., Uyi, O. O., & Ekaye, S. O. (2020). Exposure of the African mound building termite, *Macrotermes bellicosus* workers to commercially formulated 2, 4-D and atrazine caused high mortality and impaired locomotor response. *Plos one*, 15(3), e0230664.
5. Govorushko, S. (2019). Economic and ecological importance of termites: A global review. *Entomological Science*, 22(1), 21-35.
6. Heyde, A., Guo, L., Jost, C., Theraulaz, G., & Mahadevan, L. (2021). Self-organized biotectonics of termite nests. *Proceedings of the National Academy of Sciences*, 118(5), e2006985118.
7. Jouquet, P., Bottinelli, N., Shanbhag, R. R., Bourguignon, T., Traoré, S., & Abbasi, S. A. (2016). Termites: the neglected soil engineers of tropical soils. *Soil Science*, 181(3/4), 157-165.
8. Kagezi, G. H., Twesigye, V., Musasizi, J. K., Ssebugenyi, I., Namara, E., Ssenoga, G., ... & Arinaitwe, G. (2023). Farmers' Knowledge and Perception of the Pest Status and Management Options for Termites in Buikwe District, Central Uganda.
9. KalovotoDamariis, M., Kimiti Jacinta, M., & Manono Bonface, O. (2020). Influence of women empowerment on adoption of agroforestry technologies to counter climate change and variability in semi-arid Makueni County, Kenya. *International Journal of Environmental Sciences & Natural Resources*, 24(2), 47-55.
10. Khasabulli, B. D., Mutisya, M. D., Anyango, S. P., Manono, B. O., & Odhiambo, D. G. (2023). Soil Microbial Biomass, Microbial Population and Diversity in Maize-Banana Based Agroforestry System in Kisii County, Kenya. *Asian J. Res. Crop Sci*, 8(4), 230-239.
11. Kinyuru, J. N., Konyole, S. O., Roos, N., Onyango, C. A., Owino, V. O., Owuor, B. O., ... & Kenji, G. M. (2013). Nutrient composition of four species of winged termites consumed in western Kenya. *Journal of food composition and analysis*, 30(2), 120-124.
12. Lepage, M., Abbadie, L., & Mariotti, A. (1993). Food habits of sympatric termite species (Isoptera, Macrotermitinae) as determined by stable carbon isotope analysis in a Guinean savanna (Lamto, Côte d'Ivoire). *Journal of Tropical Ecology*, 9(3), 303-311.

- 224 13. Lopez-Hernandez, D. (2023). Termite mound as nutrient hot-spots in savannah with  
 225 emphasis in P cycling and the potential use of mounds as soil amendment.  
 226 *Pedobiologia*, 150888.
- 227 14. Manono, B. O., Moller, H., Bengé, J., Carey, P., Lucock, D., & Manhire, J. (2019).  
 228 Assessment of soil properties and earthworms in organic and conventional farming  
 229 systems after seven years of dairy farm conversions in New Zealand. *Agroecology and*  
 230 *Sustainable Food Systems*, 43(6) 678-704.
- 231 15. Manono, B. O. (2016c). New Zealand dairy farm effluent, irrigation, and soil biota  
 232 management for sustainability: Farmer priorities and monitoring. *Cogent Food &*  
 233 *Agriculture*, 2(1), 1221636.
- 234 16. Manono, B. O. (2016b). Carbon dioxide, nitrous oxide and methane emissions from the  
 235 Waimate District (New Zealand) pasture soils as influenced by irrigation, effluent  
 236 dispersal and earthworms. *Cogent Environmental Science*, 2(1), 1256564.
- 237 17. Manono, B. (2016a). Agro-ecological role of earthworms (Oligochaetes) in sustainable  
 238 agriculture and nutrient use efficiency: a review. *Journal of Agriculture and Ecology*  
 239 *Research International*, 8(1), 1-18.
- 240 18. Manono, B. O., & Moller, H. (2015). Effects of stock type, irrigation and effluent dispersal  
 241 on earthworm species composition, densities and biomasses in New Zealand pastures.  
 242 *Pedobiologia*, 58(5-6), 187-193.
- 243 19. Manono, B. O. (2014). Effects of irrigation, effluent dispersal and organic farming on  
 244 earthworms and soil microbes in New Zealand dairy farms (Doctoral dissertation,  
 245 University of Otago).
- 246 20. Marzi, M., Shahbazi, K., Kharazi, N., & Rezaei, M. (2020). The influence of organic  
 247 amendment source on carbon and nitrogen mineralization in different soils. *Journal of*  
 248 *Soil Science and Plant Nutrition*, 20, 177-191.
- 249 21. Ndiaye, D., Lepage, M., Sall, C. E., & Brauman, A. (2004). Nitrogen transformations  
 250 associated with termite biogenic structures in a dry savanna ecosystem. *Plant and Soil*,  
 251 265, 189-196.
- 252 22. Nyeko, P., & Olubayo, F. M. (2005). Participatory assessment of farmers' experiences of  
 253 termite problems in agroforestry in Tororo District. University of Nairobi.
- 254 23. Okoba, B. O., & De Graaff, J. (2005). Farmers' knowledge and perceptions of soil  
 255 erosion and conservation measures in the Central Highlands, Kenya. *Land Degradation*  
 256 *& Development*, 16(5), 475-487.
- 257 24. Oliver-Villanueva, J. V., Gascón-Garrido, P., & Ibiza-Palacios, M. D. S. (2013).  
 258 Evaluation of thermally-treated wood of beech (*Fagus sylvatica* L.) and ash (*Fraxinus*  
 259 *excelsior* L.) against Mediterranean termites (*Reticulitermes* spp.). *Eur. J. Wood Wood*  
 260 *Prod*, 71, 391-393.
- 261 25. Quevedo, H. D., Brandani, C. B., Bento, C. B., Pitombo, L. M., Ferreira Filho, P. J., & do  
 262 Carmo, J. B. (2021). Greenhouse gas emissions from termite mounds in a transition  
 263 area between the Cerrado Savanna and the Atlantic Forest in Brazil. *Acta Oecologica*,  
 264 110, 103690.

265 26. Sileshi, G., Akinnifesi, F. K., Ajayi, O. C., Chakeredza, S., Mng'omba, S., & Nyoka, B. I.  
266 (2008). Towards sustainable management of soil biodiversity in agricultural landscapes  
267 in Africa. Biodiversity, 9(1-2), 64-67.

268