

Effect of gibberellic acid and zinc sulphate on growth, yield and quality of Cucumber (*Cucumis sativus* L.)

Comment [b1]: c

Abstract

An experiment entitled **Effect of gibberellic acid and zinc sulphate on growth, yield and quality of cucumber (*Cucumis sativus* L.)** was conducted at Horticulture Research Field, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during February-May, 2023 on Cucumber. Experiment was laid out in factorial randomized block design with twenty treatment combinations and was replicated thrice. The experiment consisted of two factors. Factor H: Hybrids (TMCU-1107, TMCU-1125, TMCU-3112 and Saira-934) and Factor T: Different concentrations of GA + ZnSO₄ (T₀ - Control (water spray), T₁ - GA 100ppm + ZnSO₄ 0.25%, T₂ - GA 200ppm + ZnSO₄ 0.50%, T₃ - GA 300ppm + ZnSO₄ 0.75% and T₄ - GA 400ppm + ZnSO₄ 1%). TMCU-1125 recorded significantly higher yield attributes of cucumber over other hybrids in parameters like vine length at 80 DAS (102.6 cm), Days to 50% flowering (40), Days to 1st harvest (46), Number of fruits per plant (11), Fruit length (18.4 cm), Average fruit weight (237.5 g), Average fruit yield per plant (2.6 kg) and ascorbic acid content (0.8 mg/100g). Among different treatments, GA @300ppm + ZnSO₄ @0.75% resulted in significantly higher vine length at 80 DAS (98.4 cm), Days to 50% flowering (43), Days to 1st harvest (50), Number of fruits per plant (9), Fruit length (16.6 cm), Average fruit weight (232.0 g), Average fruit yield per plant (2.0 kg) and ascorbic acid content (0.5 mg/100g). The interaction effect of T₈ (TMCU-1125 + GA @300ppm + ZnSO₄ @0.75%) is best suited for getting higher growth, yield, quality of cucumber.

Comment [b2]: within inverted comma

Comment [b3]: c

Comment [b4]: d

Comment [b5]: d

Comment [b6]: n

Comment [b7]: f

Comment [b8]: a

Comment [b9]: a

Comment [b10]: d

Comment [b11]: n

Comment [b12]: f

Comment [b13]: a

Comment [b14]: a

Comment [b15]: ??? (interaction)

Comment [b16]: G

Keywords: gibberellic acid, zinc, growth, yield, and quality.

1. Introduction

Vegetable are important nutritive components of the daily diet because their nutritive value as a vital source of micronutrient has been well recognized. Vegetables play an important role in the balance diet by providing not only energy but also supplying vital protective nutrients either mineral and vitamins. Thus, vegetables are getting increasingly higher importance in India as well as in the world due to their relevance in achieving nutritional security from emerging nutritional problems in human beings. Today, India is the second largest producer of vegetables in the world after China. According to recommendations given by Indian Council of Medical Research (ICMR) an average man with vegetarian or non-vegetarian food habit should consume (300 g) vegetables per day, which include 125g leafy vegetable, 100g root vegetable and 75g other vegetables, but its availability of vegetable in India is only 225g.

Comment [b17]: b

Plant growth regulators can enhance the uptake of utilization of micronutrients in plant. certain PGRs can improve root growth and development, leading to increased exploration of soil for micronutrients. Foliar spray of plant growth regulators also promote growth, increases yield and quality in plants and micronutrients can serve as supplementary sources of essential elements for plants.

Comment [b18]: k

Comment [b19]: c

Gibberellins commonly known as gibberellic acids first came to the attention of western scientists in 1950s, they had been discovered much earlier in Japan. Gibberellic acid (also called gibberellin A3, GA, and GA3) is a hormone found in plants and fungi. Gibberellins (GAs) play important role in the development of lateral roots (LRs), which are critical for plant productivity. Therefore, it is of great importance for cucumber production to study the role of GAs in LR development. Here, the results showed that GAs regulated cucumber LR development in a concentration-dependent manner.

Zinc sulphate is the inorganic compound with the formula $ZnSO_4$. The addition of fertilizers to supplement natural soil fertility is a routine practice in modern agriculture, although temperate and tropical soils commonly remain deficient in micronutrients, particularly zinc (Zn). This work aimed to study the influence of foliar spray of zinc (as zinc sulphate) on growth parameters.

Comment [b20]: .

2. Materials and Methods

The investigation was carried out at the Horticulture Research Field, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj during Feb-May, 2023. The experiment was conducted in Factorial Randomized Block Design with 20 treatments in three replications viz. TC₁: H₁T₀ . TMCU-1107+control(waterspray), TC₂: H₁T₁ TMCU-1107+GA₃100ppm+ZnSO₄0.25%, TC₃: H₁T₁ TMCU-1107+GA₃200ppm+ZnSO₄0.5%, TC₄: H₁T₁ TMCU-1107+GA₃300ppm+ZnSO₄0.75%, TC₅: H₁T₁ TMCU-1107+GA₃400ppm+ZnSO₄1%, TC₆: H₂T₁ TMCU-1125+control(waterspray), TC₇: H₂T₁ TMCU-1125+GA₃100ppm+ZnSO₄0.25%, TC₈: H₂T₁ TMCU-1125+GA₃200ppm+ZnSO₄0.5%, TC₉: H₂T₁ TMCU-1125+GA₃300ppm+ZnSO₄0.75%, TC₁₀: H₂T₁ TMCU-1125+GA₃400ppm+ZnSO₄1%, TC₁₁: H₃T₁₀ . TMCU-3112+control(waterspray), TC₁₂: H₃T₁ TMCU-3112+GA₃100ppm+ZnSO₄0.25%, TC₁₃: H₃T₁ TMCU-3112+GA₃200ppm+ZnSO₄0.5%, TC₁₄: H₃T₁ TMCU-3112+GA₃300ppm+ZnSO₄0.75%, TC₁₅: H₃T₁ TMCU-3112+GA₃400ppm+ZnSO₄1%, TC₁₆: H₄T₁ Saira-934+control(waterspray), TC₁₇: H₄T₁ Saira-934+GA₃100ppm+ZnSO₄0.25%, TC₁₈: H₄T₁ Saira-934+GA₃200ppm+ZnSO₄0.5%, TC₁₉: H₄T₁ Saira-934+GA₃300ppm+ZnSO₄0.75%, TC₂₀: H₄T₁ Saira-934 +GA₃400ppm+ZnSO₄1%. with four

cucumber hybrids TMCU-1107, TMCU-1125, TMCU-3112 and Saira-934. Crop was planted with the spacing of 150×50 cm with the application of FYM@18 tonnes+NPK100:50:50 as basal dose along with spraying of GA₃+ZnSO₄ which was done at 15 and 30 days after emergence of two true leaf stage. The data was recorded for the following parameters viz. Vine length (cm), days to 50% flowering, days to first harvest, number of fruits per plant, avg. Number of picking, avg. fruit length (cm), average fruit weight (g), fruit girth (cm), average yield per plant (kg), total yield (t/ha), TSS(°Brix) and ascorbic acid content (mg/100g). TSS was measured with the help of hand refractometer and Vitamin C was measured by nitrilization method.

Comment [b21]: v

Comment [b22]: ??

3. RESULTS AND DISCUSSION

The data on yield attributes of cucumber after application of gibberellic acid and zinc sulphate was recorded and is presented in Table 1. Among different hybrids, Hybrid H2 (TMCU- 1125) recorded significantly longer vine at 80 DAS (102.6 cm), lesser number of days to 50% flowering (40), days to 1st harvest (46), number of fruits per plant (11), fruit length (18.4 cm), average fruit weight (237.5 g), average fruit yield per plant (2.6 kg) and ascorbic acid content (0.8 mg/100g) followed by vine at 80 DAS (101.3 cm), days to 50% flowering (41), days to 1st harvest (47), number of fruits per plant (10), fruit length (17.9 cm), average fruit weight (236.4 g), average fruit yield per plant (2.4 kg) and ascorbic acid content (0.7 mg/100g) in hybrid H1 (TMCU-1107) and shorter vine at 80 DAS (99.1 cm), more number of days to 50% flowering (42), late harvest (49), lower number of fruits per plant (9), lower fruit length (17.4 cm), average fruit weight (234.3 g), average fruit yield per plant (2.2 kg) and ascorbic acid content (0.6 mg/100g) in Hybrid H4 (Saira-934). Among different treatments, significantly longer vines at 80 DAS (107.1 cm), lesser number of days to 50% flowering (36), days to 1st harvest (42), number of fruits per plant (13), fruit length (19.9 cm), average fruit weight (244.2 g), average fruit yield per plant (3.2 kg) and ascorbic acid content (1.8 mg/100g) were recorded in (GA₃ @300 ppm + ZnSO₄ @0.75%), followed by vine at 80 DAS (104.9 cm), days to 50% flowering (40), days to 1st harvest (46), number of fruits per plant (12), fruit length (18.9 cm), average fruit weight (240.0 g), average fruit yield per plant (2.8 kg) and ascorbic acid content (0.8 mg/100g) in (GA₃ @200 ppm + ZnSO₄ @0.50%) and shorter vine at 80 DAS (92.3 cm), more number of days to 50% flowering (46), late harvest (52), lower number of fruits per plant (6), lower fruit length (16.0 cm), average fruit weight (227.3 g), average fruit yield per plant (1.5 kg) and ascorbic acid content (0.4 mg/100g) in control (water spray).

Comment [b23]: delete

Interaction data revealed that treatment T8 (TMCU-1125+ GA₃ @300 ppm + @0.75%) recorded significantly longer vine at 80 DAS (107.8 cm), lesser number of days to 50% flowering (35), days to 1st harvest (41), number of fruits per plant (14), fruit length (20.9 cm), average fruit weight (245.7 g), average fruit yield per plant (3.4 kg) and ascorbic acid content (1.3 mg/100g) followed by in T3 (TMCU-1107 + GA₃ @300 ppm + ZnSO₄ @0.75%) vines at 80 DAS (107.6 cm), days to 50% flowering (36), days to 1st harvest (42), number of fruits per plant (13), fruit length (19.9 cm), average fruit weight (244.8 g), average fruit yield per plant (3.3 kg) and ascorbic acid content (1.0 mg/100g) in hybrid H1 (TMCU-1107) and shorter vine at 80 DAS (88.4 cm), more number of days to 50% flowering (48), late harvesting (54), lower number of fruits per plant (5), lower fruit length (15.2 cm), average fruit weight (225.5 g), average fruit yield per plant (1.2 kg) and ascorbic acid content (0.3 mg/100g) in Hybrid H4 (Saira-934).

Comment [b24]: check

Significantly longer vine was recorded with the application of (GA₃ @300 ppm + ZnSO₄ @0.75%) which might be due to combined effect of gibberellic acid and zinc sulphate. Gibberellic acid promotes stem

Comment [b25]: Significantly

Comment [b26]: delete

elongation, leaf expansion, and overall plant growth. It stimulates cell division and elongation, which can lead to increased plant height and leaf size. It also helps in overcoming growth inhibition under stress conditions, thus promoting better growth even in unfavorable environments. While, zinc is an essential micronutrient required for various metabolic processes in plants, including photosynthesis, enzyme activity, and hormone synthesis. It plays a crucial role in the synthesis of auxins, which are important for root development, flowering, and fruiting. The combination of gibberellic acid and zinc sulphate ensures that cucumber plants have optimal conditions for growth, development, and productivity. Thereby promoting better overall plant health, yield, and fruit quality of cucumber.

Comment [b27]: W

UNDER PEER REVIEW

Table1.Effect of gibberellic acid and zinc sulphate of different cucumber hybrids ongrowth, yieldandqualityof cucumber

Hybrids → Treatments ↓	Vinlength at 80 DAS					Days to50%flowering					Days to first harvest					No. offruits/ plant				
	H1	H2	H3	H4	Mean T	H1	H2	H3	H4	Mean T	H1	H2	H3	H4	Mean T	H1	H2	H3	H4	Mean T
T0	93.4	96.9	90.6	88.4	92.3	45	44	46	48	46	52	51	53	54	52	7	7	6	5	6
T1	98.6	99.8	97.9	97.2	98.4	43	42	43	44	43	50	49	50	51	50	9	9	8	8	9
T2	105.4	105.6	104.9	103.6	104.9	40	39	40	41	40	46	44	46	47	46	12	12	12	11	12
T3	107.6	107.8	107.0	105.9	107.1	36	35	37	38	36	42	41	42	44	42	13	14	13	13	13
T4	101.7	102.7	100.6	100.3	101.3	41	41	42	42	41	48	47	48	49	48	10	11	10	10	10
Mean H	101.3	102.6	100.2	99.1		41	40	42	42		47	46	48	49		10	11	10	9	

Vinlength at 80 DAS			Days to50%flowering			Days to first harvest			No. offruits/ plant		
Factors	SE(d)±	CD0.05	Factors	SE(d)±	CD0.05	Factors	SE(d)±	CD0.05	Factors	SE(d)±	CD0.05
Factor(H)	0.018	0.036	Factor(H)	0.043	0.088	Factor(H)	0.054	0.110	Factor(H)	0.018	0.036
Factor(T)	0.020	0.040	Factor(T)	0.048	0.098	Factor(T)	0.060	0.123	Factor(T)	0.020	0.040
Factor(HX T)	0.039	0.080	Factor(HX T)	0.097	0.197	Factor(HX T)	0.121	0.245	Factor(HX T)	0.039	0.080

Hybrids →	Fruit length (cm)					Averagefruit weight (g)					Averagefruit yield /plant (kg)					Ascorbicacid content(mg/100g)				
	H1	H2	H3	H4	Mean T	H1	H2	H3	H4	Mean T	H1	H2	H3	H4	Mean T	H1	H2	H3	H4	Mean T
T0	16.3	16.5	15.9	15.2	16.0	227.7	229.4	226.5	225.5	227.3	1.5	1.7	1.4	1.2	1.5	0.4	0.5	0.3	0.3	0.4
T1	16.5	16.9	16.4	16.7	16.6	232.4	233.4	231.7	230.6	232.0	2.1	2.1	1.9	1.8	2.0	0.5	0.5	0.4	0.4	0.5
T2	19.0	19.2	18.8	18.6	18.9	240.5	241.5	239.5	238.5	240.0	2.9	2.9	2.8	2.7	2.8	0.9	0.9	0.9	0.8	0.8
T3	19.9	20.9	19.6	19.3	19.9	244.8	245.7	243.7	242.5	244.2	3.3	3.4	3.1	3.1	3.2	1.0	1.3	0.9	0.9	1.8
T4	17.9	18.4	17.2	17.2	17.8	236.4	237.4	235.5	234.5	236.0	2.4	2.6	2.2	2.2	2.4	0.6	0.8	0.7	0.6	0.7
Mean H	17.9	18.4	17.6	17.4		236.4	237.5	235.4	234.3		2.4	2.6	2.2	2.2		0.7	0.8	0.6	0.6	

Fruit length (cm)			Averagefruit weight (g)			Averagefruit yield /plant (kg)			Ascorbicacid content(mg/100g)		
Factors	SE(d)±	CD _{0.05}	Factors	SE(d)(±)	CD _{0.05}	Factors	SE(d)±	CD _{0.05}	Factors	SE(d)±	CD _{0.05}
Factor(H)	0.012	0.024	Factor(H)	0.104	0.051	Factor(H)	0.004	0.009	Factor(H)	0.023	0.046
Factor(T)	0.013	0.026	Factor(T)	0.116	0.057	Factor(T)	0.005	0.010	Factor(T)	0.026	0.052
Factor(HX T)	0.026	0.053	Factor(HX T)	0.233	0.114	Factor(HX T)	0.010	0.019	Factor(HX T)	0.051	0.104

Note: H1- (TMCU-1107), H2-(TMCU-1125), H3- (TMCU- 3112) and H4 - (Saira- 934)

T-Different concentrations of gibberellic acid and zinc sulphate

Conclusion

Based on the results of the present investigation entitled “Effect of gibberellic acid and zinc sulphate on growth, yield and quality of cucumber (*Cucumis sativus* L.)” it is concluded that among the different treatment combinations of hybrids, gibberellic acid and zinc sulphate, treatment T8 (TMCU - 1125 + GA3 @300 ppm + ZnSO4 @0.75%) recorded significantly better results in vine length, days to 50% flowering, days to 1st harvest, number of fruits per plant, fruit length, average fruit weight, average fruit yield per plant, and ascorbic acid content as well as significantly higher Benefit Cost Ratio of 3.41.

References

- Dalai, S., Singh, M.K., Singh, K.V., Kumar, M., Malik, S. and Kumar, V. (2015).** Effect of foliar application of GA₃ and NAA on growth, flower-ingyield and yield attributes of cucumber [*Cucumis Sativus* L.]. *Annals of Horticulture*, **8**(2):181-194.
- Eidani, A. and AL-Shibani, J. (2019).** Effect of biofertilizer with trichodermaharzianu Fungus and zinc foliar spraying on cucumber plant (*Cucumis Sativus* L.) and Growth Characters.*Al-Qadisiyah Journal For Agriculture Sciences*, **9**(2):.215-224.
- P Pal, P., Yadav, K., Kumar, K. and Singh, N., (2016).** Effect of gibberellic acid and potassium foliar sprays on productivity and physiological and biochemical parameters of parthenocarpic cucumber cv. ‘Seven Star F’. *Journal of Horticultural Research*,**24**(1):.93-100.
- Rahman, M.A., Sikder, S., Bahadur, M.M. and Pramanik, S.K. (2020).** Influence of gibberellic acid (GA₃) on growth, flowering, and fruit yield of cucumber. *Journal of Science and Technology*, **18**(2):.33-42
- Sidhu, M.K., Raturi, H.C., Singh, S.K., Bora, L. and Shukla, Y.R. (2020).** Effect of foliar application of boron, iron and zinc on growth, yield and economics of cucumber grown under naturally ventilated polyhouse condition. *Vegetable Science*, **47**(1):.110-115.
- Sinha, K.P., Kumar, A., Afreen, S., Kumari, A., Singh, V.K. and Singh, S.P. (2013).**Study on effectofmicronutrientsand plant growthregulatoronproductionofqualityseedincucumber(*Cucumissativus* L.).Seedling,p.5.
- Toman, S.S., Al-Taey, D.K.A., Al-Tawaha, A.R., Sirajuddin, S.N., Rasyid, I. and Hassan, A.A.A.H. (2020),** April. Effect of foliar application and mineral fertilizer on growth parameters and content auxins, GA and CK in cucumber leaves. In IOP Conference Series: Earth and Environmental Science (Vol. **492**(1): 012009). IOP Publishing

Comment [b28]: (data)

Comment [b29]: (data)

Comment [b30]: (data)

Comment [b31]: (data)

Comment [b32]: (data)

Comment [b33]: (data)

Comment [b34]: (data)

Comment [b35]: flowering

Comment [b36]: harzianum

Comment [b37]: italics