

Original Research Article

OPTIMIZATION OF KALMEGH (*Andrographis paniculata* Nees.) CULTIVATION THROUGH PLANT SPACING AND NUTRIENT MANAGEMENT

Abstract:

Kalmegh is being used from the immortal time for treating the cold, fever, snake bite, diabetics, liver diseases etc. An field experiment was conducted during *kharif* season of 2019-2020 at Horticulture Farm, Main Agricultural Research Station, College of Agriculture, University of Agricultural Sciences, Raichur, Karnataka, India, to study the response of plant geometry and integrated nutrient management on growth and yield of kalmegh. The experiment was laid out in a spilt plot design with 15 treatments and 3 replications. The results revealed that spacing of 30cm x 45 cm with imposition of 50% RDN + 50% vermicompost equivalent to recommended N found to be significantly higher number of primary branches and leaves per plant, leaf area and dry matter accumulation over the other treatments. The fresh and dry herbage yield per hectare were resulted in the significantly higher in the spacing of 30cm x 15 cm with the application of 50% RDN + 50% vermicompost equivalent to recommended N.

Key words: Spacing, application, kalmegh

Introduction:

India is known to be “Home for Medicinal Plants” as it grown more than 8000 species of medicinal plants. From ancient time the medicinal plants are used for treatment of many diseases by tradition knowledge (TK). Traditional knowledge (TK) associated with medicinal herbs and cultivation, innovation and preservation of medicinal herbs is a highly gendered activity in most countries. Kalmegh (*Andrographis paniculata* Nees.) belonging to family Acanthaceae which is commonly known as “King of Bitters” because of bitter taste. Its major biological activity is antidiabetic, but it has also been reported to have anti-angiogenetic, antibacterial, anti-cancer anti-inflammatory, antimalarial, antioxidant and hepatoprotective activities. A recent study has been documented that, kalmegh is having anti-HIV properties due

to presence of important active constituent Andrographolide (Sajwan, 2008). The therapeutic activities of the plant are attributed to andrographolide and diterpenides. The whole herb is the source of several diterpenoids of which andrographolide is important and is distributed all over the plant in different proportions. The leaves contain the maximum andrographolide (2.50%) compare to stem (2%) and capsule (0.15%) (Makwana *et al.*, 2010). The use and marketing demand of herbal medicine is increasing for the safety and prevention concerns of human health. Commercial cultivation of medicinal plants with good agriculture practices is most important. In this concern to meet the demand of growing population with good quality of produce has been achieved through the good agriculture practices as well as good agriculture management. In this view, the investigation has been conducted to different plant geometry with optimize the nutrient supply through integrated use of organic and inorganic fertilizers for higher yield per unit area of kalmegh.

Material and methods:

Experimental site:

An investigation was conducted during *kharif* season of 2019-2020 at Horticulture Farm, Main Agricultural Research Station, College of Agriculture, University of Agricultural Sciences, Raichur, Karnataka, India which is situated in North Eastern Dry Zone (Zone-II) of Karnataka at 16° 15' N latitude, 77° 21' East longitudes at an altitude of 400 m above mean sea level. The soil of the experimental site was sandy clay loam in texture having high organic carbon, low available nitrogen, high available phosphorus and medium in exchangeable potassium with slightly alkaline in reaction.

Experimental Design and Details of treatments

The experiment was laid out in Split plot design with 3 replications. There are 15 treatment combinations comprising of three main plots as different spacing *viz.*, $S_1 = 30\text{cm} \times 15\text{ cm}$, $S_2 = 30\text{cm} \times 30\text{ cm}$, $S_3 = 30\text{cm} \times 45\text{ cm}$ and five subplots as different organic and inorganic fertilizer combinations *viz.*, $F_1 = 100\% \text{ RDF } (75:75:50 \text{ NPK kg ha}^{-1})$, $F_2 = \text{FYM alone equivalent to recommended N}$, $F_3 = \text{Vermicompost alone equivalent to recommended N}$, F_4

= 50% RDN + 50% FYM equivalent to recommended N and $F_5 = 50\% \text{ RDN} + 50\%$ Vermicompost equivalent to recommended N.

Crop measurements:

Seeds were sown in nursery, before sowing the seeds were soaked in water for 12 hours than treated with Carbendazim (2g kg^{-1} of seeds) and 40 – 45 days old seedlings were used for transplanting. All the standard recommended cultivation practices were followed during the cultivation of kalmegh. Observation on growth parameters were recorded on five randomly selected plants in each replication of different treatments at 30, 60 and 90 days of transplanting (DAP). Harvesting of kalmegh was done at maturity (110 DAT).

Statistical analysis

The data collected on different parameters during the course of investigation were subjected to Fisher's method of analysis of variance and interpretation of data was done as per the procedure described by Panse and Sukhatme (1985). The level of significance used in 'F' test was $p = 0.05$.

Results and discussion:

Growth attributes of the kalmegh as influenced by the geometry and integrated nutrient management are presented in the Table 1(a) and 1(b). The growth pattern was progressively increasing during all the growing stages with respect to all the growth parameters. The significantly taller plants (56.62 cm) and higher leaf area index (1.61) was observed in the narrow spacing with imposition of 50% RDN + 50% vermicompost equivalent to recommended N (S_1F_5) over S_1F_1 . The taller plants were noticed under narrow spacing compare to wider spacing. The narrow spaced plants grown vertically than horizontal might be due to more competition for the radiation to rescue the photosynthetic demand of plant and these plants grow taller by using the basic resources such as water, sunlight and nutrients. The same results have been observed by Tansi *et al.*, 2017 in stevia. The higher leaf area index was observed in narrow spacing which might be due to the lesser ground area of the crop leading to the higher leaf area index where as

more ground area of crop resulted in the lesser leaf area index (Anupama *et al.*, 2017 in kasurimethi).

The higher number of primary branches per plant (24.49), number of leaves per plant (150.78), leaf area (814.19cm²) and dry matter accumulation (36.10 g plant⁻¹) was noticed with

UNDER PEER REVIEW

Table 1(a). Influence of plant geometry and integrated nutrient management on growth attributes of kalmegh

Treatments	Plant height (cm)						Number of Primary Braches per plant						Number of leaves per plant					
	F ₁	F ₂	F ₃	F ₄	F ₅	Mea n	F ₁	F ₂	F ₃	F ₄	F ₅	Mea n	F ₁	F ₂	F ₃	F ₄	F ₅	Mea n
	S ₁	48.	54.	55.	54.	56.	53.9	16.	21.	21.	22.	24.	21.2	120.	130.	131.	133.	142.
	80	00	94	21	62	1	54	20	41	79	17	2	51	9	57	81	03	76
S ₂	47.	54.	54.	55.	56.	53.7	17.	22.	22.	23.	24.	22.0	123.	131.	136.	139.	141.	134.
	94	44	43	73	15	4	60	14	74	45	28	4	56	66	97	34	07	52
S ₃	47.	49.	51.	52.	54.	51.0	17.	22.	24.	23.	24.	22.5	125.	144.	146.	148.	150.	142.
	07	33	4	70	70	4	17	95	38	49	49	0	63	01	09	42	78	99
Mean	47.	52.	53.	54.	55.		17.	22.	22.	23.	24.		123.	135.	138.	140.	144.	
	94	59	92	21	82		10	1	84	24	31		23	52	21	52	63	
	S.Em.±		C.D. at 5%				S.Em.±		C.D. at 5%				S.Em.±		C.D. at 5%			
Main plot (S)	0.51		2.00				0.18		0.71				1.00		3.93			
Subplot (F)	0.28		0.82				0.20		0.58				0.75		2.19			
F at same level of S	0.48		1.41				0.35		1.01				1.30		3.80			
S at same or different	3.34		11.72				1.79		5.7				7.68		25.70			

Table 1(b). Influence of plant geometry and integrated nutrient management on growth attributes of kalmegh

Treatments	Leaf area (cm ²)					Mean	Leaf area index					Mean	Dry matter accumulation (g plant ⁻¹)					Mean
	F ₁	F ₂	F ₃	F ₄	F ₅		F ₁	F ₂	F ₃	F ₄	F ₅		F ₁	F ₂	F ₃	F ₄	F ₅	
S ₁	602.	667.	684.	709.	766.	686.	1.3	1.4	1.5	1.5	1.6	1.48	22.	23.	23.	22.	23.	23.1
	54	57	17	19	97	09	2	3	1	1	1		51	88	47	78	29	8
S ₂	617.	671.	712.	738.	761.	700.	0.8	0.9	0.9	0.9	0.9	0.90	22.	27.	28.	28.	27.	27.0
	82	47	23	48	79	36	0	0	0	3	6		66	44	27	95	78	2
S ₃	628.	734.	759.	786.	814.	744.	0.5	0.5	0.5	0.6	0.6	0.58	31.	33.	33.	34.	36.	34.0
	16	44	69	61	19	62	4	7	9	0	2		83	71	75	9	10	6
Mean	616.	691.	718.	744.	780.		0.8	0.9	1.0	1.0	1.0		25.	28.	28.	28.	29.	
	17	16	7	76	99		9	7	0	1	6		66	35	49	88	06	
	S.Em.±		C.D. at 5%			S.Em.±		C.D. at 5%			S.Em.±		C.D. at 5%					
Main plot (S)	5.24		20.57			0.02		0.09			0.78		3.06					
Subplot (F)	3.95		11.54			0.01		0.03			0.47		1.38					
F at same level of S	6.85		19.98			0.02		0.06			0.82		2.38					
S at same or different level of F	40.29		134.74			0.15		0.52			5.32		18.47					

Table 2. Influence of plant geometry and integrated nutrient management on yield of kalmegh

Treatments	Fresh herbage yield (t ha ⁻¹)						Dry herbage yield (t ha ⁻¹)					
	F ₁	F ₂	F ₃	F ₄	F ₅	Mean	F ₁	F ₂	F ₃	F ₄	F ₅	Mean
S ₁	9.84	10.62	11.66	13.47	15.81	12.28	3.78	4.62	5.80	6.11	7.37	5.54
S ₂	5.52	6.21	7.21	8.07	8.90	7.18	2.85	2.85	4.42	4.20	4.20	3.70
S ₃	4.16	5.10	6.69	6.66	6.91	5.91	1.38	1.92	2.48	2.99	3.06	2.32
Mean	6.51	7.31	8.52	9.40	10.54		2.67	3.13	4.23	4.43	4.88	
		S.Em.±		C.D. at 5%				S.Em.±		C.D. at 5%		
Main plot (S)		0.32		1.28				0.08		0.31		
Subplot (F)		0.17		0.50				0.07		0.20		
F at same level of S		0.30		0.86				0.12		0.34		
S at same or different level of F		2.09		7.38				0.66		2.16		

wider spaced plant with the consortia of application of 50% RDN + 50% vermicompost equivalent to recommended N (S_3F_5) over S_1F_1 . This might be due the increases in the photosynthetic activity due to penetration of more sunlight on plant leads to vigorous growth of the herbage (Pal *et al.*, 2019 in kalmegh). Application chemical fertilizer in combination with vermicompost resulted higher uptake of the nutrient (especially nitrogen) which enhances the cell multiplication, elongation and differentiation leads to increased plant growth which might be results in more number of branches and leaves per plant (Lokesh *et al.*, 2018 in coriander). As number of leaves shown higher in wider spacing due to more light intensity, surface area, less competition for nutrients leads to higher leaf area also with the high chlorophyll content and leaf production which produced due to the application nitrogen application ultimately helps increasing the photosynthetic activity (Vahidi *et al.*, 2018 in henna and Jat *et al.*, 2021 in Ashwagandha). Dry matter accumulation was increased progressively with the advancement in plant age till the maturity found to be better translocation of carbohydrate and their utilization for the production of more number of leaves and number of branches reported by Divya *et al.* (2018) and Shivani *et al.* (2023) in kalmegh.

The increment in the number of branches and number of leaves per plant resulted in the higher fresh and dry herbage yield (Table 2). Narrow spaced plants with application found to be higher 50% RDN + 50% vermicompost equivalent to recommended N (S_1F_5) noticed the significantly higher fresh and dry herbage yield per hectare (15.81 and 7.37 kg ha⁻¹) over S_3F_1 . The concept of HDP followed through 30 x 15 cm which accommodated more number of plants per unit area would result in more fresh and dry herbage yield per unit area. Increased biomass also noticed by the integrated application of 50% RDN + 50% vermicompost equivalent to recommended N to supply the N usage of crop would increased the better growth parameters, herbage yield of good quality in kalmegh. The higher dry matter accumulation attributed due to the more number of leaves and branches per plant owing to availability of more nutrient continuously from organic and inorganic source of nutrient resulted in the increased fresh herbage yield which supported with the findings of Himanshu *et al.* (2018) and Amala *et al.*, 2023 in kalmegh. Similar findings are noticed by Cheena *et al.* (2020), Shwetha *et al.* (2021), Shakywa *et al.* (2022) and Sadbhawana *et al.* (2024) in kalmegh.

Conclusion: Medicinal plants are boon of nature and demand for herbal extracts is progressively increasing in developing countries. Production and productivity of the crop can be increased by the adoption of standardized production practices. The adoption of spacing of 30 x 15 cm with the consortia of application 50% vermicompost and 50% recommended N resulted in the higher number of branches and number of leaves per plant resulted in the higher fresh and dry herbage yield potentially offers improved soil health or sustainability compared to inorganic fertilizers alone.

References:

1. Amala D, Raj Kumar M, Kiran Kumar A, Naveen Kumar B, Gouthami P and Sathish G, Effect of Organic Manures and Biofertilizers on Quality and Soil Analysis of Kalmegh (*Andrographis paniculata* Wall. Ex.Nees.) Var. CIM Megha. Int. J. Plant Soil Sci., 2023;35:696-703.
2. Anupama G, Hegde LN., Hegde NK., Devappa V, Mastiholi, A. B. Nishani S. Effect of nitrogen and spacing levels on physiological and yield parameters of kasurimethi (*Trigonella corniculata* L.) var. Pusa Kasuri. Int. J. Curr. Microbiol. App. Sci.2017;6(9):723-733.
3. Cheena J, Krishna Veni V, Padma M and Sreenivas M. Improvement in the Growth and Yield of Kalmegh influenced by Pruning and Nutrition levels. International Journal of Current Microbiology and Applied Sciences. 2020;9(12):203-212.
4. Himanshu V, Negi MS, Archana J, Babita B, Anil S, Mahapatra BS. Jai P. Growth attributes of kalmegh (*andrographis paniculataburm.f.*) as influenced by integrated nutrient management under Tarai conditions of Uttarakhand. Int. J. Chem. Stud. 2018;6(5): 2947-2949.
5. Jat RS, Basak BB, Gajbhiye NA. Organic manures and biostimulants fostered soil health and increased the harvest quality of the medicinal herb ashwagandha. Agronomy journal. 2021;113:504-514.

6. Lokesh B, Ramar A, Jegadeeswari V. Effect of biostimulant and growth regulators on herbage yield and quality in coriander var. CO (CR) 4. *Int. J. Chem. Stud.* 2018;6(5): 1373-1376.
7. Makwana PD, Patel DH, Patel JJ, Patel, HK. Effect of different organic manures and spacing on yield and yield attributes of Kalmegh-Panchang (*Andrographis paniculata* Wall. Ex. Nees.) under middle Gujarat Conditions. *Int. J. Plant Sci.* 2010;5(1): 30-32.
8. Panse VG, Sukhatme, PV. *Statistical methods for agricultural workers II*nd Enlarged Ed. ICAR., New Delhi. 1985;135-136.
9. Sadbhawana J, Kailash CM, Om Singh and Patidar DK. Responses of INM on growth and yield of Kalmegh. *Med. Plants - Int. J. Phytomed.* 2024;16(1):86-91.
10. Sajwan, B. S. Medicinal plants scenario in india-government initiatives towards resource security and quality assurance for increased trade and commerce. In: International conclave on medicinal plants for asean and bimstec countries Lamphelpat, held at Imphal, India. 2008;11-25.
11. Shakywa SK, Meena KC, Soni N, Patidar DK and Patidar BK. Response to Organic Manures and Plant Geometry in Kalmegh (*Andrographis paniculata* Nees.): Way to Reduce Exploitation of Forest. *Annals of Forest Research.* 2022;65(1):7633-7641.
12. Shivani C, Meena KC, Nitin S, Patidar DK, Kachouli BK, Patidar BK and Ajay H. Response of recommended dose of fertilizers with organic manures on growth, yield and economics of kalmegh (*Andrographis paniculata* Nees.): A way to reduced use of chemical Fertilizers. *Pharma Innovation.* 2023; 12(3): 119-124.
13. Shwetha C, Padma E, Naram Naidu L, Krishna U. Effect of integrated nutrient management on growth and yield of Kalmegh (*Andrographis paniculata* Nees) variety CIM Megha. *Pharma Innovation.* 2021;10(8):835- 838.
14. Tansi LS, Samadpourrigani E, Gedik S. Effects of different plant density and cutting times on yield of stevia under the cukurova conditions. *Int. J. Sec. Metabol.* 2017;4(3):355-358.
15. Vahidi A, Alizadeh A, Baghizadeh A, Ansari H. Effect of mycorrhiza, vermicompost and chemical fertilizer application on yield and lawson content of henna as medicinal plant under water deficit condition. *J. Water Research in Agric.* 2018;32(1):107-120.

UNDER PEER REVIEW