

Original Research Article

PERFORMANCE OF *Beauveria bassiana* (Balsamo) Vuillemin STORED AS OIL CULTURES UNDER DIFFERENT TEMPERATURE REGIMES

ABSTRACT:

Oil cultures of *Beauveria bassiana* (Balsamo) Vuillemin were prepared from the laboratory-maintained isolate by using two vegetable oils viz., Rice Bran oil, Sesamum oil and two mineral oils viz., Liquid Paraffin oil and Heavy grade Mineral Oil. Prepared Oil cultures were stored under three different temperature regimes viz., at room temperature ($28 \pm 3^{\circ}\text{C}$), refrigerated condition (4°C) and freezing temperature (-20°C). These 12 cultures stored under different temperatures along with crude culture and untreated control were evaluated for viability and virulence against third instar *Spodoptera litura* (Fabricius) larva on the day of preparation and upto 180 days under laboratory conditions. Among all oil cultures, Sesamum oil cultures and Liquid Paraffin oil cultures stored at -20°C performed well with highest conidial germination of 87.85% and 83.57% and highest larval mortality of 90.00% and 86.67% on the day of preparation (0 DAS). After 180 DAS of storage, for the above two formulations conidial germination was in the range of 62.87% to 61.43% and larval mortality was 60.00%. Comparatively, least performance was observed in rice bran oil formulation stored at room temperature with conidial germination and larval mortality of 77.14% and 80.00% (0 DAS) and 41.43% and 40.00% (180 DAS) respectively.

Usage of oils for the preservation of entomopathogenic fungal based formulations prevent conidial desiccation and improve adhesion and afford protection to fungal conidia from the UV of sunlight improving field performance.

Keywords:

Beauveria bassiana, oil cultures, vegetable oils, mineral oils, different temperature regimes, storage, viability, virulence.

1. INTRODUCTION

According to FAO's 2022 report, India consumed over 61,000 tonnes (t) of pesticides in 2020 (FAO, 2022). These pesticides in addition to creating pesticide residues, environmental pollution issues and other issues, the widespread use of synthetic pesticides inevitably lead to the development of insecticide resistance, harmful effect on natural enemies, insecticide-induced resurgence of the pests and biodiversity decline (Dong *et al.*, 2021; Waqas *et al.*, 2021; Guo *et al.*, 2020). Considering these limitations, environmentally benign pest management methods have become increasingly popular to rebuild the balance of cropland ecosystems for the sustainable control of crop pests (Yan li *et al.*, 2022). Pest management by biocontrol agents are assuming prominence and have been considered as an important strategy in insect population reduction (Ummidi and Vadlamani, 2014). Among biocontrol agents, microbial pathogens such as bacteria, fungi, viruses, nematodes and protozoans are promising agents for the effective control of insects.

Entomopathogenic fungi are regarded as the most significant among the numerous microbial agents due to their wide host range, eco-friendly nature, effectiveness against target pest *etc.* They are cheaper in long run, show lesser residual effects, and are able to overcome the problem of resistance (Sharma *et al.*, 2023). Entomopathogenic fungi are a specialized group of soil-dwelling

microbial organisms that infects and kill insects and other arthropods through cuticle penetration, with traits and modes of action that render them as effective biopesticides that they play a crucial role in insect pest management (Mantzoukaset *et al.*, 2022). EPF is advantageous in pest control because direct ingestion of fungal propagules is not needed by insects, thus also becoming active against the non-feeding stages of insects (Bateman *et al.*, 1998). They cause insect mortality by nutritional deficiency, destruction of tissues and by release of toxins. The entry of entomopathogenic fungi through the insect cuticle occur by a combination of mechanical pressure and enzymatic degradation (Ramanujam *et al.*, 2014). They have the ability to synthesize extracellular cuticle-degrading enzymes like chitinase, protease and lipase that work together to overcome the chitinous and proteinaceous components of the insect cuticle, penetrating the germ tube inside thus, enabling hyphal access to the haemolymph and colonizing the insects to death (Gebremariam *et al.*, 2022; Seema *et al.*, 2013; Beys-da-Silva *et al.*, 2014). As, they have a complex mode of action that makes it difficult for pests to develop resistance against them (Dusfour, 2019).

Several species of entomopathogenic fungi have now been industrialized (de Faria and Wraight, 2007). According to a recent report, out of 171 commercial EPF products, majority of them were based on *Beauveria bassiana* (33.9 %) followed by *Metarhizium anisopliae* (33.9 %), *Isaria fumosorosea* (5.8%) and *B. brongniartii* (4.1%) (Islam *et al.*, 2021). *Beauveria bassiana* is the asexual form (anamorph) of *Cordyceps bassiana* fungus that infects a huge variety of insects, which is used to control crop infestations caused by aphids, thrips and whiteflies (Money, 2016) as well as lepidopteran pests (Singh, 2015). Among the cyclic hexadepsipeptide mycotoxins produced by the different EPF, beauvericin produced by them have shown the most effective larvicidal properties (Wang *et al.*, 2012). Even though, effective strains were isolated there was still problem with the development of suitable cost effective formulations which could overcome harsh environmental conditions and ensure biological and chemical stability as well as viability during storage for extended periods (Rachappa *et al.*, 2005).

One among the technique is the usage of oils for the preservation of entomopathogenic fungi based formulations. It has been suggested that oil formulations can prevent conidial desiccation and improve adhesion of conidia to the hydrophobic surface of insect cuticle and enabling the opportunistic spread of conidia in to high humidity locations such as the inter-segmental membranes (Inyang *et al.*, 2000; Devi and Prasad, 1996; Bateman *et al.*, 1993). In field conditions, higher temperature, lower humidity and Ultraviolet radiation (UV) exposure pose detrimental effect on fungal conidia. This situation warrants shift to the use of oil based formulations which showed good control of insect pests under field condition (Batta, 2003). Oil affords protection to fungal conidia from the UV of sunlight thereby improvement in the field performance (Moore *et al.*, 1993).

While, cryoprotectant preservation at -196°C submerged in liquid nitrogen following freezing at a controlled rate is generally accepted as the optimal storage technique for preserving both cell viability and biological properties *i.e.* infectivity and virulence, it is considerably expensive and time consuming which is not affordable by many laboratories (Humber, 1997; Lastra *et al.*, 2002).

Maintenance at optimum temperature for long term preservation of EPF culture is of utmost importance.

Therefore, it is necessary for the development of suitable and affordable formulation technologies for the proper maintenance and preservation of different entomopathogenic fungal cultures. There have been relatively fewer studies conducted on optimising the storage conditions for *Beauveria bassiana* oil cultures, which could remain viable for extended periods and be suitable for later retrieval. Therefore, the present study was undertaken to evaluate the performance of *B. bassiana* stored as oil cultures under different temperature regimes.

2. MATERIAL AND METHODS

Place and duration of study: Insect Pathology laboratory, Central Instrumentation Laboratory and Insectary, Department of Entomology Sri Venkateswara Agricultural College, Tirupati during between September, 2023 and April, 2024.

2.1 Preparation of *Beauveria bassiana* oil cultures

For preparation of oil cultures, 4 oils were chosen viz., 2 vegetable oils (Rice bran oil, Sesamum oil) and 2 mineral oils (Heavy grade Mineral oil, Liquid paraffin oil). Sabouraud's Dextrose Agar with Yeast Extract Medium (SDAY). was prepared, poured into test tubes, autoclaved at 121°C at 15 Psi for 15 minutes and made as slants. After slants got cooled down, discs or loop of spores and mycelia of laboratory maintained culture of *B. bassiana* were inoculated into slants under aseptic condition and incubated at 25 ± 2°C temperature. Ten to twelve days after inoculation, satisfactory growth and sporulation of *B. bassiana* were obtained in almost all slants (Fig. 1). Each fungal slant was poured with autoclaved vegetable and mineral oils viz., Rice bran oil, Sesamum oil, Mineral oil and Liquid paraffin oil respectively until the entire fungi grown in slant media got submerged with oil (Fig. 2).



Fig. 1. Slant culture of *B. bassiana*



Fig. 2. *B. bassiana* oil cultures

2.2 Storage of *Beauveria bassiana* oil cultures at different temperature regimes

The prepared oil cultures of *B. bassiana* with Rice bran oil, Sesamum oil, Mineral oil and Liquid paraffin oil were maintained at room temperature ($28 \pm 3^{\circ}\text{C}$), refrigerated condition (4°C) in Refrigerator at Insect Pathology laboratory and freezing temperature (-20°C) in Deep Freezer at Central instrumentation Lab of the college.

A total of treatments 14 treatments (including control) of *B. bassiana* oil cultures were prepared.

- T₁: Mineral oil culture of *B. bassiana* stored at Room temperature (28 ± 3°C)
T₂: Mineral oil culture of *B. bassiana* stored at Refrigerator temperature (4°C)
T₃: Mineral oil culture of *B. bassiana* stored at freezing temperature (-20°C)
T₄: Liquid paraffin oil culture of *B. bassiana* stored at Room temperature (28 ± 3°C)
T₅: Liquid paraffin oil culture of *B. bassiana* stored at Refrigerator temperature (4°C)
T₆: Liquid paraffin oil culture of *B. bassiana* stored at freezing temperature (-20°C)
T₇: Rice bran oil culture of *B. bassiana* stored at Room temperature (28 ± 3°C)
T₈: Rice bran oil culture of *B. bassiana* stored at Refrigerator temperature (4°C)
T₉: Rice bran oil culture of *B. bassiana* stored at freezing temperature (-20°C)
T₁₀: Sesamum oil culture of *B. bassiana* stored at Room temperature (28 ± 3°C)
T₁₁: Sesamum oil culture of *B. bassiana* stored at Refrigerator temperature (4°C)
T₁₂: Sesamum oil culture of *B. bassiana* stored at freezing temperature (-20°C)
T₁₃: Unformulated culture of *B. bassiana* (Not preserved in oil)
T₁₄: Untreated control

2.3 Evaluation of *B. bassiana* oil cultures stored under different temperature regimes

The viability of *B. bassiana* conidia and its efficacy was studied on the day of preparation (0 DAS) and at monthly intervals upto 180 days (DAS).

2.3.1 Assessment of viability of oil cultures

Serial dilutions of 1×10^5 spores ml⁻¹ of all oil cultures were employed for viability experiments. Spore count was taken with the help of haemocytometer. Two to three drops of spore suspension were placed in the cavity slide. The cavity slide was prepared by arranging moistened cotton in petriplates and incubated in the incubator at 22°C. After 24 hrs, the spore suspension was observed under microscope for counting total number of spores and number of germinated spores. The germination percentage of *Beauveria bassiana* conidia at different treatments were calculated.

2.3.2 Assessment of virulence of oil cultures against third instar *Spodoptera litura* larvae

From each treatment, 0.5 ml of oil culture was poured into 100 ml of distilled water and added with 100µl of Triton- X 100 (0.1 %) into conical flasks to make spray suspension. Triton- X 100 (0.1 %) as a wetting agent for uniform dispersion of spores with oil. Spore suspensions of 1×10^7 spores per ml were standardized after assessing the number of spores in the suspension with an improved Neubauer haemocytometer. Spray suspensions were sprayed on castor leaves with atomizer and air dried. Freshly moulted third instar larvae were allowed to feed on them. The treatments were replicated thrice to confirm the reproducibility of the results. The bioassays were conducted at room temperature under laboratory conditions. From next day, larvae were fed with fresh castor leaves. Daily observations on post treatment changes in larvae and larval mortality were recorded.

2.4 Analysis of the Data

All the recorded observations *i.e.* conidial viability and larval mortality were converted to percentage values. The percentage values were converted to angular transformed values. The observations were statistically analysed by using SPSS v 16 software. Data were subjected to analysis of variance (ANOVA) at $P < 0.01$ level of significance. Means were compared by Duncan's Multiple Range Test (DMRT) (Saheb *et al.*, 2021).

The conidial germination was expressed as per cent conidial viability by using the formula;

$$\text{Per cent conidia viability} = \frac{\text{Number of conidia germinated}}{\text{Total conidia}} \times 100$$

The larval mortality was expressed as per cent larval mortality by using the formula;

$$\text{Per cent larval mortality} = \frac{\text{Number of larvae dead due to infection}}{\text{Total number of larvae treated}} \times 100$$

3. RESULTS AND DISCUSSION

A total of twelve types of oil cultures were prepared by using four oils and these cultures were maintained at three different temperatures. Additionally, an unformulated crude suspension of *Beauveria bassiana* and untreated control were maintained. The post treatment larval changes and the results of viability and virulence were presented below.

After releasing the larvae onto treated leaves, it was observed that there was a noticeable reduction in their tendency to feed, particularly on leaves treated with sesame oil. When the larvae were fed with fresh leaves the day after treatment, they exhibited a reduced inclination to feed. The room temperature ($28 \pm 3^\circ\text{C}$) was highly conducive for mycosis development. Freshly moulted 3rd instar larvae were significantly more susceptible to EPF treatments compared to late 3rd and 4th instar *S. litura* larvae. After 3-4 days of treatment with *B. bassiana*, the infected larvae exhibited sluggish movement and consumed less food material. The larval integument got shrunken and larval bodies had become exceedingly smooth. Majority of larvae were dead within 1 week of inoculation with fungal spores. There was sparse growth of fungal mycelium on the surface of larval integument and whitish mycelial growth of fungus was conspicuous from 4th day after inoculation. The entire larval body was covered with puffy whitish mycelia and whitish fungal spores were produced on to larval surface. As the time progressed, the puffiness of the fungus on the larval body reduced gradually. The larval body eventually turned into a hard and mummified cadaver. The stiffness of the cadaver after death might be due to excessive fungal growth inside larval body. Larval mortality occurred more rapidly within 5-6 days with oil formulations, compared to 8-10 days with unformulated crude suspensions. The results were in agreement with Ummidi and Vadlamani (2014) who conducted bioassay studies of oil formulations of *B. bassiana* against *S. litura*. Based on compatibility

studies, almond oil, olive oil, gingelly oil, castor oil based formulations were chosen for bioassay. He reported that all the four formulations displayed higher mortalities of the target pest compared to unformulated conidia, with *B. bassiana* showed higher mortality with gingelly oil and almond oil.

In the present study it was observed that, the infected larvae progressed to next instar and pupated earlier compared to untreated larvae *i.e.* significant decrease in larval period was observed due to infection in comparison with untreated control. Some affected larvae metamorphosed to malformed pupae and adult. These results corroborate with the findings of Torrado-Leon *et al.* (2006) who documented the interference in the moulting process of *Bemisia tabaci* (Gennadius) nymphs when treated with *B. bassiana*. More than 30% of the imago emerged from treated nymphs were unable to detach completely from the exuvium.

It was noticed that on the day of preparation, the highest mean per cent conidial viability of 87.85% was recorded in Sesamum oil cultures (T₁₀, T₁₁, T₁₂), followed by Liquid paraffin oil cultures (T₄, T₅, T₆) and Rice bran oil cultures (T₇, T₈, T₉) which recorded 83.57% and 80.71% respectively (Fig. 3). Comparatively lower conidial viability of 77.14% was observed in oil cultures formulated using Mineral oil (T₁, T₂, T₃). The unformulated culture (T₁₃) recorded a conidial viability of 63.57% (Table. 1). The data regarding the mean percent larval mortality indicated that the maximum larval mortality of 90.00% was recorded in Sesamum oil cultures (T₁₀, T₁₁, T₁₂), followed by Liquid paraffin oil cultures (T₄, T₅, T₆) and Rice bran oil cultures (T₇, T₈, T₉) which recorded 86.67% and 83.33% respectively (Fig. 4). Comparatively lower larval mortality of 80.00% was observed in oil cultures formulated using Mineral oil (T₁, T₂, T₃). The unformulated culture (T₁₃) recorded a larval mortality of 63.33%. There was no larval mortality recorded in untreated control T₁₄. (Table. 2)

In the investigations after thirty days of storage, the highest mean per cent conidial viability of 82.14% was recorded with Sesamum oil cultures stored at -20°C (T₁₂). The treatment T₁₁ (80.71%) was on par with T₆ (79.28%). The treatments T₉ (76.42%), T₈ (75.71%) and T₃ & T₅ (74.28%) were on par with each other. These were followed by the treatments T₂ and T₄ & T₁₀ (70.71%). Comparatively lower conidial viability of 67.85% was observed in oil cultures formulated using Mineral oil (T₁) and Rice Bran oil at room temperature (T₇). The unformulated culture (T₁₃) recorded a conidial viability of 61.43% (Table. 1). The observations of larval mortalities of *Beauveria bassiana* oil cultures after being stored for 30 days revealed that the highest per cent larval mortality of 86.67% was observed in T₁₂ followed by T₆ (83.33%) and T₉ (80.00%). The next superior treatments T₃, T₈ & T₁₁ (76.67%) and T₅ (73.33%) were on par with each other. These were followed by the treatments T₂, T₄ & T₁₀ (70.00%) and T₁ & T₇ (66.67%). Comparatively lower mortality of 60.00% was observed in T₁₃. The per cent larval mortality in untreated control was 0.00% (Table. 2).

Among the treatments after sixty days of storage, the treatment (T₁₂) Sesamum oil cultures stored at -20°C was recorded with the highest mean per cent conidial viability of 79.28%. T₁₁ (76.42%) was on par with T₉ (75.71%). The next better treatment was T₆ (74.28%). The treatments T₃ & T₅ (70.71%), T₈ (69.28%) and T₄ (67.85%) were on par with each other. These were followed by the

treatments T₂&T₁₀ (65.71%), T₇ (64.28%) and T₁ (61.43%) were on par with each other. The unformulated culture (T₁₃) recorded a conidial viability of 58.57% (Table. 1). After 60 days of storage, highest mean per cent larval mortality of 83.33% was observed in T₁₂ followed by T₁₁ (80.00%), T₆& T₉ (76.67), T₃, T₅ & T₈ (70.00%) and T₂, T₄& T₁₀ (66.67%). The treatments T₇ (63.33%) and T₁ (60.00%) were on par with each other. Comparatively lower mortality of 56.67% was observed in T₁₃. There was no larval mortality recorded in untreated control T₁₄ (Table. 2).

It was observed that among all the treatments after ninety days of storage, the treatment T₁₂ was recorded with the highest mean per cent conidial viability of 75.71% and it was on par with T₁₁ (74.28%) and T₉ (73.57%). The next superior treatment was T₆ (70.71%). The treatments T₃& T₅ (65.00%) was on par with T₈ (65.71%). T₁₀ (63.57%), T₂ (62.87%) and T₄ (61.43%) were on par with each other. These were followed by the treatments, T₇& T₁ (57.87%). The unformulated culture (T₁₃) recorded a conidial viability of 51.43% (Table. 1). The results of mean per cent larval mortality after ninety days of storage of oil cultures revealed that the highest per cent larval mortality was recorded in T₁₂ (76.67%), followed by T₉& T₁₁ (73.33%). The treatments T₆ (70.00%) and T₃ & T₅ (66.67%) were on par with each other. These treatments were followed by T₈ & T₁₀ (63.33%) and T₂& T₄ (60.00%) which were on par with each other. Comparatively lower per cent larval mortalities were observed in T₁& T₇ (56.67%) and T₁₃ (50.00%). The per cent larval mortality in untreated control was 0.00% (Table. 2)

The data on observation of mean per cent conidial viability after 120 DAS among all the treatments, T₁₂ was recorded with the highest mean per cent conidial viability of 74.28%. T₁₁ (67.85%) was on par with T₉ (69.28%). The next superior treatment T₆ (65.71%), T₃ (64.28%) and T₅ (61.43%) were on par with each other. The treatments T₂ & T₈ (58.57%), T₁₀ (57.87%) and T₄& T₇ (54.29%) were statistically indifferent from each other. These were followed by the treatment T₁ (51.43%). The unformulated culture (T₁₃) recorded a conidial viability of 45.71% (Table. 1). The observations after 120 days of storage of *B.bassiana* oil cultures revealed that maximum of 73.33% mean per cent larval mortality was recorded in T₁₂ followed by T₉ (70.00%) and T₆ & T₁₁ (66.67%). T₃ (63.33%) was on par with T₅ (60.00%). These treatments were followed by T₂, T₈ and T₁₀ (56.67%), T₄& T₇ (53.33%) followed by T₁ (50.00%) and T₁₃ (43.33%). There was no mortality recorded in untreated control T₁₄ (Table. 2).

Table 1: Viability of *Beauveria bassiana* conidia stored under different temperature regimes after formulating as oil cultures (2023-2024)

S. No.	Treatment	Mean Per cent Conidia germination						
		0 DAS	30 DAS	60 DAS	90 DAS	120 DAS	150 DAS	180 DAS
1	T ₁	77.14 ^c (61.44)	67.85 ^{cd} (55.46)	61.43 ^{de} (51.61)	57.87 ^d (49.53)	51.43 ^{cd} (45.82)	44.29 ^{de} (41.72)	41.43 ^f (40.07)
2	T ₂	77.14 ^c (61.44)	70.71 ^{bcd} (57.23)	65.71 ^{cde} (54.16)	62.87 ^{cd} (52.46)	58.57 ^{bcd} (49.93)	51.43 ^{bcdde} (45.82)	48.57 ^{def} (44.18)
3	T ₃	77.14 ^c (61.44)	74.28 ^{abc} (59.53)	70.71 ^{bcd} (57.23)	65.00 ^{bc} (53.73)	64.28 ^{abc} (53.30)	58.57 ^{abcd} (49.93)	54.29 ^{bcd} (47.46)
4	T ₄	83.57 ^b (66.09)	70.71 ^{bcd} (57.23)	67.85 ^{bcd} (55.46)	61.43 ^{cd} (51.61)	54.29 ^{bcd} (47.46)	49.29 ^{bcdde} (44.59)	45.71 ^{def} (42.54)

5	T ₅	83.57 ^b (66.09)	74.28 ^{abc} (59.53)	70.71 ^{bcd} (57.23)	65.00 ^{bc} (53.73)	61.43 ^{abc} (51.61)	58.57 ^{abcd} (49.93)	51.43 ^{cde} (45.82)
6	T ₆	83.57 ^b (66.09)	79.28 ^{ab} (62.92)	74.28 ^{abc} (59.53)	70.71 ^{ab} (57.23)	65.71 ^{abc} (54.16)	63.75 ^{abc} (52.98)	61.43 ^{ab} (51.61)
7	T ₇	80.71 ^{bc} (63.95)	67.85 ^{cd} (55.46)	64.28 ^{de} (53.30)	57.87 ^d (49.53)	54.29 ^{bcd} (47.46)	48.57 ^{cde} (44.18)	44.29 ^{ef} (41.72)
8	T ₈	80.71 ^{bc} (63.95)	75.71 ^{abc} (60.47)	69.28 ^{bcd} (56.34)	65.71 ^{bc} (54.16)	58.57 ^{bcd} (49.93)	55.71 ^{abcd} (48.28)	51.43 ^{cde} (45.82)
9	T ₉	80.71 ^{bc} (63.95)	76.42 ^{abc} (60.95)	75.71 ^{ab} (60.47)	73.57 ^a (59.06)	69.28 ^{ab} (56.34)	64.28 ^{ab} (53.30)	58.57 ^{abc} (49.93)
10	T ₁₀	87.85 ^a (69.60)	70.71 ^{bcd} (57.23)	65.17 ^{de} (53.83)	63.57 ^{cd} (52.87)	57.87 ^{bcd} (49.53)	49.29 ^{bcd} (44.59)	47.85 ^{def} (43.77)
11	T ₁₁	87.85 ^a (69.60)	80.71 ^{ab} (63.95)	76.42 ^{ab} (60.95)	74.28 ^a (59.53)	67.85 ^{ab} (55.46)	62.87 ^{abc} (52.46)	54.29 ^{bcd} (47.46)
12	T ₁₂	87.85 ^a (69.60)	82.14 ^a (65.00)	79.28 ^a (62.92)	75.71 ^a (60.47)	74.28 ^a (59.53)	67.85 ^a (55.46)	62.87 ^a (52.46)
13	T ₁₃	63.57 ^d (52.87)	61.43 ^d (51.61)	58.57 ^e (49.93)	51.43 ^e (45.82)	45.71 ^d (42.54)	39.29 ^e (38.82)	32.85 ^g (34.97)
14	T ₁₄	0.00 ^e (0.00)	0.00 ^e (0.00)	0.00 ^e (0.00)	0.00 ^e (0.00)	0.00 ^e (0.00)	0.00 ^e (0.00)	0.00 ^e (0.00)
	General mean	75.10	68.01	64.24	60.36	55.97	50.98	46.79
	F	Sig.	Sig.	Sig.	Sig.	Sig.	Sig.	Sig.
	Sig.	0.000	0.000	0.000	0.000	0.000	0.000	0.000
	SE(m)±	0.72	0.37	0.61	0.59	0.57	0.56	0.56
	C.D.	2.10	1.07	1.78	1.72	1.67	1.64	1.63
	C.V.	2.09	1.16	2.02	2.05	2.10	2.20	2.31

DAS: Days After Storage, Values are the means of three replications,

Values in parentheses are angular transformed values

Means in a column followed by same superscript are not significantly different according to DMRT at $P \leq 0.05$.

SE(m)± = Standard error of mean **C.V.** = Coefficient of variation

C.D. = Critical Difference at 1% level of significance

T₁: Mineral oil culture of *B. bassiana* stored at Room temperature ($28 \pm 3^\circ\text{C}$), T₂: Mineral oil culture of *B. bassiana* stored at Refrigerator temperature (4°C), T₃: Mineral oil culture of *B. bassiana* stored at freezing temperature (-20°C), T₄: Liquid paraffin oil culture of *B. bassiana* stored at Room temperature ($28 \pm 3^\circ\text{C}$), T₅: Liquid paraffin oil culture of *B. bassiana* stored at Refrigerator temperature (4°C), T₆: Liquid paraffin oil culture of *B. bassiana* stored at freezing temperature (-20°C), T₇: Rice bran oil culture of *B. bassiana* stored at Room temperature ($28 \pm 3^\circ\text{C}$), T₈: Rice bran oil culture of *B. bassiana* stored at Refrigerator temperature (4°C), T₉: Rice bran oil culture of *B. bassiana* stored at freezing temperature (-20°C), T₁₀: Sesamum oil culture of *B. bassiana* stored at Room temperature ($28 \pm 3^\circ\text{C}$), T₁₁: Sesamum oil culture of *B. bassiana* stored at Refrigerator temperature (4°C), T₁₂: Sesamum oil culture of *B. bassiana* stored at freezing temperature (-20°C), T₁₃: Unformulated culture of *B. bassiana* (Not preserved in oil), T₁₄: Untreated control.

Table 2: Virulence of *Beauveria bassiana* conidia stored under different temperature regimes after formulating as oil cultures against third instar larva of *Spodoptera litura* under laboratory conditions (2023-2024)

S. No	Treatment	Mean per cent larval Mortality of third instar <i>Spodoptera litura</i> larvae						
		0 DAS	30 DAS	60 DAS	90 DAS	120 DAS	150 DAS	180 DAS
1	T ₁	80.00 ^a (63.44)	66.67 ^{de} (54.74)	60.00 ^{de} (50.77)	56.67 ^{cd} (48.33)	50.00 ^{de} (45.00)	43.33 ^{cd} (41.17)	40.00 ^{cd} (39.23)
2	T ₂	80.00 ^a (63.44)	70.00 ^{cde} (56.79)	66.67 ^{cde} (54.74)	60.00 ^{bcd} (50.77)	56.67 ^{bcd} (48.83)	50.00 ^{abcd} (45.00)	46.67 ^{abc} (43.09)
3	T ₃	80.00 ^a	76.67 ^{bcd}	70.00 ^{bcd}	66.67 ^{abc}	63.33 ^{abcd}	56.67 ^{abc}	53.33 ^{abc}

		(63.44)	(61.12)	(56.79)	(54.74)	(52.73)	(48.83)	(46.91)
4	T ₄	86.67 ^a (68.59)	70.00 ^{cde} (56.79)	66.67 ^{cde} (54.74)	60.00 ^{bcd} (50.77)	53.33 ^{cde} (46.91)	50.00 ^{abcd} (45.00)	43.33 ^{bcd} (41.17)
5	T ₅	86.67 ^a (68.59)	73.33 ^{bcd} (58.91)	70.00 ^{bcd} (56.79)	66.67 ^{abc} (54.74)	60.00 ^{abcd} (50.77)	56.67 ^{abc} (48.83)	50.00 ^{abc} (45.00)
6	T ₆	86.67 ^a (68.59)	83.33 ^{ab} (65.90)	76.67 ^{abc} (61.12)	70.00 ^{abc} (56.79)	66.67 ^{abc} (54.74)	63.33 ^{ab} (52.73)	60.00 ^a (50.77)
7	T ₇	83.33 ^a (65.90)	66.67 ^{de} (54.74)	63.33 ^{de} (52.73)	56.67 ^{cd} (48.83)	53.33 ^{cde} (46.91)	46.67 ^{bcd} (43.09)	43.33 ^{bcd} (41.17)
8	T ₈	83.33 ^a (65.90)	76.67 ^{bcd} (61.12)	70.00 ^{bcd} (56.79)	63.33 ^{bcd} (52.73)	56.67 ^{bcd} (48.83)	53.33 ^{abcd} (46.91)	50.00 ^{abc} (45.00)
9	T ₉	83.33 ^a (65.90)	80.00 ^{abc} (63.44)	76.67 ^{abc} (61.12)	73.33 ^{ab} (58.91)	70.00 ^{ab} (56.79)	63.33 ^{ab} (52.73)	56.67 ^{ab} (48.83)
10	T ₁₀	90.00 ^a (71.57)	70.00 ^{cde} (56.79)	66.67 ^{cde} (54.74)	63.33 ^{bcd} (52.73)	56.67 ^{bcd} (48.83)	50.00 ^{abcd} (45.00)	46.67 ^{abc} (43.09)
11	T ₁₁	90.00 ^a (71.57)	76.67 ^{bcd} (61.12)	80.00 ^{ab} (63.44)	73.33 ^{ab} (58.91)	66.67 ^{abc} (54.74)	60.00 ^{abc} (50.77)	53.33 ^{abc} (46.91)
12	T ₁₂	90.00 ^a (71.57)	86.67 ^a (68.59)	83.33 ^a (65.90)	76.67 ^a (61.12)	73.33 ^a (58.91)	66.67 ^a (54.74)	60.00 ^a (50.77)
13	T ₁₃	63.33 ^d (52.73)	60.00 ^e (50.77)	56.67 ^e (48.83)	50.00 ^d (45.00)	43.33 ^e (41.17)	36.63 ^d (37.25)	30.00 ^d (33.21)
14	T ₁₄	0.00 ^c (0.00)	0.00 ^f (0.00)	0.00 ^f (0.00)	0.00 ^e (0.00)	0.00 ^f (0.00)	0.00 ^e (0.00)	0.00 ^e (0.00)
	General mean	77.38	68.33	64.76	59.76	55.00	49.76	45.24
	F	Sig.	Sig.	Sig.	Sig.	Sig.	Sig.	Sig.
	Sig.	0.000	0.000	0.000	0.000	0.000	0.000	0.000
	SE(m)±	1.85	1.74	1.62	1.65	1.52	1.43	1.36
	C.D.	5.40	5.06	4.73	4.81	4.43	4.16	3.96
	C.V.	5.21	5.47	5.33	5.76	5.63	5.65	5.73

DAS: Days After Storage, Values are the means of three replications,

Values in parentheses are angular transformed values

Means in a column followed by same superscript are not significantly different according to DMRT at $P \leq 0.05$.

SE(m)± = Standard error of mean **C.V.** = Coefficient of variation **C.D.** = Critical Difference at 1% level of significance

T₁: Mineral oil culture of *B. bassiana* stored at Room temperature ($28 \pm 3^\circ\text{C}$), T₂: Mineral oil culture of *B. bassiana* stored at Refrigerator temperature (4°C), T₃: Mineral oil culture of *B. bassiana* stored at freezing temperature (-20°C), T₄: Liquid paraffin oil culture of *B. bassiana* stored at Room temperature ($28 \pm 3^\circ\text{C}$), T₅: Liquid paraffin oil culture of *B. bassiana* stored at Refrigerator temperature (4°C), T₆: Liquid paraffin oil culture of *B. bassiana* stored at freezing temperature (-20°C), T₇: Rice bran oil culture of *B. bassiana* stored at Room temperature ($28 \pm 3^\circ\text{C}$), T₈: Rice bran oil culture of *B. bassiana* stored at Refrigerator temperature (4°C), T₉: Rice bran oil culture of *B. bassiana* stored at freezing temperature (-20°C), T₁₀: Sesamum oil culture of *B. bassiana* stored at Room temperature ($28 \pm 3^\circ\text{C}$), T₁₁: Sesamum oil culture of *B. bassiana* stored at Refrigerator temperature (4°C), T₁₂: Sesamum oil culture of *B. bassiana* stored at freezing temperature (-20°C), T₁₃: Unformulated culture of *B. bassiana* (Not preserved in oil), T₁₄: Untreated control.

Fig.3. Viability of *Beauveria bassiana* conidia stored under different temperature regimes after formulating as oil cultures (2023-2024)





Fig.4. Virulence of *Beauveria bassiana* conidia stored under different temperature regimes after formulating as oil cultures against third instar larva of *Spodoptera litura* under laboratory conditions (2023-2024)



The results after 150 days of storage of different oil cultures of *Beauveria bassiana* revealed that among the treatments T₁₂ was recorded with the highest mean per cent conidial viability of 67.85%, followed by T₉ (64.28%). T₆ (63.75%) was on par with T₁₁ (62.87%). The treatment T₃ & T₅ (58.57%) on par with T₈ (55.71%). The treatments T₂ (51.43%) and T₁₀ & T₄ (49.29%) were statistically indifferent. These were followed by the treatments T₇ (48.57%) and T₁ (44.29%). The unformulated culture (T₁₃) recorded a conidial viability of 39.29% (Table. 1). Storage of oil cultures for 150 days in the laboratory showed that highest mean per cent larval mortality of 66.67% was recorded in T₁₂ followed by T₆ & T₉ (63.33%). T₁₁ (60.00%) and T₃ & T₅ (56.67%) were on par with each other. The treatments T₈ (53.33%) and T₂, T₄ & T₁₀ (50.00%) were significantly indifferent with each other. These were followed by T₇ (46.67%), T₁ (43.33%) and T₁₃ (36.63%). The per cent larval mortality in untreated control was 0.00%. (Table. 2).

The results of the percent viability after 180 days of storage clearly shows that, the highest mean per cent conidial viability of 62.87% was recorded with T₁₂ followed by T₆ (61.43%), T₉ (58.57%), T₃ & T₁₁ (54.29%) and T₅ & T₈ (51.43%). The treatments T₂ (48.57%), T₁₀ (47.85%) and T₄ (45.71%) were on par with each other. These were followed by the treatments T₇ (44.29%) and T₁ (41.43%). The unformulated culture (T₁₃) recorded a conidial viability of 32.85% (Table. 1). The data after 180 days of storage of *Beauveria bassiana* oil cultures clearly revealed that highest mean larval mortality of 60.00% was recorded with T₁₂ & T₆ followed by T₉ (56.67%). The treatments T₃ & T₁₁ (53.33%), T₅ & T₈ (50.00%) and T₂ & T₁₀ (46.67%) were on par with each other. These were followed by the treatments T₄ & T₇ (43.33%) followed by T₁ (40.00%) and T₁₃ (30.00%). There was no larval mortality recorded in untreated control T₁₄ (Table. 2).

Comparatively higher mean per cent conidial viabilities of *Beauveria bassiana* oil cultures were observed in the treatments T₁₂, T₆, T₉ and T₁₁, from the day of preparation to 180 days after storage. The viability varied as follows: T₁₂ (87.85% to 62.87%), T₆ (83.57 to 61.43%), T₉ (80.71 to 58.57%) and T₁₁ (87.85 to 54.29%). Comparatively lower per cent conidial viability were recorded in T₁ (77.41 to 41.43%) and T₁₃ (63.57 to 32.85%). The highest mean per cent larval mortalities were observed in Treatments T₁₂, T₆, T₉ and T₁₁ when treated against third instar *S. litura* larvae at monthly intervals. The mean per cent larval mortalities ranged as follows: T₁₂ (90.00% to 60.00%), T₆ (86.67 to 60.00%), T₉ (83.33 to 56.67%) and T₁₁ (90.00 to 53.33%). Comparatively lower per cent larval mortalities were recorded in T₁ (80.00 to 40.00%) and T₁₃ (63.33 to 30.00%).

The pathogenicity levels correlated with conidial viabilities, showing a gradual decrease in both viability and virulence over the 180-day storage period.

From the present study, it can be concluded that the performance of *Beauveria bassiana* was best with oil formulations compared to unformulated crude cultures. Among the oils used, Sesamum oil formulations were the most effective followed by Liquid paraffin oil, Rice bran oil and mineral oil.

Considering the storage conditions, the formulations stored at freezing temperature (-20°C) performed better to those stored at Refrigerated temperature (4°C) and Room temperature (28 ± 3°C).

Sesamum oil cultures had been found out to be best among all oil cultures, this may be due to Sesquiterpene present in it which have insecticidal properties that serve as a synergist for pyrethroid insecticides. (Wei *et al.*, 2022, Baker *et al.*, 2018 and Sirato-Yasumoto *et al.*, 2001).

The performance of oil cultures were best at lower temperatures compared to room temperature, which may be due to reduced metabolism and growth at lower temperatures thereby maintaining viability for an extended period for extended period in such lower temperatures. Murugasridevi *et al.* (2021) prepared oil-based formulation by dissolving 1 g of pure conidia (10^{10} conidia g⁻¹) of *B. bassiana* (Bb 112) in 100 ml of paraffin oil, along with adjuvants like polyethylene glycol and tween 80 to enhance the efficacy of the formulation and stored it under different storage conditions viz. ambient temperature (28 ± 2°C) and refrigerated condition (4°C). Shelf-life assessment showed that the formulation stored under ambient temperature and refrigerated condition recorded 587.00 and 590.77 × 10⁸ CFU ml⁻¹, respectively on the day of preparation and 167.66 and 216.66 × 10⁸ CFU ml⁻¹, respectively after 24 weeks.

The results were in agreement with Lakshmi *et al.* (2020) prepared 13 types of plant oil-based bio-formulation of *Beauveria bassiana* (TBb8 strain) using corn, soyabean, castor, rice bran, mustard, mahua, pinnai, neem, groundnut, palm, coconut and gingelly oils with different combinations and tested its efficacy against the lepidopteran fruit borer *Helicoverpa armigera* (Hubner). Among 13 oils, corn oil was found superior by maintaining a population of 7.0 × 10⁹ cfu/ml after 210 days of storage. Oil formulation of TBb8 isolate (1 × 10⁸ spore/ml) treated *H. armigera* showed larval mortality of 70.0%, pupation period of 13.7 days, pupal malformation of 17.9% and adult malformation was recorded as 16.2%.

Sandhu *et al.* (1993) evaluated the effects of temperature on conidial viability and virulence of *B. bassiana* against third instar *Helicoverpa armigera* (Hubner) larvae over a 24-month period. He stored unformulated conidia of *B. bassiana* at five different temperatures (0°, 10°, 20°, 30° and 40°C). Based on the experimental results they reported that conidia survived longest at lower temperatures (0–20°C). At higher temperatures, (30–40°C) conidia did not survive.

Oliveira *et al.* (2011) evaluated the effect of preservation of *B. bassiana* isolates for one year on growth, production and viability of spores as well as macro- and micro-morphology by using 3 storage methods viz. freezing at -20°C in an aqueous glycerol solution; freeze-drying (lyophilized) and maintaining at ambient temperature with continual sub-culturing in PDA medium. They reported that glycerol-freeze method at -20°C is the most suitable method for long-term preservation of fungal species.

4. CONCLUSIONS

The results indicated that *B. bassiana* could be a potent biocontrol agent for managing lepidopteran pests. The combinations of oils and controlled storage temperature presents a promising strategy in enhancing the effectiveness of *B. bassiana* compared to unformulated crude culture. This approach not only enhances the practical application but also supports sustainable pest management practices by providing a reliable and environmentally friendly alternative to chemical pesticides. Further research should be conducted in this line at field level to identify the most effective oil type and concentrations suiting different environment conditions and pest targets, as well as optimizing the application methods to ensure uniform coverage and maximum reach to pests. Additionally, research on compatibility between entomopathogenic fungal oil cultures and other biocontrol agents or agricultural practices will improve its effectiveness and reliability.

REFERENCES

1. FAO, 2022. Pesticides Use, Pesticides Trade and Pesticides Indicators-Global, Regional and Country Trends, 1990–2020. *FAOSTAT Analytical Briefs*, no. 46.
2. Dong, H., Zhu, K.H., Zhao, Q., Bai, X.P., Zhou, J.C. and Zhang, L.S. Morphological defense of the egg mass of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) affects parasitic capacity and alters behaviours of egg parasitoid wasps. *Journal of Asia-Pacific Entomology*.2021;24 (3): 671-678.
3. Waqas, M.S., Shi, Z., Yi, T.C., Xiao, R., Shoaib, A.A., Elabasy, A.S. and Jin, D.C. Biology, ecology, and management of cotton mealybug *Phenacoccusolenopsis* Tinsley (Hemiptera: Pseudococcidae). *Pest Management Science*.2021; 77 (12): 5321-5333.
4. Guo, J., Wu, S., Zhang, F., Huang, C., He, K., Babendreier, D. and Wang, Z. Prospects for microbial control of the fall armyworm *Spodoptera frugiperda*: a review. *BioControl*.2020; 65 :647-662.
5. Yan-li, Z., Hui, D., Li-sheng, Z., Zu-min, G. and Jin-cheng, Z. 2022. High virulence of a naturally occurring entomopathogenic fungal isolate, *Metarhizium (Nomuraea) rileyi*, against *Spodoptera frugiperda*. *Journal of Applied Entomology*:2022; 146 (6): 659-665.
6. Ummidi, V.R.S. and Vadlamani, P. Preparation and use of oil formulations of *Beauveria bassiana* and *Metarhizium anisopliae* against *Spodoptera litura* larvae. *African Journal of Microbiology Research*.2014; 8 (15): 1638-1644.
7. Sharma, A., Sharma, S. and Yadav, P.K. Entomopathogenic fungi and their relevance in sustainable agriculture: A review. *Cogent Food & Agriculture*.2023; 9 (1)
8. Mantzoukas, S., Kitsiou, F., Natsiopoulou, D and Eliopoulos, P.A. Entomopathogenic fungi: interactions and applications. *Encyclopedia*. 2022; 2 (2): 646-656.
9. Bateman, R.P., Douro-Kpindou, O.K., Kooyman, C., Lomer, C. and Ouambama, Z. Some observations on the dose transfer of mycoinsecticide sprays to desert locusts. *Crop Protection*. 1998;17(2): 151-158.
10. Ramanujam, B., Rangeshwaran, R., Sivakmar, G., Mohan, M. and Yandigeri, M.S. Management of insect pests by microorganisms. *Proceedings of the Indian National Science Academy*. 2014; 80 (2): 455-471.
11. Gebremariam, A., Chekol, Y. and Assefa, F.Extracellular enzyme activity of entomopathogenic fungi, *Beauveria bassiana* and *Metarhizium anisopliae* and their pathogenicity potential as a bio-control agent against whitefly pests, *Bemisia tabaci* and *Trialeurodes vaporariorum* (Hemiptera: Aleyrodidae). *BMC Research Notes*.2022; 15 (1): 117.
12. Seema, Y., Neeraj, T. and Krishan, K. Mass production of entomopathogens *Beauveria bassiana* and *Metarhizium anisopliae* using rice as a substrate by diphasic liquid-solid fermentation technique. *International Journal of Advanced Biological Research*. 2013; 3 (3): 331-335.
13. Beys-da-Silva, W.O., Santi, L., Berger, M., Calzolari, D., Passos, D.O., Guimarães, J.A., Moresco, J.J. and Yates, J.R. Secretome of the biocontrol agent *Metarhizium anisopliae* induced by the cuticle of the cotton pest *Dysdercus peruvianus* reveals new insights into infection. *Journal of Proteome Research*. 2014; 13 (5):2282-2296.
14. Dusfour, I., Vontas, J., David, J.P., Weetman, D., Fonseca, D.M., Corbel, V., Raghavendra, K., Coulibaly, M.B., Martins, A.J., Kasai, S. and Chandre, F. Management of insecticide resistance in

- the major *Aedes* vectors of arboviruses: Advances and challenges. *PLoS neglected tropical diseases*.2019; 13 (10): e0007615.
15. de Faria, M.R. and Wraight, S.P. Mycoinsecticides and mycoacaricides: a comprehensive list with worldwide coverage and international classification of formulation types. *Biological control*. 2007; 43 (3): 237-256.
 16. Islam, W., Adnan, M., Shabbir, A., Naveed, H., Abubakar, Y.S., Qasim, M., Tayyab, M., Noman, A., Nisar, M.S., Khan, K.A. and Ali, H. Insect-fungal-interactions: A detailed review on entomopathogenic fungi pathogenicity to combat insect pests. *Microbial Pathogenesis*. 2021; 159: 105122.
 17. Money, N.P. Fungi and biotechnology. *The fungi*.2016; 401-424. Academic Press, London.
 18. Singh, H.B., Keswani, C., Ray, S., Yadav, S.K., Singh, S.P., Singh, S and Sarma, B.K. *Beauveria bassiana*: biocontrol beyond lepidopteran pests. *Biocontrol of Lepidopteran Pests: Use of Soil Microbes and their Metabolites*.2015;219-235.
 19. Wang, Q. and Xu, L. Beauvericin, a bioactive compound produced by fungi: a short review. *Molecules*. 2012. 17 (3): 2367-2377.
 20. Rachappa, V., Lingppa, S., Patil, R.K. and Hugar, P.S. Suitability of media and containers for mass production of *Metarhizium anisopliae*. *Karnataka Journal of Agricultural Sciences*. 2005;18 (3): 680.
 21. Inyang, E.N., McCartney, H.A., Oyejola, B., Ibrahim, L. and Archer, S.A. Effect of formulation, application and rain on the persistence of the entomogenous fungus *Metarhizium anisopliae* on oilseed rape. *Mycological Research*.2000; 104 (6): 653-661.
 22. Devi, P.V. and Prasad, Y.G. Compatibility of oils and antifeedants of plant origin with the entomopathogenic fungus *Nomuraearileyi*. *Journal of invertebrate pathology*. 1996; 68 (1): 91-93.
 23. Bateman, R.P., Carey, M., Moore, D.E. and Prior, C. The enhanced infectivity of *Metarhizium flavoviride* in oil formulations to desert locusts at low humidities. *Annals of Applied Biology*. 1993;122 (1): 145-152.
 24. Batta, Y.A. Production and testing of novel formulations of the entomopathogenic fungus *Metarhizium anisopliae* (Metschnikoff) Sorokin (Deuteromycotina: Hyphomycetes). *Crop Protection*. 2003;22 (2): 415-422.
 25. Moore, D., Bridge, P.D., Higgins, P.M., Bateman, R.P. and Prior, C. Ultra-violet radiation damage to *Metarhizium flavoviride* conidia and the protection given by vegetable and mineral oils and chemical sunscreens. *Annals of Applied Biology*. 1993; 122 (3): 605-616.
 26. Humber, R.A. Fungi: preservation of cultures. In *Manual of techniques in insect pathology*. Academic Press.1997; 269-279.
 27. López Lastra, C.C., Hajek, A.E. and Humber, R.A. Comparing methods of preservation for cultures of entomopathogenic fungi. *Canadian Journal of Botany*. 2002; 80 (10): 1126-1130.
 28. Saheb, Y.P., Manjula, K.K., Devaki, Devi, R.S.J., Reddy, B.R. and Anokhe, A. Evaluation of *Metarhizium (Nomuraea) rileyi* Rice Bran Oil Formulations against 3rd instar *Spodoptera litura* under Laboratory. *Biological Forum – An International Journal*.2021; 13 (3): 601-607.
 29. Torrado-León, E., Montoya-Lerma, J. and Valencia-Pizo, E. Sublethal effects of *Beauveria bassiana* (Balsamo) Vuillemin (Deuteromycotina: Hyphomycetes) on the whitefly *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae) under laboratory conditions. *Mycopathologia*. 2006;162: 411-419.
 30. Wei, P., Zhao, F., Wang, Z., Wang, Q., Chai, X., Hou, G. and Meng, Q. Sesame (*Sesamum indicum* L.): A comprehensive review of nutritional value, phytochemical composition, health benefits, development of food, and industrial applications. *Nutrients*. 2022;14(19): 4079.
 31. Baker, B.P. and Grant, J.A. Active ingredients eligible for minimum risk pesticide use: Overview of the profiles.2018
 32. Sirato-Yasumoto, S., Katsuta, M., Okuyama, Y., Takahashi, Y. and Ide, T. Effect of sesame seeds rich in sesamin and sesamol on fatty acid oxidation in rat liver. *Journal of Agricultural and Food Chemistry*. 2001. 49(5): 2647-2651.
 33. Murugasridevi, K., Jeyarani, S. and Kumar, S.M. Evaluation of shelf life and pathogenicity of oil based formulation of *Beauveria bassiana* against chilli thrips and mites. *Journal of Entomological Research*. 2021. 45 (4): 679-685.
 34. Lakshmidevi, P., Gopalakrishnan, C. and Antony, R.S. Bio-formulation based on plant oil of the future *Beauveria bassiana* (Balsamo) for fruit borer control in tomatoes. *Journal of Entomology and Zoology Studies*. 2020; 8: 200-206.
 35. Sandhu, S.S., Rajak, R.C. and Agarwal, G.P. Studies on prolonged storage of *Beauveria bassiana* conidia: effects of temperature and relative humidity on conidial viability and virulence

- against chickpea borer, *Helicoverpa armigera*. *Biocontrol Science and Technology*. 1993; 3 (1): 47-53.
36. Oliveira, I., Pereira, J.A., Bento, A. and Baptista, P. Viability of *Beauveria bassiana* isolates after storage under several preservation methods. *Annals of Microbiology*. 2011; 61 (2): 339-344.

UNDER PEER REVIEW