

# Structural and Potential Biological Properties of SBDTC and SMDTC Containing Metal Complexes: A Brief Review

## Abstract

S-benzylthiocarbamate (SBDTC) and S-methylthiocarbamate (SMDTC) are working as a starting material for the synthesis of various Schiff base ligands. SBDTC and SMDTC can act as NS, SNNS, SNNNS, and NOS chelating agents and have biological and chemical potential for synthesizing many transition and inner transition metal complexes. The metal complexes prepared from SBDTC, SMDTC, and their derivatives have shown promising biological activities like antibacterial, cytotoxicity, antioxidant, antitumor, and anticancer activity. Last few decades, a lot of works have been reported on SBDTC and SMDTC containing Schiff base metal complexes. In this review, it is considered to show the various synthesis procedures, and structural and biological activities of SBDTC and SMDTC-containing metal complexes.

Keywords: Schiff base, metal complexes, SBDTC, SMDTC, antibacterial activity, antioxidant properties, anticancer activity.

## 1. Introduction

Nitrogen-oxygen, Nitrogen-Sulfur, Sulfur-Nitrogen-Nitrogen-Sulfur, and Nitrogen-Oxygen-Sulfur chelating agents have drawn prime interest to the researchers for the synthesis of new metal complexes that may provide an important path for the development of chemotherapeutic agents and to improve the field of bioinorganic chemistry [1,2]. It was also reported that S-methylthiocarbamate (SMDTC), and S-benzylthiocarbamate (SBDTC), a nitrogen-sulfur donor, have enormous interest in bioinorganic chemistry. From then onwards, many Schiff bases derived from SMDTC and SBDTC have been reported for their multipurpose coordination chemistry and increasingly important biological activities [3-8]. Because of their unique electrical characteristics, metal complexes with donor atoms in the locations of N, S, and O play a vital role in biological processes. It was reported that the presence of metal ions bonded to biologically active compounds may enhance their activities [9,10]. The existing functional group ( $-NCS-R$ ) of these compounds not only confers chelating properties but also imparts interesting electronic and steric effects to the resulting Schiff bases. As a consequence, SMDTC and SBDTC Schiff bases exhibit boosted stability, reactivity, and selectivity in a wide range of chemical transformations and applications. Di-thiocarbamate and its substituted derivatives have become promising interest in the past few decades. It exists in two tautomeric forms, thione and thiol. The thione group ( $C=S$ ) is relatively unstable in the monomeric system and tends to turn to a stable  $C-S$  single bond by enethionization if at least one hydrogen atom is adjacent to the  $C=S$  bond. The molecule can act as a NOS uni negatively charged tridentate chelating agent [11,12]. Dithiocarbamic acid and the ligands obtained from S-alkyl and S-benzyl esters produce an interesting series of ligands. Various NO, NS, SNNS, SNNNS donor Schiff base ligands have been synthesized by condensation reaction of numerous aldehydes and ketones with SMDTC and SBDTC. The numbers of these types of compounds continue to increase due to the intriguing investigation that different ligands show different biological activity. Metal complexes of Schiff

bases derived from S-alkyldithiocarbazates have been the theme of great contemporary importance because of their fascinating physico-chemical properties and possibly useful biological action [13–18]. However, most preceding revisions on metal-dithiocarbazates were established on tri, tetra, penta, and hexadentate chelating agents [20,21]. Pentadentate ligands derived from S-alkyldithiocarbazates have not received much devotion. Transition metal complexes of the pentadentate ligand formed from 2,6-diacetylpyridine, and S-methyldithiocarbamate (R=CH<sub>3</sub>) have been reported by Majumder *et al.* [22]. Thus, the aims of this review are to show the various synthesis procedure of Schiff bases derived from SBDTC and SMDTC and their metal complexes and studied their structural properties and biological activity to find the potential agent against various pathogenic bacteria and fungi.

## 2. Structural Properties

Two novel quadridentate Schiff base ligands formed from 2,5-hexanedione and S-alkyldithiocarbamic acids and their Ni(II), Zn(II), and Cd(II) complexes having the general formula [M(SNNS)] ((SNNS) is the dinegatively charged ligands) have been synthesized and characterized by M.A. Ali *et al.* The Ni(SNNS) complexes are diamagnetic and square-planar (figure-1). The Zn(SNNS) complexes are assigned polymeric structures with mercaptosulfur-bridging. The Cd(SNNS) complexes presumably have polymeric structures [23].




Figure 1: Structure of Ni(II) complex.

Tris- and bis-ligand complexes of Ni(II) with SBDTC having the general formulae Ni(SBDTC)<sub>3</sub>X<sub>2</sub>·H<sub>2</sub>O (X = Cl, Br and NO) and Ni(SBDTC)<sub>2</sub>X<sub>2</sub> (X = Cl and NCS) respectively, were synthesized and characterized by M.A. Ali *et al.* The ligand acts as a bidentate sulphur-nitrogen chelating agent (figure-2). Both the tris- and bis ligand cationic Ni(II) complexes have high spin magnetic moments of ca. 3.10 B.M. Based on magnetic and spectral information octahedral structures are assigned to these complexes. Under alkaline conditions complexes of the deprotonated ligand having the formulae M(SBDTCA)<sub>2</sub> [M= Ni(II) and Zn(II); SBDTCA= anion of SBDTC] were isolated. The Ni(II) complex is square-planar [24].





Figure2: Structure of ligands

Ni(II), Cu(II), and Co(II) complexes of Schiff bases formed from SBDTC and some aldehydes and ketones were prepared by M.T.H. Tarafder *et al.* [25]. The Schiff base S-methyl-P-N-(2-hydroxyphenyl) methylenedithiocarbazate,  $o\text{-HO.GHGH} = \text{NNHC}(=\text{S})\text{SCH}_3$ , and the corresponding chloro, 5-bromo, 5-nitro, and 3-methoxy derivatives were prepared from the appropriate salicylaldehyde and SMDTC. The acetylaceton Schiff base 2-hydroxy-4-methyl-5,6-diaza-7-thiono-8-thianona-2,4-diene,  $\text{CHC}(\text{OH}) = \text{CHC}(\text{CH}_3) = \text{NNHC}(=\text{S})\text{SCH}_3$ , was also prepared. The Cu(II) ion form stable complexes (figure-3) with the synthesized ligands [26].




Figure3: Structure of the Schiff base Copper(II) complexes

E. Zangrando *et al.* synthesized Ni(II) complex from Schiff base via the condensation reaction of SMDTC with 4-benzyloxybenzaldehyde. The crystallographic results show that the Schiff base exists uninegetively bidentate (NS) in thione tautomeric form with respect to the C=N bond of the group and the Ni(II) complex(figure-4) exhibited a distorted square planar geometry [27].




Figure4: Structure of the Schiff base Nickel(II) complexes

Y. Ji *et al.* synthesized a cobalt (II) complex of general formula  $[\text{Co}(\text{PMBP-smdtc})_2] \cdot 2\text{H}_2\text{O}$ . The novel tridentate Schiff base was derived from the condensation of 1-phenyl-3-methyl-4-benzoyl-5-pyrazolone and SMDTC. The single-crystal X-ray diffraction studies indicated that the complexes showed an octahedrally coordinated compound [28]. Novel bidentate Schiff bases (figure-5) having nitrogen-sulfur donor sequence were synthesized from condensation of racemate camphor, (R)-camphor, and (S)-camphor with methyl hydrazinecarbodithioate. The NS Schiff bases formed complexes of the general formula,  $[\text{M}(\text{NS})_2]$  or  $[\text{M}(\text{NS})_2 \cdot \text{H}_2\text{O}]$  where M is Bi(III) or Ag(I), the expected geometry is octahedral for Bi(III) complexes while Ag(I) is expected to oxidize to Ag(II) forming square planar complexes [29].




Figure5: Synthesis route of novel Schiff base ligand

A one-dimensional indented chain structure is generated when two water molecules make an H-bond along the b-axis, bridging the molecules together. The monobasic bidentate Schiff base S-methyl  $\beta$ -N-(4-methoxyphenylmethyl)methylenedithiocarbamate (NS) and its Bis-chelated metal complexes  $[\text{M}(\text{NS})_2]$ ,  $[\text{M} = \text{Co}(\text{II}), \text{Ni}(\text{II}) \& \text{Cu}(\text{II})]$ , were synthesized and characterized by R. Singh *et al.* [30]. M.A. Ali *et al.* synthesized new Cu(II) complexes of the general formula,

[Cu(mpsme)X] (mpsme=anionic form of the methylpyruvate Schiff base of S-methyldithiocarbazate. based on magnetic and spectral evidence, an oxygen-bridged square-planar structure is assigned to this complex. The crystal and molecular structures of the Schiff base, Hmpsme, and its chlorocopper(II) complex, [Cu(mpsme)Cl] have been determined by X-ray diffraction. The [Cu(mpsme)Cl] complex has a distorted square-planar structure with the ligand coordinated to the Cu(II) ion as a uni-negatively charged tridentate chelating agent via the carbonylic oxygen atom, the azomethine nitrogen atom, and the thiolato sulfur atom. The fourth coordination position around the Cu(II) ion is occupied by the chloride ligand[31]. The distortion from regular square-planar geometry is attributed to the restricted bite size of the ligand. A. Alim *et al.* [32] synthesized Cu(II), Ni(III) Zn(II), Cd(II), Co(II), Sb(III) and Fe(III) containing bidentate Schiff base (figure-6), [Ni-(4-chloro-benzylidene)-hydrazecarbodithioicacid methyl ester] derived from the condensation of SMDTC and P-chlorobenzaldehyde.




Figure6: Structure of the Schiff base metal complexes

2, 6-diacetylpyridinebis(S-benzylidithiocarbazate) ( $H_2SNNNS$ ) pentadentate ligand (figure-7) reacts with Zn(II) and Cd(II) ions forming stable complexes. The crystal and molecular structure of the Zn(II) complex has been determined by X-ray diffraction. The complex is a dimer in which the pyridine nitrogen atom, the azomethine nitrogen atom, and the thiolate sulfur atom from one ligand coordinate to one of the Zn(II) ions whereas the azomethine and thiolate sulfur atoms from another ligand complete penta coordination around the Zn(II) ion, the ligands being coordinated in their deprotonated forms. The coordination geometry of each Zn(II) can be considered as intermediate between a square-pyramidal and trigonal-bipyramidal. The Cd(II) complex is also assigned with a dimeric structure [33].




Figure7: Thione-thiol tautomerism in the ligand (a) thione form (b) thiol form

Ni(II), Cu(II), Pd(II), and Pt(II) complexes of the thiophene-2-aldehyde Schiff bases of SMDTC and SBDTC have been synthesized and characterized by M.A. Ali *et al.* [34]. A.A. Alshaher *et al.* produced a variety of complexes of dithiocarbazate Schiff bases (abbreviated as NiSMdiAP, CoSMdiAP, FeSMdiAP, MnSMdiAP, and ZnSMdiAP, where SMdiAP indicates the Schiff base). S-methyl dithiocarbazate and diacetylpyridine were condensed to create the Schiff base. Every complex functions effectively as a catalyst for oxidation reactions [35]. The novel pyrimidine based NNS tridentate Schiff base ligands (figure-8) has been prepared from condensation of 2-S-methylmercapto-6-methylpyrimidine-4-carbaldehyde and S-methyl/S-benzyl dithiocarbazate by S. Roy *et al.* All the complexes exhibited a distorted octahedral arrangement with an  $N_4S_2$  chromophore around the central metal ion. Each ligand molecule binds the metal ion using azomethine pyrimidyl nitrogen and thiolato sulfur. Among the complexes, the Ni(II) complex, behaves as a neutral tridentate, and the other complexes the Co(III) and Fe(III) complexes, the ligand molecules behave as monoanionic tridentate. All the complexes were investigated by single crystal X-ray diffraction and indicated a distortion from an octahedral geometry of the coordination environment was observed [36].




Figure8: Synthesis route of novel Schiff base ligand

New bis-chelated Zn(II) and Cd(II) complexes of empirical formula,  $[M(\text{mpsme})_2]$  (mpsme) the anionic form of the tridentate ONS donor ligand formed from methylpyruvate and S-methyldithiocarbazate) have been synthesized by M.A. Ali *et al.* [37]. Both complexes are allotted a distorted octahedral geometry in which the ligands are organized meridionally around the metal ions. The distortion from systematic octahedral geometry is attributable to the constrained bite angles of the ligand. Condensation of 2,6-diacetylpyridine (dap) with SMDTC in a 1:2 molar ratio yields a bicondensed pentadentate Schiff base ( $H_2\text{dapsme}$ ) which reacts with  $K_2MCl_4$  ( $M = \text{Pd(II)}, \text{Pt(II)}$ ) giving stable complexes of empirical formula,  $[M(\text{dapsme})] \cdot 0.5\text{Me}_2\text{CO}$  [38]. These complexes have been characterized by a variety of physico-chemical techniques and Single X-ray crystal measurement. The complex has a distorted square-planar structure in which the ligand is coordinated to the Pd(II) ion as a uninegatively charged tridentate chelating agent via the pyridine nitrogen atom, the azomethine nitrogen atom, and the thiolato sulfur atom; the oxygen atom of the acetyl group does not participate in coordination. Two new Co(III) complexes with S-benzyl-N-(5-methylpyrazole-3-yl)methylenedithiocarbazate ( $H_2L$ ),  $[\text{Co}(\text{HL})_2]\text{NO}_3 \cdot \text{EtOH}$  and  $[\text{Co}(\text{HL})(L)] \cdot \text{H}_2\text{O}$ , have been synthesized and characterized using single-crystal X-ray diffraction and spectroscopic (electronic, IR and NMR) techniques. Both complexes exhibited a distorted octahedral geometry and ligands in the mer configuration [39]

### 3. Biological Properties

#### 3.1: Antimicrobial Activity

SMDTC and SBDTC when condensed with glyoxal, five coordinated ligands will produce that also make stable crystalline complexes of the general formula,  $M(SNNS)$  ( $M = Ni(II), Cu(II), Zn(II)$  and  $Cd(II)$ ;  $SNNS =$  ligand dianion). The Schiff bases and metal complexes display good antifungal and antibacterial properties. All the complexes showed activities against two pathogenic bacteria (*E. coli* and *B. subtilis*) and three phytopathogenic fungi (*D. oryzae*, *I. padwickii*, and *F. moniliforme*) [40]. M.E. Hossain *et al.* prepared  $Cu(II)$  complexes containing Schiff base ligands (figure-9) derived from the condensation reactions of 2-benzoylpyridine with SMDTC and SBDTC respectively. The antifungal and antibacterial properties of the Schiff bases and their  $Cu(II)$  complexes have been evaluated against three pathogenic fungi and two bacteria. The Schiff bases and  $Cu(II)$  complexes display moderate antifungal activity, but their activities are less than that of the commercially important antifungal agent nystatin [41].




Figure9: Thione-thiol tautomerism in the ligand (a) thione form (b) thiol form

M.K.E. Zahan *et al.* synthesized a series of metal complexes of  $Mn(II)$ ,  $Fe(III)$ ,  $Co(II)$ ,  $Ni(II)$ ,  $Cu(II)$ , and  $Sb(III)$  containing the bidentate Schiff base derived from the condensation of S-methyldithiocarbamate and cinnamaldehyde. All the complexes of Schiff base exhibited antibacterial activity against four gram-positive (*Bacillus subtilis*, *Staphylococcus aureus*, *Bacillus megatherium*, *Streptococcus- $\beta$ -haemolyticus*) and four gram-negative (*Escherichia coli*, *Shigella dysenteriae*, *Shigella sonnei*, *Shigella shiga*) and antifungal activity against four *Aspergillus* (*Candidus*, *Penicillium Marneffeii*, *Candida Albicans*, *Aspergillus niger*) [42]. M.L. Low *et al.* [43] synthesized a series of  $Cu(II)$  complexes from SMDTC and SBDTC with methyl levulinate, levulinic acid, and 4-carboxybenzaldehyde. The synthesized ligands and their respective  $Cu(II)$  complexes showed moderate antibacterial activity against both Gram-negative and Gram-positive bacteria. The Copper complexes were highly effective against *Staphylococcus aureus* with a minimum inhibition concentration (MIC) of 0.5-1  $\mu M$ .  $Cu(II)$  complexes of the general formula  $[Cu(Ap-SMe)X]$  or  $[Cu(Ap-SBz)X]$  (where HAp-SMe and HAp-SBz, respectively, represent the 2-acetylpyridine Schiff bases of SMDTC or SBDTC;  $X = Cl, Br, NO_3$ ) have been prepared and characterized by M.E. Hossain *et al.*. The geometry of the complex was a distorted square pyramid with the NNS tridentate ligand and an oxygen atom of the nitrate ion occupying the basal plane. The fifth coordination position was occupied by oxygen from the nitrate ion. The anti-fungal and anti-bacterial properties of the Schiff bases and their copper(II) complexes have been evaluated against the phytopathogenic fungi *A. solanyi*, *F. equiseti*, and *M. phaseolina* and the pathogenic bacteria *E. coli* and *S. aureus*. The fungi toxicity of the five-coordinated  $[Cu(Ap-SBz)NO_3]$  complex approaches that of nystatin, whereas the Schiff base HAp-SMe and its

copper(II) complex  $[\text{Cu}(\text{Ap-SMe})\text{NO}_3]$  display significant antibacterial activity against *E. coli* and *S. aureus*[44]. A new Schiff Base ligand  $[(\text{SBDTC})_2\text{AP}]$  has been synthesized by the 1:2 condensation of acetophenone and SBDTC. A group of selected ions such as Cu(II), Cd(II), Zn(II), Ni(II), and Co(II) complexes of  $[(\text{SBDTC})_2\text{AP}]$  have been prepared and characterized by N.Ameramaet *al.*The Schiff base(figure-10), and its metal complexes have been evaluated for their biological activities against *Staphylococcus aureus*, *S.Typhy M.H.*, and *Aeromonas sp.*All metal complexes were inactive against *S.Typhy M.H.*[45]




Figure10: Synthesis route of Schiff base

A new tridentate nitrogen–oxygen–sulfur Schiff base has been prepared from the condensation reaction of S-methyldithiocarbamate and methylisatin by M.A.F.A. Manan *et al.*The Schiff base was found to be selectively active against the selected Gram-positive bacterial strains (*Bacillus subtilis* and *Staphylococcus aureus*) [46].The nitrogen–oxygen–sulfur tridentate Schiff base ligand prepared from condensation of S-benzylthiocarbamate (SBDTC) and salicylaldehyde was reported by N.A. Bituet *al.*The complexes of Zr(IV), Th(IV), and U(VI) ions have been synthesized from the derived Schiff base. The antibacterial activity of all the compounds has been evaluated using Kanamycin (K-30) as a standard against two Gram-negative (*Escherichia coli* and *Shigella dysenteriae*), and two Gram-positive (*Staphylococcus aureus* and *Bacillus subtilis*) bacteria. Against every examined organism, the Zr(IV) complex is less effective against bacteria than the U(VI) and Th(IV) complexes [47].Cu(II) complexes of formulas,  $[\text{Cu}(\text{NNS})(\text{NO}_3)(\text{H}_2\text{O})]\cdot\text{H}_2\text{O}$  and  $[\text{Cu}(\text{NNS})_2]$  (NNS=anionic form of the 6-methyl-2-formylpyridine 4-N-dimethylthiosemicarbazone) have been synthesized and characterized by M.A. Aliet *al.*. The ligand and the metal complexes were screened for antifungal activity against the *phytopathogenic fungi*, *A. alternata*, *F. equiseti*, and *M. phaseolina* but the complexes are less active than the free ligand due to less solubility in non-aqueous solvents. The copper complexes may have less solubilities in non-aqueous solvents than the Schiff base [48]. The di-2-pyridylketone Schiff base of S-methyldithiocarbamate (Hdpksme) reacts with Ni(II) salts giving both mono- and bis-chelated complexes were synthesized by M.A. Aliet *al.* The ligand and its Ni(II) complexes exhibited weak antimicrobial activity against the pathogenic bacteria, *S. dysenteriae*, *B. cereus*, *S. aureus*, and *E. coli* and the fungi, *A. alternata* and *M. phaseolina* [49].A tridentate ONS Schiff base derived from the condensation of SBDTC with salicylaldehyde has been synthesized. The metal complexes of general formula  $[\text{M}(\text{ONS})\text{X}]$ ,  $[\text{M}=\text{Ni}(\text{II}), \text{Cu}(\text{II}), \text{Cr}(\text{III}), \text{Sb}(\text{III}), \text{Zn}(\text{II}), \text{Zr}(\text{IV}) \text{ or } \text{U}(\text{VI}) \text{ with } \text{X}=\text{H}_2\text{O}, \text{Cl}]$ .Antimicrobial tests indicate that the Schiff base and five of the metal complexes of Cu(II), Ni(II), U(VI), Zn(II), and Sb(III) are strongly active against bacteria. Ni(II) and Sb(III) complexes were the most effective against *Pseudomonas aeruginosa*, while the Cu(II) complex proved to be best against *Bacillus cereus*. Antifungal activities were also noted with the Schiff base and the U(VI) complex. These compounds showed positive results against *Candida albicans* fungi, however, none of them were effective against *Aspergillus ochraceous* fungi [50]. M.A. Ali and A.H. Mirza prepared Ni(II) complex with NOS coordinating Schiff base ligand derived from the condensation reaction of N-

methyl-S-methyldithiocarbazate with 2-hydroxybenzaldehyde or 5-bromo-2-hydroxybenzaldehyde. The synthesized ligands and Ni(II) complexes were screened for antifungal activity against three pathogenic fungi. The ligands display moderate fungi toxicities against these organisms but their nickel(II) complexes are less active than the free ligands. The chelation of nickel(II) complexes with dithiocarbazate lessens the antifungal activity [51]. Cu(II) complexes containing Schiff base ligand  $\alpha$ -N-methyl-S-methyl-b-N-(2-pyridyl)methylenedithiocarbazate(NNS) were synthesized and characterized by M.A. Ali *et al.* Antimicrobial tests indicated that Schiff base is inactive against the bacteria, *Bacillus subtilis* (mutant defective DNA repair), *Pseudomonas aeruginosa*, methicillin resistant *Staphylococcus aureus* and *Bacillus subtilis* (wild type) and weakly active against the fungi, *Candida albicans*, *Candida lypolytica*, *Saccharomyces cereviseae* and *Aspergillus ochraceous* but its Cu(II) complexes, Cu(NNS) $X_2$  are strongly active against these organisms [52]. Mono- and bis-chelated Cd(II) complexes of the di-2-pyridylketone Schiff base of S-methyldithiocarbazate (Hdpsme) have been synthesized by M. A. Ali and A. H. Mirza. The Schiff base and its Cd(II) complexes were screened for antibacterial activity against *Shigella dysenteriae*, *Bacillus cereus*, *Staphylococcus aureus*, and *Escherichia coli*. They are also mildly fungitoxic against the phytopathogenic fungi, *Alternaria alternata* and *Macrophomina phaseolina* [53].

### 3.2 Anti-Cancer Activity

M.L. Low *et al.* prepared Cu(II), Zn(II), and Re(I) complexes of the general formula of Cu(SBCM) $_2$ , Zn(SBCM) $_2$ , and Re $_2$ (SBCM) $_2$  derived from Schiff base of the S-Benzyldithiocarbazate and 3-acetylcoumarin. X-ray diffractometry (XRD) indicated that Rhenium exhibited a centrosymmetric dimeric complex via the Re-S-Re bridge and Cu(II), Zn(II) coordinated to a bidentate ligand of unintuitively thiolate and azomethine nitrogen atom of Schiff base. Cytotoxicity assesses the apparent enhancement of SBCM-H's bioactivity following complexation. Cu(II) and Re(I) complexes have been discovered to exhibit activity against cell lines that represent human breast adenocarcinoma cancer cell lines MCF-7 and MDA-MB-231 [54]. M.X. Li *et al.* have synthesized a series of Cu(II) and Zn(II) complexes of 2-benzoylpyridine Schiff bases derived from S-methyldithiocarbazate(L<sup>1</sup>), and S-phenyldithiocarbazate (L<sup>2</sup>) of general formula of Cu $_2$ (L<sup>1</sup>) $_2$ (CH $_3$ COO)](ClO $_4$ )(1), [Zn $_2$ (L<sup>1</sup>) $_2$ (ClO $_4$ ) $_2$ ](2), [Zn(L<sup>2</sup>) $_2$ ](3). A single-crystal X-ray diffraction study showed that the Cu(II) complex (1) adopts square-pyramidal and Zn(II) complex (2) distorted pentagonal bipyramidal geometry, Zn(II) complex (3) octahedral coordination geometry. It should be emphasized that Zn(II) complex (3) effectively inhibits the K562 leukemia cell line lower than HL<sup>2</sup> and Cu(II) complex (1) inhibits the K562 leukemia cell line lower than HL<sup>1</sup>. Importantly, IC $_{50}$  values of Zn(II) complexes 2 and 3 are higher in QSG7701 cell line than in K562 leukemia cell line, indicating both of them have tumor cells selectivity [55]. M.L. Low *et al.* prepared two novel Schiff base and their Cu(II) complexes. The Schiff base synthesized from the condensation reaction of S-methyl- and S-benzyldithiocarbazate with 2,5-hexanedion. Cu(II) complexes are strongly active against human breast adenocarcinoma cancer cell lines MDA-MB-231 and MCF-7 [56]. M.K.B. Breaket *et al.* synthesized a series of macrocyclic Schiff base ligands and their Cu(II) complexes (1-9) via a condensation reaction of various dicarbonyls with S-methyl

dithiocarbazate and *S*-benzyl dithiocarbazate. Among the complexes, Complex (1) exhibited the most cytotoxic effect among the complexes against MDA-MB-231 and MCF-7 breast cancer cells, with IC<sub>50</sub> values of 1.7 μM and 1.4 μM, respectively[57]. A series of novel metal complexes possessing general formula of [M(L)<sub>2</sub>] [M= Cu(II)(1), Ni(II)(2), and Zn(II)(3) and Schiff base derived from *S*-(4-methylbenzyl)dithiocarbazate and 2-methoxybenzaldehyde by E.N.M. Yusof *et al.* According to the cytotoxic findings, complex (1) significantly reduced the growth of MCF-7 and MDA-MB-213 cancer cell lines. The compounds demonstrated strong hydrogen bonding, Van der Waals force, and hydrophobic DNA-binding capabilities[58]. The Schiff bases RR'C=NNHC(S)SCH<sub>3</sub> derived from *S*-methylthiocarbazate have been prepared. The mass spectra of the Schiff bases having R = Me and R' = Ph, 2-furyl, and 2-thienyl and R = R' = Ph are similar and display strong peaks for the molecular ion M and the ions M-CH<sub>3</sub>S, M-CH<sub>3</sub>SH, R'C=NH, R'CN, R'H, R', and RI-H. A mechanism of fragmentation is proposed. Complexes of the Schiff bases of the type Met(RR'C=NN=CSSCH<sub>3</sub>)<sub>2</sub> (Met = Ni, Cu, Zn, Pd, and Pt) have been isolated in most instances. When R' = Bu" or Bu', nickel ion catalyses the hydrolysis of the Schiff base with the concomitant formation of the *S*-methylthiocarbazate complex [Ni(H, NNHCSSCH<sub>3</sub>,)] (NO). The metal complexes are being tested for anti-cancer activity [59]. C.L. Chena *et al.* have synthesized transition metal complexes of Co(II) and Mn(II) containing 2-acetylpyridine *S*-methylthiocarbazate. Biological activity studies carried out in vitro against the K562 leukemia cancer cell line have shown that the free ligand and its metal complexes exhibited significant and different antitumor activity since they exhibit IC<sub>50</sub> values in the μM range [60]. More than 70 relieved thiosemicarbazones and several metal complexes of each have been analyzed for their antifungal activity. Their action is significantly affected by the replaced groups involved at both <sup>1</sup>N and <sup>4</sup>N of the thiosemicarbazone moiety. The greatest activity occurs for 2-substituted pyridine thiosemicarbazones with differences experiential for 2-formylpyridine, 2-acetylpyridine, and 2-benzoylpyridine derivatives and their metal complexes. Further, there are activity differences for <sup>4</sup>N-alkyl-, <sup>4</sup>N-aryl-, <sup>4</sup>N-dialkyl- and 3-azacyclothiosemicarbazones and their metal complexes as well as changes in the substituent size among each of these subgroups. Cu(II) complexes are often more active than the uncomplexed thiosemicarbazones, with the latter showing similar activity to Ni(II) complexes in many instances. The reduction potential of the thiosemicarbazone ligand in a Cu(II) complex, the strength of the ligand field, and various spectral properties can be correlated to the inhibitory activity [61]. M.A. Ali *et al.* synthesized Pd(II) and Pt(II) complexes containing Schiff bases of *S*-methyl- and *S*-benzylthiocarbazate and characterized them by a variety of physicochemical techniques. The complex has a distorted cis-square planar structure with the ligands coordinated to the Pd(II) ions as uninegatively charged bidentate NS chelating agents via the azomethine nitrogen and the mercaptide sulfur atoms. The distortion from a regular square-planar geometry is attributed to the restricted bite angles of the ligands. Antimicrobial tests indicate that the Schiff bases exhibit strong activities against the pathogenic bacteria, *Bacillus subtilis* (mutant defective DNA repair), methicillin-resistant *Staphylococcus aureus*, *B. subtilis* (wild type) and *Pseudomonas aeruginosa* and the fungi, *Candida albicans* (CA), *Candida lyptica* (2075),

*Saccharomyces cerevisiae* (20341) and *Aspergillus ochraceus* (398) the activities exhibited by these compounds being greater than that of the standard antibacterial and antifungal drugs, *streptomycin* and *nystatin*, respectively. The Pd(II) and Pt(II) complexes were inactive against most of these organisms but, the microbe, *Pseudomonas aeruginosa* shows strong sensitivity to the Pt(II) complexes. Screening of the compounds for their cytotoxicities against *T-lymphoblastic leukemia* cancer cells has shown that the acetone Schiff base of S-methyldithiocarbamate (Hasme) exhibits a very weak activity, whereas the S-benzyl derivative (Hasbz) is inactive. However, the Pd(II) complexes exhibit strong cytotoxicities against this cancer; their activities being more than that of the standard anticancer drug, tamoxifen. The [Pt(asme)] complex exhibits very weak cytotoxicity, 2 whereas [Pt(asbz)] is inactive against leukemic cells [62]. The transition metal complexes were synthesized from the Schiff base by L.Z. Zhang *et al.* The two novel Schiff bases derived from S-methyldithiocarbamate (HL<sup>1</sup>) and S-benzylthiocarbamate (HL<sup>2</sup>) with pyridine-2-carboxaldehyde. Among the Schiff base and their complexes, only HL<sup>1</sup> and manganese complex II show significant antitumor activity against K562 leukemia cell line, since they exhibit IC<sub>50</sub> values in the μM range [63].

### 3.3 Antioxidant Activity

A series of metal complexes were synthesized from the novel Schiff base ligand by S.A. Elsayed *et al.* The Schiff base was derived from the condensation reaction of pyridoxal and SBDTC. The antioxidant activity of the complexes was measured by 2,2-diphenyl-1-picrylhydrazyl radical (DPPH). Among the complexes, the oxovanadium (IV) exhibited the highest scavenging activity against DPPH with lowest value of IC<sub>50</sub> [64]. M.T.H. Tarafder *et al.* prepared Bis-chelated Cu(II), Ni(II) and Zn(II) complexes containing two new isomeric Schiff bases, S-methyl-b-N-(2-furylmethyl)methylenedithiocarbamate (NS') and S-methyl-b-N-(5-methyl-2-furyl)methylenedithiocarbamate (NS''). The antioxidant activity of the complexes was tested by FTC methods which indicated the peroxide value in the primary stage of lipid oxidation. The data evaluated that SBDTC exhibited a higher antioxidant activity than the α-tocopherol (vitamin E) [65]. M.T.H. Tarafder *et al.* prepared bidentate and quartidentate Schiff base ligands derived from the condensation reaction of S-benzylthiocarbamate with 2,3 butanedione. Spectroscopic evidence suggested that the prepared ligand acts as an NS and NNSS coordinating agent during complexation with Cu(II) and Ni(II) ions. Antioxidant activity has shown that nickel and copper complexes were more active than those of Vitamin E [66].

### Conclusions

This review shows various synthesis modes of Schiff bases derived from SBDTC and SMDTC and their metal complexes. This article also explains how the Schiff base metal complexes are potentially active against various pathogenic bacteria and fungi. Some of the metal complexes are potentially effective against cancer cells. Improvements in bioinorganic chemistry are important for enlightening the design of compounds to moderate toxic side-effects and understand their mechanisms of action. This review reveals that the Schiff bases and their

metal complexes derived from SBDTC and SMDTC could be a suitable approach to progress innovative therapeutic tools for medical treatment.

## References

- [1]. B.M. Ismael, M.A. El-Nawawy, H.A. Gomaa, O. El-Gammal, "Divalent transition metal complexes of multidentate nitrogen, oxygen and sulfur-containing ligand: Design, spectroscopic, theoretical molecular modeling and antioxidant like activity", *Egyptian Journal of Chemistry*, 2022; 65(9): 639-655. <https://doi.org/10.21608/ejchem.2022.118264.5331>
- [2]. B. Tang, L. Zheng, X. Dai, H. Chen, "Nitrogen/oxygen co-doped porous carbons derived from a facilely synthesized Schiff-base polymer for high-performance supercapacitor", *Journal of energy storage*, 2019; 26: 100961. <https://doi.org/10.1016/j.est.2019.100961>
- [3]. A.A.A. Emara, A.M. Taha, M.M. Mashaly, O.M.I. Adly, "Synthesis, spectroscopic, thermal and anti-microbial studies of transition metal complexes of hydrazone derived from 4,6-diacetylresorcinol and S-methyldithiocarbamate", *European Journal of Chemistry*, 2015; 6(2): 107–116. <https://doi.org/10.5155/eurjchem.6.2.107-116.1162>
- [4]. S.S. Bhuyar, H.D. Junej, L.M.J. Paliwal, R.G. Chaudhary, "Synthesis, characterization and thermal degradation of some coordination polymers with terephthalaldehyde bis (S-methyldithiocarbamate)", *Journal of the Chinese Advanced Materials Society*, 2015; 3(1): 17-31. <https://doi.org/10.1080/22243682.2014.981859>
- [5]. T.B.S.A. Ravoof, K.A. Crouse, M.I.M. Tahir, A.R. Cowley, M.A. Ali. "Synthesis, characterization and bioactivity of mixed-ligand Cu(II) complexes containing Schiff bases derived from S-benzylidithiocarbamate and saccharinate ligand and the X-ray crystal structure of the copper-saccharinate complex containing S-benzyl-b-N-(acetylpyrid-2-yl)methylenedithiocarbamate", *Polyhedron*, 2007; 26(6): 1159-1165. <http://dx.doi.org/10.1016/j.poly.2006.03.007>
- [6]. T.B.S.A. Ravoof, K.A. Crouse, E.R.T. Tiekink, M.I.M. Tahir, E.N.M. Yusof, R. Rosli, "Synthesis, characterization and biological activities of S-2- or S-4-methylbenzyl-β-N-(di-2-pyridyl)methylenedithiocarbamate and Cu(II), Ni(II), Zn(II) and Cd(II) complexes", *Polyhedron*. 2017; 133: 383-392. <https://doi.org/10.1016/j.poly.2017.05.053>
- [7]. R. Zahan, M.S. Rahi, M.C. Sheikh, R. Miyatake, E. Zangrando, T. Naz, M.A.A.A. Islam, M.A. Reza. "Design, synthesis and X-ray structural studies of novel [acetonitrile-3-benzyl-2,4-dihydroxyphenylmethylene)hydrazinecarbodithioato-3-N', S, O] nickel(II) complex that potently inhibit cell proliferation through regulation of apoptosis related genes", *Applied Organometallic Chemistry*, 2019; 33(1): e4601. <https://doi.org/10.1002/aoc.4601>
- [8]. M.A.A.A. Islam, M.C. Sheikh, M.A. Mumit, R. Miyatake, M.A. Alam, M.O.A. Mondal, "Synthesis, characterization and antimicrobial activity of a bidentate NS Schiff base of S-

- benzyl dithiocarbazate and its divalent complexes”, *Journal of Coordination Chemistry*, 2016; 69(23): 3580–3592. <https://doi.org/10.1080/00958972.2016.1233329>
- [9]. M.K.E.Zahan, M.F. Hossen, R.H. Ansary, R. Zamir, M.A. Asraf, “Synthesis, characterization and biological activity evaluation of Ni(II) and Cu(II) complexes of a Schiff base derived from S-benzyl dithiocarbazate (SBDTC) and Cinnamaldehyde”, *International Journal of Chemistry Studies*, 2020; 4(2): 18-24.
- [10]. J. Lin, M. Lin, Y. Su, M. Liu, H. Zeng, Y. Cai, “Synthesis and spectroscopic studies of the cobalt(II) complex with methyl 2-pyridylmethylidenehydrazinecarbodithioate (HNNS)”, *Polyhedron*, 2007; 32:338-344. <https://doi.org/10.1007/s11243-006-0173-3>
- [11]. O.M.I. Adly, H.F.E. Shafiy, “New metal complexes derived from S-benzyl dithiocarbazate (SBDTC) and chromone-3-carboxaldehyde: synthesis, characterization, antimicrobial, antitumor activity, and DFT calculations. *Journal of Coordination Chemistry*, 2019; 72(2):218-238. <https://doi.org/10.1080/00958972.2018.1564912>
- [12]. M.A. Ali, S.E. Livingstone, “Metal complexes of sulphur-nitrogen chelating agents”, *Coordination Chemistry Reviews*, 1974; 13(2-3):101-132. [https://doi.org/10.1016/S0010-8545\(00\)80253-2](https://doi.org/10.1016/S0010-8545(00)80253-2)
- [13]. M.A. Ali, A.H. Mirza, G.A. Fong, “Synthesis, characterization and X-ray crystal structures of the bis–ligand zinc(II) and cadmium(II) complexes of the methylpyruvate Schiff base of S-methyl dithiocarbazate”, *Transition Metal Chemistry*, 2004; 29: 613-619. <https://doi.org/10.1007/s11243-004-2511-7>
- [14]. M.A. Ali, A.H. Mirza, M.Haniti, S.A. Hamid, N. Aminath, P.V. Bernhardt, “Synthesis, characterization and X-ray crystal structures of thiolate sulfur-bridged dimeric copper(II) complexes of the 2-aminoacetophenone Schiff base of S-methyl dithiocarbazate”, *Polyhedron*, 2012; 47: 79-86. <http://dx.doi.org/10.1016/j.poly.2012.08.024>
- [15]. M.A. Ali, H.J.H.A. Bakar, A.H. Mirza, S.J. Smith, L.R. Gahan, P.V. Bernhardt, “Preparation, spectroscopic characterization and X-ray crystal and molecular structures of nickel(II), copper(II) and zinc(II) complexes of the Schiff base formed from isatin and S-methyl dithiocarbazate (His-sme)”. *Polyhedron*, 2008; 27: 71-79. <https://doi.org/10.1016/j.poly.2007.08.022>
- [16]. M.A. Ali, A.H. Mirza, F.H. Bujang, M.H.S.A. Hamid, P.V. Bernhardt, “Synthesis, characterization and X-ray crystallographic structural study of copper(II) and nickel(II) complexes of the 2-quinoline carboxaldehyde Schiff base of S-methyl dithiocarbazate (Hqaldsme)”, *Polyhedron*, 2006; 25(17): 3245-3252. <https://doi.org/10.1016/j.poly.2006.05.027>
- [17]. M. Nazimuddin, M.A. Ali, F.E. Smith. “The preparation and characterization of some nickel(II) and copper(II) complexes of ONS ligands”, *Polyhedron*, 1991; 10(12): 1327-1332. [https://doi.org/10.1016/S0277-5387\(00\)81265-6](https://doi.org/10.1016/S0277-5387(00)81265-6)
- [18]. M. Rahman, M.A. Ali, P. Bhattacharjee, M. Nazimuddin, “Synthesis and characterization of some metal complexes containing a quadridentate NSSN and a sexidentate ONSSNO ligand”, *Polyhedron*, 1991; 10(8): 823-827. [https://doi.org/10.1016/S0277-5387\(00\)86115-X](https://doi.org/10.1016/S0277-5387(00)86115-X)

- [19]. S.M.M.H. Majumder, M.A. Ali, F.E.Smith, M.A.U.Mridha, "Synthesis, characterization and fungitoxicities of metal chelates of a pentadentate  $N_3S_2$  ligand", *Polyhedron*, 1988; 7(21): 2183-2187. [https://doi.org/10.1016/S0277-5387\(00\)81800-8](https://doi.org/10.1016/S0277-5387(00)81800-8)
- [20]. Y. Tian, C. Duan, Z. Lu, X. You, H. Fun, S. Kandasamy, "Crystal structure and spectroscopic studies on metal complexes containing ns donor ligands derived from S-benzyl dithiocarbazate and p-dimethylaminobenzaldehyde", *Polyhedron*, 1996; 15(13): 2263-2271. [https://doi.org/10.1016/0277-5387\(95\)00477-7](https://doi.org/10.1016/0277-5387(95)00477-7)
- [21]. M.A. Ali, M. Nazimuddin, R. Shaha, R.J. Butcher, J.C. Bryan, "Synthesis and characterization of bis-chelated nickel(II) complexes of the methylpyruvate Schiff bases of S-alkyl dithiocarbazates and the X-ray crystal structure of the  $[Ni(ONSM_e)_2]$  complex", *Polyhedron*, 1998; 17(22): 3955-3961. [https://doi.org/10.1016/S0277-5387\(98\)00189-2](https://doi.org/10.1016/S0277-5387(98)00189-2)
- [22]. S. Bhattacharjee, R. Bhattacharyya, "Synthesis, characterization, electrochemistry and oxo-transfer kinetics of oxomolybdenum-(VI), -(V) and -(IV) complexes with ONS donors", *Journal of Chemical Society, Dalton Transactions*, 1992; 1357-1364. <https://doi.org/10.1039/DT9920001357>
- [23]. M.A. Ali, S.M.G. Hossain, S.M.M.H. Majumder, M.N. Uddin, M.T.H. Tarafder, "Synthesis and characterization of some new nickel(II), zinc(II) and cadmium(II) complexes of quadridentate SNNS ligands", *Polyhedron*, 1987; 6(8): 1653-1656. [https://doi.org/10.1016/S0277-5387\(00\)80766-4](https://doi.org/10.1016/S0277-5387(00)80766-4)
- [24]. M.A. Ali, M.T.H. Tarafder, "Metal complexes of sulphur and nitrogen-containing ligands: complexes of S-benzyl dithiocarbazate and a Schiff base formed by its condensation with pyridine-2-carboxaldehyde", *Journal of Inorganic and nuclear Chemistry*, 1977; 39(10): 1785-1791. [https://doi.org/10.1016/0022-1902\(77\)80202-9](https://doi.org/10.1016/0022-1902(77)80202-9)
- [25]. M.T.H. Tarafder, M.A.J. Miah, R.N. Bose, "Metal complexes of some Schiff bases derived from S-benzyl dithiocarbazate", *Journal of Inorganic and nuclear Chemistry*, 1981; 43(1): 3151-3157. [https://doi.org/10.1016/0022-1902\(81\)80079-6](https://doi.org/10.1016/0022-1902(81)80079-6)
- [26]. M.T.H. Tarafder, S.E. Livingstone, D.J. Phillips, "Metal chelates of dithiocarbazic acid and its derivatives. VI. antiferromagnetic and ferromagnetic interactions in some copper(II) complexes of salicylaldehyde and acetylacetone Schiff bases derived from s-methyl dithiocarbazate", *Inorganica Chimica Acta*, 1973; 7(2): 179-186. [https://doi.org/10.1016/S0020-1693\(00\)94808-4](https://doi.org/10.1016/S0020-1693(00)94808-4)
- [27]. E. Zangrando, M.T. Islam, M.A.A.A.A. Islam, M.C. Sheikh, M.T.H. Tarafder, R. Miyatake, R. Zahan, M.A. Hossain, "Synthesis, characterization and bio-activity of nickel(II) and copper(II) complexes of a bidentate NS Schiff base of S-benzyl dithiocarbazate", *Inorganica Chimica Acta*, 2015; 427: 278-284. <https://dx.doi.org/10.1016/j.ica.2014.12.014>
- [28]. Y. Ji, L. Liu, D. Jia, Y. Qiao, K. Yu, "Synthesis and crystal structure of Schiff base transition metal complex  $[Co(PMBP-smdtc)_2] \cdot 2H_2O$ ", *Journal of Chemical Crystallography*, 2002; 32(12): 505-509. <https://doi.org/10.1023/A:1021105128641>

- [29]. I.H.T. Al-Karkhi, A.K. Yaseen, H.T. Sadeq, "Synthesis and characterization of novel methyl 2-(1,7,7-trimethylbicyclo[2.2.1]hept-2-ylidene)hydrazinecarbodithioate Schiff bases derived from methylhydrazine carbodithioate and their bi(iii) and ag(ii) complexes", *Baghdad Science Journal*, 2013; 10(3): 609-617. <https://doi.org/10.21123/bsj.10.3.607-617>
- [30]. R. Singh, Y.C.Goswami, R. goswami, "Semi-empirical & Experimental Investigation on coordination behavior of S-methyl-β-N-(4-methoxyphenylmethyl)methylenedithiocarbazate Schiff base towards Co(II), Ni(II) and Cu(II) metal ions". *Journal of Chemistry*. 2011; 2(1): 42-50.
- [31]. M.A. Ali, A.H. Mirza, R.J. Butcher, "Synthesis and characterization of copper(II) complexes of the methylpyruvate Schiff base of S-methyldithiocarbazate (Hmpsme) and the X-crystal structures of Hmpsme and [Cu(mpsme)Cl]", *Polyhedron*, 2001; 20: 1037–1043. [https://doi.org/10.1016/S0277-5387\(01\)00730-6](https://doi.org/10.1016/S0277-5387(01)00730-6)
- [32]. A. Alim, M.K.E.Zahan, M.M. Haque, M.T.H Tarafder, "Synthesis and characterization of some metal complexes of Cu(II), Ni(II), Zn(II), Cd(II), Sn(II), Co(II), Sb(III) and Fe(III) containing bidentate Schiff base of SMDTC". *Science Journal of Chemistry*. 2014; 3(3): 35-39. <https://doi.org/10.11648/j.sjc.20150303.11>
- [33]. M.A. Ali, A.H. Mirza, C.W. Voo, A.L. Tan, P.V. Bernhardt, "The preparation of zinc(II) and cadmium(II) complexes of the pentadentate N<sub>3</sub>S<sub>2</sub> ligand formed from 2,6-diacetylpyridine and S-benzyl dithiocarbazate (H<sub>2</sub>SNNNS) and the X-ray crystal structure of the novel dimeric [Zn<sub>2</sub>(SNNNS)<sub>2</sub>] complex", *Polyhedron*, 2003; 22: 3433–3438. <https://doi.org/10.1016/j.poly.2003.08.004>
- [34]. M. A. Ali, A. H. Mirza, R. J. Butcher, M. Rahman, "Nickel(II), copper(II), palladium(II) and platinum(II) complexes of bidentate SN ligands derived from S-alkyldithiocarbazates and the X-ray crystal structures of the [Ni(tasbz)<sub>2</sub>] and [Cu(tasbz)<sub>2</sub>].CHCl<sub>3</sub> complexes", *Transition Metal Chemistry*, 2000; 25: 430–436. <https://doi.org/10.1023/A:1007009810572>
- [35]. A.A. Alshaheri, M.I.M. Tahir, M.B.A. Rahman, T. Begum, T.A. Saleh, "Catalytic oxidation of cyclohexane using transition metal complexes of dithiocarbazate Schiff base", *Chemical Engineering Journal*, 2017; 327: 423-430. <http://dx.doi.org/10.1016/j.cej.2017.06.116>
- [36]. S. Roy, T.N. Mandal, A.K. Barik, S. Pal, S. Gupta, A. Hazra, R.J. Butcher, A.D. Hunter, M. Zeller, "Metal complexes of pyrimidine derived ligands—Syntheses, characterization and X-ray crystal structures of Ni(II), Co(III) and Fe(III) complexes of Schiff base ligands derived from S-methyl/S-benzyl dithiocarbazate and 2-S-methylmercapto-6-methylpyrimidine-4-carbaldehyde", *Polyhedron*, 2007; 26: 2603–2611. <https://doi.org/10.1016/j.poly.2007.01.006>
- [37]. S.K. Kar, M.A. Ali, A.H. Mirza, G.A. Fong, "Synthesis, characterization and X-ray crystal structures of the bis-ligand zinc(II) and cadmium(II) complexes of the methylpyruvate Schiff base of S-methyldithiocarbazate", *Transition Metal Chemistry*, 2004; 29: 613–619. <https://doi.org/10.1007/s11243-004-2511-7>

- [38]. M.A. Ali, A.H. Mirza, S.T.W. Keng, "Palladium(II) and platinum(II) complexes of Schiff bases derived from 2,6-diacetylpyridine and S-methyldithiocarbazate and the X-ray crystal structure of the [Pd(mdapsme)Cl] complex", *Transition Metal Chemistry*, 2003; 28: 241–245. <https://doi.org/10.1023/a:1022960632103>
- [39]. P. Bera, R.J. Butcher, S. Chaudhuri, N. Saha, "Synthesis and spectroscopic characterisation of cobalt(III) complexes with S-benzyl-N-(5-methylpyrazole-3-yl)methylenedithiocarbazate (H<sub>2</sub>L): X-ray structures of [Co(HL)<sub>2</sub>]NO<sub>3</sub>·EtOH (1) and [Co(HL)(L)]·H<sub>2</sub>O (2)", *Polyhedron*, 2002; 21(1): 1–6. [https://doi.org/10.1016/S0277-5387\(01\)00933-0](https://doi.org/10.1016/S0277-5387(01)00933-0)
- [40]. M.A. Ali, M.T.H. Tarafder, M.A. Khair, "Synthesis, characterization and biological activities of some new nickel(II), copper(II), zinc(II) and cadmium(II) complexes of quadridentate SNNS ligands", *Transition Metal Chemistry*, 1992; 17: 133-136. <https://doi.org/10.1007/BF02910804>
- [41]. M.E. Hossain, M.N. Alam, J. Begum, M.A. Ali, M. Nazimuddin, F.E. Smith, R.C. Hynes, "The preparation, characterization, crystal structure and biological activities of some copper(II) complexes of the 2-benzoylpyridine Schiff bases of S-methyl- and S-benzyl-dithiocarbazate", *Inorganic Chimica Acta*, 1996; 249(2): 207-213. [https://doi.org/10.1016/0020-1693\(96\)05098-0](https://doi.org/10.1016/0020-1693(96)05098-0)
- [42]. M.K.E. Zahan, M.S. Islam, M.A. Bashar, "Synthesis, characteristics, and antimicrobial activity of some complexes of Mn(II), Fe(III) Co(II), Ni(II), Cu(II), and Sb(III) containing bidentate Schiff base of SMDTC" *Russian Journal of General Chemistry*, 2015; 85(3): 667–672. <https://doi.org/10.1134/S1070363215030238>
- [43]. M.L. Low, L. Maigre, P. Dorlet, R. Guillot, J. Pagès, K.A. Crouse, C. Policar, N. Delsuc, "Conjugation of a new series of dithiocarbazate Schiff base copper(II) complexes with vectors selected to enhance antibacterial activity", *Bioconjugate Chemistry*, 2014; 25(12): 2269–2284. <https://doi.org/10.1021/bc5004907>
- [44]. M.E. Hossain, M.N. Alam, M.A. Ali, M. Nazimuddin, F.E. Smith, R.C. Hynes, "The synthesis, characterization and bioactivities of some copper(II) complexes of the 2-acetylpyridine Schiff bases of s-methyl- and s-benzyl-dithiocarbazate, and the x-ray crystal structure of the nitrato(s-benzyl-β-n-(2-acetylpyridyl) methylenedithiocarbazato)copper(II) complex", *Polyhedron*, 1996; 15(5-6): 973-980. [https://doi.org/10.1016/0277-5387\(95\)00310-X](https://doi.org/10.1016/0277-5387(95)00310-X)
- [45]. N. Amiram, U. Othman, "Synthesis and characterization of Co(II), Cu(II), Cd(II), Zn(II) and Ni(II) complexes of schiff base ligand derived from S-benzyl-dithiocarbazate (SBDTC) and acetophenone with their biological activity studies", *IOSR Journal of Engineering (IOSRJEN)*, 2013; 3(8): 38-50. <https://doi.org/10.9790/3021-03853850>
- [46]. M.A.F.A. Manan, M.I.M. Tahir, K.A. Crouse, F.N.F. How, D.J. Watkin, "Synthesis, characterization and antibacterial activity of Schiff base derived from S-methyldithiocarbazate and methylisatin", *Journal of Chemical Crystallography*, 2012; 42:173–179. <https://doi.org/10.1007/s10870-011-0220-6>

- [47]. N.A. Bitu, S. Hossain, A. Kader, M.S. Islam, M.M. Haque, F. Hossen, A. Asraf, K.E. Zahan, "SBDTC derived Schiff base metal complexation: synthesis, physicochemical, and spectral studies with antibacterial activity", *Russian Journal of General Chemistry*, 2020; 90(8):1529–1532. <https://doi.org/10.1134/S1070363220080228>
- [48]. M.A. Ali, A.H. Mirza, A.M.S. Hossain, M. Nazimuddin, "Synthesis, characterization, antifungal properties and X-ray crystal structures of five- and six-coordinate copper(II) complexes of the 6-methyl-2-formylpyridine-4N-dimethylthiosemicarbazone", *Polyhedron*, 2001; 20(9-10): 1045–1052. [https://doi.org/10.1016/S0277-5387\(01\)00724-0](https://doi.org/10.1016/S0277-5387(01)00724-0)
- [49]. M.A. Ali, A.H. Mirza, M. Nazimuddin, H. Rahman, R.J. Butcher. "Mono- and bis-chelated nickel(II) complexes of the di-2-pyridylketone Schiff base of S-methyldithiocarbazate and the X-ray crystal structure of the bis[S-methyl-N-N-(di-2-pyridyl)-methylenedithiocarbazate]nickel(II) complex", *Polyhedron*, 2001; 20(19): 2431–2437. [https://doi.org/10.1016/S0277-5387\(01\)00764-1](https://doi.org/10.1016/S0277-5387(01)00764-1)
- [50]. M.T.H. Tarafder, M.A. Ali, D.J. Wee, K. Azahari, S.Silong, K.A. Crouse, "Complexes of a tridentate ONS Schiff base: synthesis and biological properties", *Transition Metal Chemistry*, 2000; 25: 456-460. <https://doi.org/10.1023/A:1007062409973>
- [51]. M.A. Ali, A.H. Mirza, M. Nazimuddin, P.K. Dhar, R.J. Butcher, "Preparation, characterization and antifungal properties of nickel(II) complexes of tridentate ONS ligands derived from N-methyl-S-methyldithiocarbazate and the X-ray crystal structure of the [Ni(ONMeS)CN]. H<sub>2</sub>O complex", *Transition Metal Chemistry*, 2002; 27: 27–33. <https://doi.org/10.1023/A:1013434113299>
- [52]. M.A. Ali, A.H. Mirza, M.S.H. Iran, "Magnetic, spectroscopic and biological properties of copper(II) complexes of the tridentate ligand, a-N-methyl-S-methyl-b-N-(2-pyridyl)methylenedithiocarbazate(NNS) and the X-ray crystal structure of the [Cu(NNS)I<sub>2</sub>] complex", *Transition Metal Chemistry*, 2002; 27: 262–267. <https://doi.org/10.1023/A:1014813315980>
- [53]. M.A. Ali, A.H. Mirza, "The preparation and characterization of mono- and bis-chelated cadmium(II) complexes of the di-2-pyridylketone Schiff base of S-methyldithiocarbazate (Hdpksme) and the X-ray crystal structure of the [Cd(dpksme)<sub>2</sub>].0.5MeOH complex", *Transition Metal Chemistry*, 2002; 27: 268–273. <https://doi.org/10.1023/A:1014894403685>
- [54]. M.L. Low, G. Paulus, P. Dorlet, R. Guillot, R. Rosli, N.Delsuc, K.A. Crouse, C. Policar, "Synthesis, characterization and biological activity of Cu(II), Zn(II) and Re(I) complexes derived from S-benzylthiocarbazate and 3-acetylcoumarin", *Biometals*, 2015; 28: 553–566. <https://doi.org/10.1007/s10534-015-9831-2>
- [55]. M.X. Li, L.Z. Zhang, C.L. Chen, J.Y. Niu, B.S. Ji, "Synthesis, crystal structures, and biological evaluation of Cu(II) and Zn(II) complexes of 2-benzoylpyridine Schiff bases derived from S-methyl- and S-phenyldithiocarbazates", *Journal of Inorganic Biochemistry*, 2012; 106(1):117-25. <https://doi.org/10.1016/j.jinorgbio.2011.09.034>

- [56]. M.L. Low, L. Maigre, M.I.M. Tahir, E.R.T. Tiekink, P. Dorlet, R. Guillot, T. Begum Ravoof, R. Rosli, J.Pagès, C. Policar, N.Delsuc, K.A. Crouse, “New insight into the structural, electrochemical and biological aspects of macrocyclicCu(II) complexes derived from S-substituted dithiocarbazateschiff bases”,*European Journal of Medicinal Chemistry*, 2016; 120: 1-12. <https://doi.org/10.1016/j.ejmech.2016.04.027>
- [57]. M.K.B. Break, T.Y. Fung, M.Z. Koh, W.Y. Ho, M.I.M. Tahir, O.A. Elfar, R.U. Syed, W.M.A. Khojali, T.M.Alluhaibi, B.Huwaimel, C. Wiart,T. Khoo, “Synthesis, crystal structure, antibacterial and in vitro anticancer activity of novel macrocyclicschiff bases and their Cu (II) Complexes derived from S-methyl and S-benzyl dithiocarbazate”,*Molecules*,2023; 28(13):5009. <https://doi.org/10.3390/molecules28135009>
- [58]. E.N.M. Yusof, M. Azam, S.S. Sirat, T.B.S.A. Ravoof, A.J. Page, A. Veerakumarasivam, T. Karunakaran,M.R. Razali, “Dithiocarbazateligand-based Cu(II), Ni(II), and Zn(II) complexes: synthesis, structural investigations, cytotoxicity, DNA binding, and molecular docking studies”,*Bioinorganic Chemistry and Applications*, 2022; 13. <https://doi.org/10.1155/2022/2004052>
- [59]. M. Das, S.E. Livingstone, “Metal chelates of dithiocarhazic acid and its derivatives. IX. metal chelates of ten new Schiff bases derived from S-methyldithiocarhazate”,*Inorganica Chimica Acta*, 1976; 19: 5-10. [https://doi.org/10.1016/S0020-1693\(00\)91065-X](https://doi.org/10.1016/S0020-1693(00)91065-X)
- [60]. C.L. Chena, X.F. Zhub, M.X. Lia, H.M. Guoc, J.Y. Niu,“Antitumor activity of manganese(II) and cobalt(III) complexes of 2-acetylpyridine Schiff bases derived from s-methyldithiocarbazate: Synthesis, characterization, and crystal structure of the manganese(II) complex of 2-acetylpyridine s-methyldithiocarbazate”,*Russian Journal of Coordination Chemistry*, 2011; 37(6): 435–438. <https://doi.org/10.1134/S1070328411050034>
- [61]. A.E. Libera, D.X. West,“Antifungal and antitumor activity of heterocyclic thiosemicarbazones and their metal complexes: current status”,*Biometals*, 1992; 5(2): 121-126. <https://doi.org/10.1007/BF01062223>
- [62]. M.A. Ali, A.H. Mirza, R.J. Butcher, M.T.H. Tarafder, “Biological activity of palladium(II) and platinum(II) complexes of the acetone Schiff bases of S-methyl- and S-benzyl dithiocarbazate and the X-ray crystal structure of the [Pd(asme)<sub>2</sub>] (asme=anionic form of the acetone Schiff base of S-methyldithiocarbazate) complex”,*Journal of Inorganic Biochemistry*, 2002; 92(3-4): 141–148. [https://doi.org/10.1016/s0162-0134\(02\)00559-7](https://doi.org/10.1016/s0162-0134(02)00559-7)
- [63]. L.Z. Zhang, T. Ding, C.L. Chen, M.X. Lia, D. Zhang, J.Y. Niu, “Biological activities of pyridine 2-carbaldehyde Schiff bases derived from S-methyl and S-benzyl dithiocarbazate and their zinc(II) and manganese(II) complexes. Crystal structure of the manganese(II) complex of pyridine 2-carbaldehyde S-benzyl dithiocarbazate”,*Russian Journal of Coordination Chemistry*, 2011; 37(5): 356–361. <https://doi.org/10.1134/S1070328411040117>
- [64]. S.A. Elsayed, A.M. Noufal, A.M.E. Hendawy, “Synthesis, structural characterization and antioxidant activity of some vanadium(IV), Mo(VI)/(IV) and Ru(II) complexes of

- pyridoxal Schiff base derivatives”, *Journal of Molecular Structure*, 2017; 1144: 120-128. <https://doi.org/10.1016/j.molstruc.2017.05.020>
- [65]. M.T.H. Tarafder, K. Chew, K.A. Crouse, A.M. Ali , B.M. Yamin , H.K. Fun, “Synthesis and characterization of Cu(II), Ni(II) and Zn(II) metal complexes of bidentate NS isomeric Schiff bases derived from S-methyldithiocarbazate (SMDTC): bioactivity of the bidentate NS isomeric Schiff bases, some of their Cu(II), Ni(II) and Zn(II) complexes and the X-ray structure of the bis[S-methyl-b-N-(2-furylmethyl)methylenedithiocarbazato]zinc(II) complex”, *Polyhedron*. 2002; 21: 2683-2690. [https://doi.org/10.1016/S0277-5387\(02\)01285-8](https://doi.org/10.1016/S0277-5387(02)01285-8)
- [66]. M.T.H. Tarafder, N. Saravanan, “Coordination Chemistry and biological activity of Ni(II) and Cu(II) ion complexes with nitrogen-sulfur donor ligands derived from S-benzylidithiocarbazate(SBDTC)”, *Transition metal chemistry*, 2001; 26: 613-618. <https://doi.org/10.1023/A:1012047001167>

UNDER PEER REVIEW