

Original Research Article

Enhancement of Drought Tolerance and Plant Growth in Tomato by *Trichoderma harzianum* Seed Treatment

ABSTRACT:

The proficiencies of soil microorganisms to drought tolerance have been known for many years, but their use as crop inoculants was not explored in agriculture commercially. One possibility to increase the drought tolerance is to use beneficial microorganisms as seed treatment. This study was conducted to investigate the effect of *Trichoderma harzianum* on the growth and development of Tomato (*Solanum lycopersicum* L.) plant under water stress condition. *Trichoderma harzianum* is applied to the seeds by treating seeds with *Trichoderma harzianum* before sowing. Parameters including germination percentage, chlorophyll content, leaf area, no of leaves, plant height, root length, plant fresh and dry weight and yield were recorded in all the treatments along with the untreated control. Tomato seeds treated with *Trichoderma harzianum* had higher seed germination percentage and chlorophyll content 95.50%, (2.41%), 42.59 SAPAD units (13.83%) respectively than untreated seed. Maximum leaf area and no of leaves 11.71cm² (19.40%), 190.75 (15.43%) respectively recorded in the plants of *Trichoderma harzianum* treated seeds in water stress condition as compared to untreated seeds. Under extreme water stress condition plant height and root length of tomato plant increased significantly 98.25cm, (19.45%), 41.25cm (42.24%) respectively in *Trichoderma harzianum* treated seeds as compare to untreated seed. Also, *Trichoderma harzianum* seed treatment in tomato showed a significantly higher plant fresh and dry weight 189.43g (8.65%) and 31.98g (47.57%) respectively in drought condition as compared to untreated seeds. *Trichoderma harzianum* seed treatment played critical role in drought condition by increasing yield 731.25 g/plant (27.62%) increase over untreated seeds. Taken together, the study recommends that in drought condition for enhanced tomato growth *Trichoderma harzianum* should be used as eco-friendly solutions by seed treatment.

Keywords: Drought tolerance, growth parameters, Tomato, Seed treatment, Root and shoot length, Plant fresh and dry weight, *Trichoderma herzianum*, Chlorophyll content and yield.

1. INTRODUCTION:

Climate change is a burning issue, and the incidence of global warming-induced by catastrophic climatic events is on the rise, causing global ecosystems including agroecosystems to suffer (Ebert and Engels, 2020). The occurrence of the abiotic stresses like drought results in a considerable reduction in the yield of crop plants throughout the globe (Khan et al., 2021). Concurrently, drought is becoming a dominant [environmental stressor for crop production globally](#) [global environmental stressor for crop production](#), limiting food security with precarious economic and sociological impacts (Trenberth et al., 2014). Due to climate change, water deficit is spreading to regions where drought was infrequent in the past (Somerville and Briscoe, 2001). Globally, the reduction in yield due to drought is likely to exceed the combined loss of all other possible causes of yield decline (Blum, 1996; Foolad, 2007). Therefore, there is a need to identify solutions to mitigate water deficit stress and its impact on food security.

The cultivation of tomato (*Solanum lycopersicum* L.) more than 7000 years ago in South America. Where small tomato fruits were first discovered and consumed orally and therapeutically. Tomatoes later spread to countries such as Peru, the Netherlands, and the United Kingdom, and are now grown and consumed in all parts of the world. There are many varieties of tomatoes and it is one of the most widely consumed vegetables in the world. Tomato is an annual plant and it is one of the most commercially important vegetable. Today, tomato is the second most consumed and widely grown nonstarchy vegetable in the world after potato. The average world production of tomato is estimated to be around 159 million tons. In the last few years, tomato consumption has increased considerably as they became an important part of the human diet. Therefore, the area of tomato cultivation has also expanded globally in the last few years. The production data shows that tomato production was 206.2 lakh tons in year 2022-23 (Spiralling Tomato Prices: Issues and Concerns, Issue No.: 03, August 2023, NABARD.).

Commented [B1]: Number

Formatted: Font: Italic

Commented [B2]: Read and deliver the real information please.

New and novel solutions for plant growth enhancements are required to ease the burden imposed on our environment and other resources. The relationship between plants and soil microbial communities is crucial to plant health (Oldroyd, 2013). Here, we look at potential solutions to these issues by investigating some of the research conducted regarding the biological inoculations of free-living plant growth-promoting microorganisms. The genus *Trichoderma*, belonging to the Hypocreaceae family (Hypocreales, Sordariomycetes, Ascomycota), is a highly studied group of filamentous fungi known for its numerous beneficial traits, particularly in agriculture as key components of microbiological inoculants. According to the most recent taxonomic concept, more than 400 described species of the genus are divided into 8 main clades (Cai and Druzhinina 2021). The beneficial effects of *Trichoderma*, however, are more pronounced when plants are under stress compared to when grown under optimal conditions (Mastouri et al., 2010; Lombardi et al., 2018).

The purpose of the present study was to investigate the effects of *Trichoderma harzianum* seed treatment in drought condition to enhance and promote yield and vegetative growth through the analysis of some indicating factors, such as germination percentage, chlorophyll content, leaf area, number of leaves, plant height, root length, plant fresh and dry weight and yield of tomato (*Solanum lycopersicum* L.) plant.

2 MATERIAL AND METHODS

2.1 Experimental site and Materials

Investigation was carried out in greenhouse during 2023-2024 at Umergaon, District Valsad, Gujarat in tomato (*Solanum lycopersicum* L.) crop. The Namdhari-NH-585th indeterminate early maturing variety of Namdhari seeds was selected for carrying present experiment.

Trichoderma harzianum (WP)-61, trade name Neemoderma-H of Shri Ram Solvent Extract Extractions (P) Ltd. having *Trichoderma harzianum* content 1% w/w (CFU Count 2×10^6 /gm. Min) plus carboxy methyl cellulose 1.0% w/w and talc powder 98.0% w/w was used for *Trichoderma harzianum* seed treatment.

Surface soil (0-15 cm) collected from farmer's field and mixed thoroughly after air dried and passed through 2 mm sieve. Well decomposed FYM is mixed in collected surface soil (10:1 ratio) and autoclaved at 15 PSI pressure and 121°C temperature for 30 minutes to sterilize. Autoclaved soil is filled in 12 inch plastic pots at the rate of 6.3 kg autoclaved soil in each pot and pots are arranged in greenhouse.

2.2 Seed treatment with *Trichoderma harzianum* and seed sowing

Tomato seeds are surface sterilized with 1% sodium hypochlorite for three minutes. Then treated seeds are washed with sterilized water and dried. The dried 30 seeds of tomato are treated with 1g *Trichoderma harzianum* in Petri plate (Fig 1.). For untreated control, seeds of tomato are soaked in distilled water. Seed treatment is given to tomato seeds as per treatment details (Table 1.). Single seed sown in each pot and pots are arranged in greenhouse. 500 ml water is applied to each pot at every alternate day with the help of measuring beaker till 15 days. 28°C to 32°C temperature and 55% to 60% relative humidity is maintained in respective greenhouse throughout the crop lifecycle (Fig 2.).

Table 1. Treatment details of seed treatment

Treatment No.	Treatment Details
T1	T. harzianum + Irrigated
T2	T. harzianum + Water Stress
T3	Untreated seed + Irrigated
T4	Untreated seed + Water Stress (Control)

Formatted Table




Fig.1 Seed treatment with *Trichoderma harzianum* Fig. 2 Overview of experiment in Greenhouse

2.3 Seed Germination Percentage

Seed germination percentage was calculated at 15 days after sowing. Seed germination percentage was calculated by using the following formula (ISTA, 2010):

Germination percentage = (Number of germinated seeds/Total number of seeds sown) X100.

2.4 Total Chlorophyll Content

Chlorophyll content was measured with the help of portable chlorophyll meter SPAD (Soil Plant Analytical Development) chlorophyll meter (MinoltaTM) at 90 days after sowing. It was used to acquire rapid estimation of chlorophyll content in SPAD units (Lah *et al.* (2011). The measurement was done at morning 10 am. It helps to avoid droplets content on leaf surface.

2.5 Leaf Area and number of Leaves

In tomato crop total number (no) of leaves are counted after five picking/harvesting. Total leaf area was measured by using methods as given below:

Leaf area = L x W x F

where L = Maximum length (cm); W = Maximum width (cm); F = Factor (0.50)

2.6 Plant Height and Root Length

The plant height and root length were measured after five picking from each pot in each replication with the help of measuring scale.

2.7 Plant Fresh Weight

At the time of harvesting plants of tomato from each treatment were softly uprooted from pots and washed off soils from the roots by running water very carefully. Plants were dried by blotting paper for removing surface moisture. Then, shoots were cut at soil line and separated. Fresh weight of root and shoot were measured on electronic balance up to three decimal digits.

2.8 Plant Dry Weight

Dry weight of tomato was measured at harvesting stage. Plants from each treatment were softly uprooted from pots and washed off soils from the roots by running water very carefully. Plants were dried by blotting paper for removing surface moisture. Then the plants were taken in brown envelop and dried in an oven set to low heat (100°F) overnight. After that, plants were cooled in a dry environment. Then, shoots were cut at soil line and separated. Shoots and root of each plant separately weighed and plant dry weight recorded.

2.9 Yield

The matured tomatoes are harvested and weight of tomato was recorded on electronic digital balance. Yield of five subsequent picking is considered in tomato for yield calculation.

2.10 Statistical Analysis

The data generated from the pot culture experiment was analyzed by completely randomized design. The data obtained was statistically analyzed and appropriately interpreted as per the methods described in "Statistical Methods for Agricultural Workers" by Panse and Sukhatme, (1985). Appropriate standard error (S.E.) and critical differences (C.D.) at 5% level were worked out as and when necessary and used for data interpretation. The data were statistically

Commented [B3]: ???

3 Results and Discussion

3.1 Seed Germination Percentage

Trichoderma harzianum treated seed of tomato has 95.50 to 94.75 percent seed germination, which is statistically similar with other all treatments, however numerically higher than the untreated seed 93.75 to 93.25 percentage (Fig 3) (Table 2). Same results are noted by Donoso et al., (2008) that inoculation of tomato seed with *Trichoderma* improved germination and other growth parameters when grown at water deficit conditions. Shores et al., (2010) found that some species of this *Trichoderma* have direct effects on plants, increasing their potential for growth and nutrient absorption, fertilizer efficiency, a higher rate and percentage of seed germination, and stimulating plant defense against biotic and abiotic damage. Mastouri et al., (2009) observed similar results that *Trichoderma harzianum* added as seed treatment (tomatoes) or as a soil treatment (Arabidopsis) largely improved the germination at osmotic potentials of up to 0.3 MPa. Plants grown from these *Trichoderma* treatments are much more resistant to water deficit conditions.




Fig 3. Effect of seed treatment with *Trichoderma harzianum* on Seed Germination




Fig 4. Effect of seed treatment with *Trichoderma harzianum* chlorophyll content

Commented [B4]: Informations in horizontal axis in the figures are not clearly readable

3.2 Total Chlorophyll Content

Drought reduces the chlorophyll content severely in plant. We recorded significant results of *Trichoderma harzianum* seed treatment, the maximum Chlorophyll Content was recorded in the treatment T2 i.e. seed treatment of *Trichoderma harzianum* with water stress condition, which gave (42.59 SPADA units) 22.02% increase over control followed by the T1 treatment plants i.e. seed treatment with *Trichoderma harzianum* with regular irrigation (40.77 SPADA units) 16.80% increase over control (Fig 4) (Table 2). The T3 treatment untreated seed with regular irrigation has (37.54 SPADA units) 7.54 % increase over control i.e. T4 treatment untreated seed with water stress condition control has (34.91 SPADA units) Chlorophyll Content. Shukla et al., (2012) observed that *Trichoderma harzianum* inoculated plants exhibit delayed wilting, increased stomatal conductance, enhanced leaf chlorophyll content, and greater net photosynthesis levels under water deficit stress conditions. Contreras-Cornejo et al., (2019) found that the plants inoculated with *Trichoderma* isolates can enhance water-deficit tolerance by improving root development, regulation of drought-induced changes in the stomatal opening, photosynthesis, and chlorophyll content in leaf.

Commented [B5]: Something is missing here. Please insert it or re-read the sentence for proper correction

3.3 Leaf Area and Number of Leaves

Significant variation of leaf area and no of leaves observed with different treatments in tomato plant (Fig 4) (Table 2). Treatment T2 i.e. seed treatment with *Trichoderma harzianum* having

water stress condition has found maximum leaf area 19.40% (11.71 cm²) increase over control followed by T1 *Trichoderma harzianum* with mean leaf area of 11.12 cm² increasing leaf area by 13.38% over control. Untreated plants with regular irrigation i.e. T3 have 10.50 cm² leaf area which is a 7.01% increase over control. The control i.e. T1 untreated seed with water stress has 9.81 cm² leaf area. Rabeendran *et al.*, (2000) carried out experiment for the evaluation of *Trichoderma* isolates in a glasshouse to check the growth promotion effects of *Trichoderma* on cabbage seedlings (*Brassica oleracea* L.) and reported that increased (P<0.05) leaf area (58.71%), shoot dry weight (91.102%) and root dry weight (100.158%) compared with the untreated control.

Among all the treatments, T2 treatment i.e. seed treatment with having water stress condition shows higher number of leaves 190.75 No of leaves per plant (190.75), which is 15.43% over control followed by treatment T2 having *Trichoderma harzianum* seed treatment with regular irrigation recording 189.75 No of leaves per plant and 14.83% increase over control (Fig 5) (Table 2). The treatment T3, i.e. untreated seed with regular irrigation have 185.25 leaves per plant having 12.10% increase over control. The control i.e. untreated seed with water stress condition (control) has 165.25 No of leaves per plant. Sundaramoorthy and Balabaskar (2013) recorded similar results that there was an increase in the plant height, number of leaves, and total yield of tomatoes in plants treated with *T. harzianum*.




Fig 5. Effect of seed treatment with *Trichoderm harzianum* on leaf area




Fig 6. Effect of seed treatment with *Trichoderm harzianum* on number of leaves

3.4 Plant Height and Root Length

In present study, we observed that drought reduces the growth drastically and *Trichoderma harzianum* not only improved these negative effects but also enhanced the plant growth significantly by increasing plant shoot and root growth (Fig 7 and 8) (Table 2.). In this experiment furthermore, we observed that the tomato plants of treatment T2 i.e. *Trichoderma harzianum* seed treatment with water stress condition has 98.25 cm plant height which is 19.45% increase over control followed by treatment T1 having *Trichoderma harzianum* with irrigation condition with tomato plant height of 95.75 cm plant height, which is 16.41% increase over control. The Treatment T3 i.e. untreated seed with regular irrigation has 92.75 cm plant height and 12.77% increase over control. The control i.e. untreated seed with water stress condition has plant has 82.25 cm plant height. Kucuk (2013) conducted experiment on wheat and reported that seed treatment with *Trichoderma harzianum* will help to increase the considerable plant height.




Fig 7. Effect of seed treatment with *Trichoderma harzianum* on Plant height



Fig 8. Effect of seed treatment with *Trichoderma harzianum* on plant height

Commented [B6]: It would be better to put the meaning of T1 to T4 after each legend (figure or table) than to repeat it each time throughout the text.

Significant improvement in root length (41.25 cm) was achieved in the plants of treatment T2 i.e. seed treatment with *Trichoderma harzianum* having water stress condition of 41.25 cm with 42.24% increase over control. The treatment T2 is followed by treatment T1 i.e. seed treatment with *Trichoderma harzianum* and regular irrigation with increase of root length by 29.31% (37.50cm) over control (Fig 9 and 10) (Table 2). Treatment T3 having untreated seed with regular watering enhancing enhanced the root length by 20.69% (35 cm) over control. The control i.e. treatment T4 having untreated seed and water stress condition has recorded 29 cm root length. Cai *et al.* (2013) reported that harzianolide produced by *Trichoderma* spp. can improve the early stage of plant development through the enhancement of root length. These morphological modifications are possible because of the ability of the *Trichoderma* spp. to act through several mechanisms such as environmental buffering (against pH, drought, waterlogging, cold and heat), phosphorus solubilization, organic matter decomposition, chitinase and siderophore production.




Fig 9. Effect of seed treatment with *Trichoderma harzianum* on Root length



Fig 10. Effect of seed treatment with *Trichoderma harzianum* on Root length

3.5 Plant Fresh Weight

The data presented in table 2 indicate that the T2 treatment i.e. seed treatment with *Trichoderma harzianum* and drought growing condition has greater plant fresh weight 189.43 g, which is 8.65% increase over control followed by T1 treatment i.e. seed treatment with *Trichoderma* and regular irrigation with 182.17g and 4.49% increase over control (fig 11) (table 2). The T3 treatment having untreated seed with regular irrigation has recorded 178.26g of plant fresh weight, which is 2.245 increase over control and control i.e. treatment T4 untreated seed with water stress condition has 174.35 g plant fresh weight, which is comparatively less than other all treatment. Chacón *et al.* (2007) found that inoculation of *T. harzianum* increased fresh weight and lateral roots in tomato plants. *Trichoderma* species are present in all soil types and are the most common cultivable fungi.




Fig 11. Effect of seed treatment with *Trichoderma harzianum* on Plant Fresh weight




Fig 12. Effect of seed treatment with *Trichoderma harzianum* on Plant Dry weight

3.6 Plant Dry Weight

As like plant fresh weight observations, we noticed that the T2 treatment i.e. seed treatment with *Trichoderma harzianum* with water stress condition has a maximum plant dry weight of 31.98 g, which is 47.57% increase over control followed by treatment T1 with i.e. *Trichoderma harzianum* seed treatment with regular irrigation 29.99 g having 34.21% increase over control (Fig 12) (Table 2). The treatment T3 having untreated seed and regular irrigation has 26.88 g of plant dry weight which is 24.03% increase over control. In control i.e. T4 treatment i.e. T4 untreated seed and water stress condition we recorded the plant dry weight record is 21.67g plant fresh weight. Camargo *et al.* (2014) reported that the application of commercial *Trichoderma* sp. in the cultivation of pea *Pisum sativum* L. (Fabaceae) significantly improved its growth and development, influencing physiological variables such as germination, leaf area, dry weight of the root and; fresh weight of the root, dry weight of the aerial part and; fresh weight of the aerial part and length of root, favoring the productive yield of the crop, when applying the same treatment to the seed.

Commented [B7]: Fresh or dry?

3.7 Yield

Results indicate that *Trichoderma harzianum* seed treatment significantly increased the yield even in drought condition. The treatment T2 exhibiting having *T. harzianum* seed treatment with water stress condition has maximum yield of 731g.25g, which is 27.62% increase over control followed by T1 treatment i.e. *Trichoderma harzianum* seed treatment with regular irrigation with 727.75g and 27.01% increase over control (Fig 13 and 14) (Table 2.). With T the treatment T3 i.e. untreated seed with regular irrigation has the yield recorded is 635.25g yield and with 10.86% increase over control. The control treatment T4 untreated seed with water stress condition has 573g yield. Mishra *et al.* (2020) found that *Trichoderma* spp. seed treatment at sowing will boosts chlorophyll accumulation, root and shoot growth, tiller number, and crop yield in rice even under drought conditions. Scudeletti *et al.* (2021) conducted field experiment in sugarcane crop and reported that *Trichoderma* spp. inoculation of sugarcane improves plant morphology and physiological factors under drought stress which contribute in increasing sugarcane yield





Fig 13. Effect of seed treatment with *Trichoderma harzianum* on Yield

Fig 14. Effect of seed treatment with *Trichoderma harzianum* on Yield

Table 2. Effect of *Trichoderma harzianum* on Germination %, Chlorophyll content, Leaf area, No of leaf/plant, Plant height, Root length, Frsh and Dry weight and Yield.

Trt . #	Treatment Details	Germination %	Chlorophyll content (SPADA units)	Leaf area (cm)	No of leaves (Nos.)	Plant height (cm)	Root Length (cm)	Fresh weight (g)	Dry Weight (g)	Yield (g)
T1	<i>T. harzianum</i> +Irrigated	94.75	40.77	11.12	185.25	95.75	37.50	182.17	29.09	727.75
T2	<i>T. harzianum</i> +Water Stress	95.50	42.59	11.71	190.75	98.25	41.25	189.43	31.98	731.25
T3	Untreated seed +Irrigated	93.75	37.54	10.50	189.75	92.75	35.00	178.26	26.88	635.25
T4	Untreated seed +Water Stress (Control)	93.25	34.91	9.81	165.25	82.25	29.00	174.35	21.67	573.00
	S.Em.±	1.64	0.53	0.49	1.65	3.21	1.63	2.83	0.97	7.59
	C.D. at 5 %	NS	1.69	1.56	5.26	10.28	5.23	9.07	3.10	24.28
	C.V. %	3.48	2.72	2.46	1.80	6.96	9.15	3.13	7.08	2.46

Formatted Table

4 Conclusion

In farming, the novel and conventional inventions boost up the yield of agriculture produce. The challenge faced by modern agriculture is to accomplish high yields in an environment-friendly manner. Hence, quick action on finding eco-friendly solutions needs to be done.

It is worldwide well known that *Trichoderma harzianum* is used as a biocontrol agent against different plant pathogens. In Existing study results indicates indicate that although seed treatment with *Trichoderma harzianum* provides an innovative, cost-effective, low toxicity, and environmentally friendly means of increasing crop yields in tomato through improving seed germination, chlorophyll content and growth under abiotic stress like drought stress conditions.

5 References

- A.W. Ebert, J.M.M. Engels 2020. Plant biodiversity and genetic resources matter. *Plants* 9(12), 1706
- Blum A. 1996. Crop responses to drought and the interpretation of adaptation. *Plant Growth Regul.* 20:135-148.
- C Somerville, J Briscoe. 2001. Genetic engineering and water. *Science*, 292, Article 2217.
- Cai F, Druzhinina IS 2021. In honor of John Bissett: authoritative guidelines on molecular identification of *Trichoderma*. *Fungal Diversity* 107: 1-69.
- Cai F, Yu G, Wang P, Wei Z, Fu L, Shen Q, Chen W. Harzianolide, 2013. a novel plant growth regulator and systemic resistance elicitor from *Trichoderma harzianum*. *Plant Physiology and Biochemistry*; 73:106-113
- Camargo, C., David F. and ávila, E.R. 2014. Efectos del *Trichoderma* sp. sobre el crecimiento y desarrollo de la arveja (*Pisum sativum* L.). *Ciencia y Agricultura*, 11, 91-100.

- Chacón S., M.I. et al. (2007)**, "Phylogeographic analysis of the chloroplast DNA variation in wild common bean (*Phaseolus vulgaris* L.) in the Americas", *Plant Systematics and Evolution*, Vol. 266, No. 3, pp. 175-195
- Contreras-Cornejo, H. A., Macías-Rodríguez, L., Cortés-Penagos, C., and López-Bucio, J. 2009.** *Trichoderma virens*, a Plant Beneficial Fungus, Enhances Biomass Production and Promotes Lateral Root Growth through an Auxin-Dependent Mechanism in Arabidopsis. *Plant Physiol.* 149:1579–1592. doi: 10.1104/pp.108.130369
- Donoso, E., Lobos, G. A., and Rojas, N. 2008.** Efecto de *Trichoderma harzianum* y compost sobre el crecimiento de plántulas de *Pinus radiata* en vivero. *Bosque Valdivia.* 29:52–57.
- Foolad MR 2007.** Genome mapping and molecular breeding of tomato. *International Journal of Plant Genomics* 2007: 1–52. DOI 10.1155/2007/64358
- Kucuk, C. 2013.** Enhanced root and shoot growth of wheat (*Triticum aestivum* L.) by *Trichoderma harzianum* from Turkey *Pak. J. Biol. Sci.* 17 122 125
- Lah MC, Nordin MNB Isa MBM, Khanif Y, Jahan MS 2011.** Composting increases BRIS soil health and sustains rice production. *Science Asia;* 37(4):291-295.
- Lombardi N., Vitale S., Turrà D., Reverberi M., Fanelli C., Vinale F., et al. 2018.** Root exudates of stressed plants stimulate and attract trichoderma soil fungi. *Mol. Plant Microbe Interact.* 31 982–994. 10.1094/MPMI-12-17-0310-R
- M.A. Khan, D. Pattnaik, R. Ashraf, I. Ali, S. Kumar, N. Donthu 2021.** Value of special issues in the Journal of Business Research: A bibliometric analysis *Journal of Business Research*, 125 (2021), pp. 295-313
- Mastouri F., Björkman T., Harman G. E. 2010.** Seed treatment with *Trichoderma harzianum* alleviates biotic, abiotic, and physiological stresses in germinating seeds and seedlings. *Phytopathology* 100 1213–1221. 10.1094/PHYTO-03-10-0091
- Mastouri F., Harman G. E. 2009.** Beneficial microorganism *Trichoderma harzianum* induces tolerance to multiple environmental and physiological stresses during germination in seeds and seedlings. In IS-MPMI 2009 XIV Congress. Quebec, Canada
- Mishra, D., Rajput, R. S., Zaidi, N. W., Singh, H. 2020.** Sheath blight and drought stress management in rice (*Oryza sativa*) through *Trichoderma* spp. *Indian Phytopathology*, 73(1), 71–77.
- Oldroyd, G.E. 2013.** Speak, friend, and enter: signalling systems that promote beneficial symbiotic associations in plants. *Nat. Rev. Microbiol.* 11: 252–263.
- Panse, V.G. and Sukhatme, P.V. 1985.** *Statistical Methods for Agricultural Workers.* Indian Council of Agricultural Research Publication, 87-89.
- Rabeendran, N., Moot, D.J., Jones, E.E., Stewart, A., 2000.** Inconsistent growth promotion of cabbage and lettuce from *Trichoderma* isolates. *N. Z. Plant Prot.* 53, 143–146.
- Scudeletti, D., Crusciol, C. A. C., Bossolani, J. W., Moretti, L. G., Momesso, L. et al. 2021.** *Trichoderma asperellum* inoculation as a tool for attenuating drought stress in sugarcane. *Frontiers in Plant Science*, 12, 645542
- Shoresh, M., Harman, G. and Mastouri, F. 2010.** Induced Systemic Resistance and Plant Responses to Fungal Biocontrol Agents. *Annual Review of Phytopathology*, 48, 21-43.
- Shukla, N., Awasthi, R. P., Rawat, L., and Kumar, J. 2012.** Biochemical and physiological responses of rice (*Oryza sativa* L.) as influenced by *Trichoderma harzianum* under drought stress. *Plant Physiol. Biochem.* 54, 78–88. doi: 10.1016/j.plaphy.2012.02.001

Sundaramoorthy, S., Balabaskar, P., 2013. Biocontrol efficacy of *Trichoderma* spp. against wilt of tomato caused by *Fusarium oxysporum* f. sp. *lycopersici*. J. Appl. Biol. Biotechnol., 1, Pp. 36–40.

Trenberth, K. E., Dai, A., van der Schrier, G., Jones, P. D., Barichivich, J., Briffa, K. R., et al. 2014. Global warming and changes in drought. Nat. Clim. Change 4, 17–22.

UNDER PEER REVIEW