

Genetic Variability Studies for Morpho-Physiological and Yield Attributing Traits in Foxtail Millet (*Setaria italica* L. Beauv.)

Abstract - Foxtail millet is an underutilized crop which has high nutritional values and wide genetic resources, possessing the ability to tolerate several abiotic stresses efficiently. Those resources must be used in order to improve desired traits, primarily yield attributes. ~~An~~ This investigation was carried out to assess the genetic parameters, such as variability, *per se* performance, heritability and genetic advance as ~~per cent~~ percent over mean for sixteen morpho-physiological, yield attributing traits in fifty foxtail millet germplasm accessions at the Regional Agricultural Research Station, Nandyal during Rabi, 2023-2024. The study revealed the wide range of variation for all ~~the studied~~ traits and that the accessions SiA 3290, SiA 4345, and SiA 4391 had potential for most traits and a high grain yield ~~per plant~~. Thus, after multilocation testing, these accessions might be exploited for commercial cultivation. The assessment revealed that the phenotypic coefficient of variation (PCV) was slightly greater than the corresponding genotypic coefficient of variation (GCV), indicating the influence of the environment on the manifestation of these traits. It was also observed that the traits *viz.*, photosynthetic rate, transpiration rate, stomatal conductance, abortive grain rate, and grain yield per plant had high PCV and GCV and these traits significantly contributed to the overall variability and scope for selection. High heritability in combination with high genetic advance as per cent of mean (GAM) was observed for photosynthetic rate, transpiration rate, stomatal conductance, 1000 grain weight, fodder yield per plant and grain yield per plant. This indicates that, these traits are mostly governed by additive genes and thus making them suitable for simple direct selection to improve these traits and further breeding efforts focused on increasing yields.

Keywords: Foxtail millet, ~~PCV, GCV, heritability, genetic variability, per se performance,~~ morpho-physiological traits and grain yield.

INTRODUCTION

Foxtail millet (*Setaria italica* L. Beauv.) cultivate around the world and also referred as Chinese millet, German millet, Italian millet and Hay millet. It is a C₄ crop with short life cycle and a relatively small genome (~515 Mb), belongs to the family Poaceae. The crop is extremely autogamous, with a 4% cross-pollination rate, and has a diploid ($2n = 2x = 18$) chromosomal number (Lata *et al.* 2013). Foxtail millet continues to play a significant role in global agriculture, feeding millions of people who rely on marginal or poor soils of Europe and Asia. Apart from human use, the grain is also utilised as feed for cattle and poultry and its stover as fodder for animals. Foxtail millet is widely known for its exceptional drought tolerance and is naturally equipped with excellent water use efficiency and nitrogen use efficiency. Further, numerous morpho-physiological traits, such as dense and deep root systems, smaller leaf area and thicker cell walls were thought to contribute to durable tolerance to a range of abiotic stresses, majorly for drought, heat, and salinity (Lata *et al.* 2013; Diao *et al.* 2014). Recently, foxtail millet has gained recognition as an important food for diabetics. It has low levels of fat (4%) and a low glycaemic index (GI = 57). Its bran is rich in linoleic and oleic acids. It is rich in dietary fiber (6.7%), minerals (4%) and micronutrients (Fe, Ca, P) and protein (11–12%) (Liang *et al.* 2010).

In general, foxtail millet productivity is low, because of local cultivars, which yield poorly, are cultivated on marginal land of arid regions. Which, necessitates the development of more stable, highly productive and adaptable foxtail millet cultivars. Genetic variability is

a key prerequisite for genetic improvement in plant breeding. Knowledge on the extent of variability existing in a crop species for different traits is crucial, because it serves as the basis for effective selection. To obtain a realistic indication of genetic variation in any trait, phenotypic variability must be partitioned into heritable and non-heritable components. The potential genetic gain from a selection process may be evaluated using estimations of heritability along with genetic advance. The aim of this research was to ~~ascertain-estimate~~ the degree of variation among different morpho-physiological and grain yield related traits. ~~Furthermore, to- and~~ identify the best suitable lines for increasing the grain yield of foxtail millet.

MATERIAL AND METHODS

Fifty foxtail millet germplasm accessions including three checks viz., SiA 3156, SiA 3223 and SiA 3159 ~~were~~ were evaluated in two replications using a ~~randomised- Randomised complete-Complete block-Block design-Design~~ (RCBD) at the Regional Agricultural Research Station, Nandyal, Andhra Pradesh, during *Rabi* season, 2023-~~20~~24. The accessions were sown at a spacing of 22.5 cm × 10 cm. Recommended agronomic practices were ~~followed to raise a good crop applied at the prime time~~. Five competitive plants per accession were selected randomly for recording observation on plant height (cm), number of productive tillers per plant, flag leaf length (cm), flag leaf width (cm), panicle length (cm), photosynthetic rate ($\mu\text{mol CO}_2 \text{ m}^{-2}\text{s}^{-1}$), transpiration rate ($\text{mmol H}_2\text{O m}^{-2}\text{s}^{-1}$), stomatal conductance ($\text{mmol H}_2\text{O m}^{-2}\text{s}^{-1}$), relative water conductance (%), abortive grain rate, fodder yield per plant (g), harvest index (%) and grain yield per plant (g), while observations on days to 50% flowering and days to maturity were recorded on a plot basis. For 1000-grain weight, a random sample of 1000 grains were counted from the threshed seed and the weight was recorded in grams. The analysis of variance was estimated by Panse and Sukhatme (1967). Heritability in broad sense (h^2_b) was calculated and classified into low (below 30%), medium (30- 60%) and high (above 60%) and genetic advance as ~~per-cent percent~~ over mean (at 5%) were computed and categorized into low (0-10%), moderate (10-20%) and high ($\geq 20\%$) as given by Johnson *et al.* (1955). Estimates of PCV and GCV was calculated following Burton and De Vane (1953) and categorized into low ($<10\%$), moderate (10-20%) and high ($>20\%$) according to Sivasubramanian and Madhavamenon (1973).

RESULT and DISCUSSIONS

~~Analysis of variance for morpho-physiological and yield attributing traits in foxtail millet germplasm accessions are presented in Table 1. In the current study the analysis of variance (ANOVA) revealed highly significant differences among foxtail millet germplasm accessions for 16 traits (Table. 1), which Results showed that, mean squares were significant for all studied traits, indicates presence of sufficient variability and scope for further selection and breeding superior and desirable genotypes. Average mean performance of the top ten high yielding accessions for a range of morpho-physiological attributes were -are presented in Table. 2. The germplasm accessions viz., SiA 3290 (11.59 g/plant), SiA 4345 (11.35 g/plant) and SiA 4391 (11.32 g/plant) performed exceptionally well for the most of the yield attributes and had the best *per se* performance, when compared to that of best standard check SiA 3159 (11.04 g/plant). Therefore, after multilocation testing, these accessions can thus be utilized for commercial exploitation. . Estimates of range of variation and genetic variability parameters~~

Commented [DR1]: In the references, Burton and Dewane (1952) , correct

Formatted: Line spacing: single

Commented [DR2]: How highly significant and the significant calculated at 5% only

Commented [DR3]: Where the previous traits in Table 2 and why enough to explain GYPP only

for the 16 traits in 50 foxtail millet accessions are illustrated in Table 3. The mean, range of variation and the estimated genetic parameters such as phenotypic coefficient of variation (PCV), genotypic coefficient of variation (GCV), heritability in broad sense and genetic advance as per cent of mean are illustrated in Table. 3. In the current investigation, significant differences in mean values for all the traits were noticed. The traits grain yield per plant ranged from 4.44 to 11.59 (g); days to 50% flowering ranged from 48.00 to 60.50; days to maturity ranged from 77.00 to 94.00; plant height ranged from 109.50 to 155.70 (cm); number of productive tillers ranged from 1.50 to 3.20; flag leaf length at flowering ranged from 23.10 to 39.70 (cm); flag leaf width at flowering ranged from 1.42 to 2.86 (cm); panicle length ranged from 12.90 to 21.90 (cm); 1000 grain weight ranged from 1.90 to 3.35 (g); fodder yield per plant ranged from 6.58 to 17.48 (g); harvest index ranged from 26.44 to 42.06 (%); photosynthetic rate ranged from 13.47 to 45.97 ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$); transpiration rate ranged from 2.98 to 6.88 ($\text{mmol m}^{-2} \text{ s}^{-1}$); stomatal conductance ranged from 0.12 to 0.34 ($\text{mmol m}^{-2} \text{ s}^{-1}$); relative water content ranged from 71.82 to 90.77 (%) and abortive grain rate ranged from 0.06 to 0.25. The experimental material had a favourable mean performance and wide range of variability for most of the traits studied and these prospective combinations might be used as potential lines aiming at simultaneous improvement for grain yield and other yield attributing traits.

Genotypic coefficient of variation cannot provide a clear estimate of the genetic gain of the trait, that can be anticipated from phenotypic-based selection, unless the heritability is known (Burton, 1952). Heritability and genetic advance as percentage of mean are the two primary selection criteria considered in the crop improvement. Heritability estimates for the different traits under investigation in this study varied from 43.60 to 97.20 *per cent*, whereas the genetic advance as percentage of mean varied from 5.92 to 45.74 *per cent*. The trends of heritability and genetic advance as percentage of mean are presented in Figure. 2. High heritability was noticed for the traits *viz.*, photosynthetic rate (97.20%), 1000-grain weight (91.00%), relative water content (90.50%), harvest index (78.60%), transpiration rate (76.30%), grain yield per plant (75.90%), stomatal conductance (75.00%), days to maturity (73.90%), flag leaf length (71.50%), panicle length (70.60%), fodder yield per plant (61.90%) and abortive grain rate (60.00%). Whereas, high genetic advance as per cent mean was also reported for stomatal conductance (45.90%), photosynthetic rate (45.74%), abortive grain rate (38.27%), transpiration rate (35.54%), grain yield per plant (35.24%), 1000 grain weight (23.25%), and fodder yield per plant (21.06%).

The estimates of heritability and genetic advance together are more accurate in assessing the potential genetic gain through selection (Johnson *et al.* 1955). Therefore, in the present study heritability and genetic advance both were estimated to access the possible extent of genetic improvement among the different accessions studied. High heritability in conjunction with high genetic advance as per cent over mean (GAM) was observed for the traits photosynthetic rate (97.20% and 45.74%); transpiration rate (76.30% and 35.58%); stomatal conductance (75.00% and 45.90%); 1000 grain weight (91.00% and 23.52%) as obtained by Harish *et al.* (2022), fodder yield per plant (61.90% and 21.06%) similar to the report by Yadav *et al.* (2024) and grain yield per plant (75.90% and 35.24%) as published by Srilatha *et al.* (2020), Harish *et al.* and Lavanya. (2022). High heritability accompanied by moderate GAM was found for flag leaf length (71.50% and 18.84%) as reported by Pallavi *et al.* (2020), panicle length (70.60% and 19.30%) similar to the results shown by Johar *et al.*

Formatted: Font: Italic

(2015) and Shingane *et al.* (2016, 2017), relative water content (90.50% and 11.72%) as realised by Bheemesh *et al.* (2018) and harvest index (78.60% and 19.52%) which was alike the results of Yadav *et al.* (2024). High heritability coupled with high GAM indicates that the traits are governed majorly by additive genes and these traits *viz.*, photosynthetic rate, transpiration rate, stomatal conductance, 1000 grain weight, fodder yield per plant and grain yield per plant can be improved through simple selection. The traits with moderate to low heritability and GAM imply that the traits are influenced by both additive and non-additive genes and thus simple selection alone might not be rewarding for the improvement of these traits, hence, in this crop recurrent selection can be employed for improvement of these traits. The Low estimates of heritability and genetic advance over the percentage of mean that were observed for 1000 grain weight (g) indicates that Selection will be inefficient because of the substantial environmental influence on the trait.

CONCLUSIONS

India has a wealth of foxtail millet genetic resources available; these resources must be utilized for harnessing the benefits. Evaluation and characterisation of the germplasm accessions can provide a substantial quantity of information about variability, which is essentially needed to improve the yield-related attributes and other desired traits. In the present study, efforts are made to unravel the variability in the germplasm lines and an analysis of variance revealed that germplasm accessions differed significantly for all the traits, providing sufficient diversity and range to select suitable genotypes. Among the tested accessions, SiA 3290, SiA 4345 and SiA 4391 were found to have potential for most of the traits and to be promising for high grain yield. All the traits showed higher PCV than corresponding GCV implying the existence of environmental influence. The traits photosynthetic rate, transpiration rate and stomatal conductance were known to possess high PCV and GCV, including the formerly mentioned traits, 1000 grain weight, fodder yield per plant and grain yield per plant has shown high heritability coupled with high GAM indicating the predominance of additive gene action on them and thus they can be improved through simple selection. Days to 50% flowering, days to maturity, plant height and relative water content possessed low to moderate variability, heritability and GAM which means that these traits are governed mostly by non-additive genes and impact of the environment is more in the expression of the traits, hence, simple selection may not be rewarding for the improvement of these traits.

REFERENCES

- Amarnath, K., Prasad, A.V.S.D., Reddy, C.V.C.M and Sreenivasulu, K.N. 2018. Genetic variability in Indian foxtail millet genetic resources (*Setaria italica* (L.) Beauv). *Green Farming*. 9 (4): 612-614.
- Bheemesh, S. J., Subba Rao, M and Reddi Shekhar, M. 2018. Genetic divergence studies for yield and physiological traits in foxtail millet. *Bulletin of Environment, Pharmacology and Life Sciences*. 7: 88-94.
- Burton, G.W. 1952. Quantitative inheritance of grasses. In: Proceedings 6th International Grassland Congress. 1: 273-283.

- Burton, G.W. and Dewane, E.W. 1952. Estimating heritability in tall fescue (*Festuca arundanaceae*) from replicated clonal material. *Agronomy Journal*. 45: 478- 481.
- Diao, X., Schnable, J., Bennetzen, J. and Li, J. 2014. Initiation of *Setaria* as a model plant. *Frontiers of Agricultural Science and Engineering*. 1: 16-20.
- Harish and Roopa Lavanya, G. 2022. Genetic variability and character association studies in foxtail millet (*Setaria italica* L.) for grain yield characters. *International Journal of Environment and Climate Change*.12(11): 2929-2938.
- Johar, J.S. 2015. A study on the nature of gene action for yield and yield components in exotic genotypes of Italian millet (*Setaria italica* (L.) Beauv.). *International Journal of Agricultural Sciences*. 5 (4): 667-670.
- Johnson, H.W., Robinson, H.F. and Comstock, R.E. 1955. Estimates of genetic and environmental variability in soybeans. *Agronomy Journal*. 47: 314–318.
- Lata, C., Gupta, S. and Prasad, M. 2013. Foxtail millet: a model crop for genetic and genomic studies in bioenergy grasses. *Critical Reviews in Biotechnology*. 33: 328–343.
- Li, H., Meng, W.J. and Liu, T.M. 1935. Problems in the breeding of millet [*Setaria italica* (L.) Beauv.]. *Journal of American Society of Agronomy*. 27:426–438.
- Liang, S., Yang, G and Ma, Y. 2010. Chemical characteristics and fatty acid profile of foxtail millet bran oil. *Journal of the American Oil Chemists' Society*. 87:63-67.
- Pallavi, N.L., Venkatesh, R., Ram, B and Suresh, B.G. 2020. Studies on correlation and path coefficient analysis in foxtail millet [*Setaria italica* (L.) Beauv]. *International Journal of Chemical Studies*. 8: 1941-1946.
- Panase, V.G. and Sukhatme, P.V. 1967. Statistical methods for agriculture workers. Indian council of Agriculture, New Delhi.
- Shingane, S., Gomashe, S., Ganapathy, K.N and Patil, J.V. 2017. Genetic variability and association analysis for grain yield and nutritional quality in foxtail millet. *International Journal of Bio-resource and Stress management*. 7(6): 1239-1243.
- Sivasubramanian, S. and Madhavamenon, M.P. 1973. Genotypic and phenotypic variability in rice. *Madras Agricultural Journal*. 60(9-12): 1093-96.
- Srilatha, J., Haritha, T., Reddy, C.V.C.M and Ramesh, D. 2020. Genetic variability studies for yield and nutritional traits in foxtail millet (*Setaria italica* (L.) Beauv.). *Journal of Pharmacognosy and Phytochemistry*. 9(6): 596-600.
- Yadav, Y., Dhoot, R., Kumhar, S.R. and Roy, C. 2024. Estimates of variability, heritability and genetic advance in foxtail millets. *Annals of Arid Zone*. 63(1): 121-123.

Commented [DR4]: Not found in text

UNDER PEER REVIEW

Table. 1. Analysis of variance (ANOVA) for morpho-physiological and yield attributing traits in foxtail millet germplasm accessions

Formatted: Justified, Indent: Left: 0", Hanging: 0.69"

Table. 1. Analysis of variance (ANOVA) for morpho-physiological and yield attributing traits in foxtail millet germplasm accessions

* Significant at 5% level of probability

S. No	Traits	Mean sum of squares		
		Replications (df = 1)	Treatments (df = 49)	Error (df = 49)
1	Days to 50% flowering	6.250	11.597*	3.026
2	Days to maturity	0.040	22.936*	3.448
3	Plant height (cm)	189.888	202.206*	55.330
4	Number of productive tillers	0.004	0.321*	0.121
5	Flag leaf length at flowering (cm)	0.774	27.795*	4.628
6	Flag leaf width at flowering (cm)	0.280	0.163*	0.064
7	Panicle length (cm)	6.452	9.921*	1.712
8	Photosynthetic rate ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$)	2.904	92.674*	1.332
9	Transpiration rate ($\text{mmol H}_2\text{O m}^{-2} \text{ s}^{-1}$)	0.271	1.949*	0.262
10	Stomatal conductance ($\text{mmol H}_2\text{O m}^{-2} \text{ s}^{-1}$)	0.001	0.007*	0.001
11	Relative water content (%)	0.325	53.056*	2.635
12	Abortive grain rate	0.001	0.004*	0.001
13	1000 grain weight (g)	0.012	0.213*	0.010
14	Fodder yield per plant (g)	2.214	8.389*	1.971
15	Harvest index (%)	0.738	31.615*	0.790
16	Grain yield per plant (g)	1.201	6.155*	0.845

Formatted Table

Table. 2. Mean performance of the top ten accessions of foxtail millet including checks for morpho-physiological and yield attributing traits

Entry	DFP	DM	PH	NPT	FLL	FLW	PL	TGW	FYPP	HI	PR	TR	SC	RWC	AGR	GYPP
SiA 3290	56.50	88.00	138.00	2.40	33.30	2.34	17.90	2.90	15.31	39.87	36.56	5.63	0.27	83.92	0.16	11.59
SiA 4345	53.50	85.50	146.80	2.55	30.40	2.30	18.10	2.75	16.24	37.91	40.43	6.49	0.29	90.77	0.17	11.35
SiA 4391	56.50	84.50	141.50	2.25	34.20	1.98	19.00	2.55	15.39	41.33	37.57	6.35	0.24	88.8	0.06	11.32
SiA 3359	56.00	89.50	129.90	1.90	31.40	1.95	19.90	2.45	14.82	38.99	36.94	4.49	0.24	87.99	0.18	10.79
SiA 4435	56.50	91.50	141.70	2.30	35.80	2.10	18.30	2.50	13.66	40.13	34.59	3.98	0.15	90.00	0.13	10.16
SiA 4436	57.00	92.00	153.00	2.30	32.30	1.88	19.20	3.10	15.93	36.53	39.27	6.88	0.33	88.10	0.14	10.14
SiA 4468	54.50	88.00	142.00	2.70	32.00	2.00	19.70	2.85	11.94	41.70	34.43	4.71	0.19	85.79	0.15	9.80
SiA 3156 (C)	54.00	89.00	143.50	1.90	33.50	2.41	19.75	3.05	12.68	40.29	33.01	5.98	0.26	86.79	0.15	9.77
SiA 3223 (C)	53.50	91.00	152.50	1.90	34.60	2.52	19.40	3.05	14.39	38.74	41.51	6.21	0.34	88.38	0.13	10.09
SiA 3159 (#C)	50.00	84.00	143.20	2.10	32.50	2.29	21.05	2.95	13.82	42.06	45.97	6.24	0.29	90.55	0.11	11.04
C.V%	3.15	2.10	5.29	15.55	6.84	12.40	7.20	3.71	10.17	5.58	3.85	11.02	14.85	1.93	19.58	11.08
S. E+ (m)	1.23	1.31	5.26	0.18	1.52	0.24	0.92	0.07	0.99	1.38	0.82	0.36	0.021	1.15	0.017	0.65
C.D at 5%	3.49	3.73	14.95	0.51	4.32	0.70	2.63	0.20	2.82	3.91	2.32	1.03	0.059	3.26	0.049	1.85

C - Checks and #C - Best performing check

DFP: Days to 50% flowering; **DM:** Days to maturity; **PH:** Plant height (cm); **NPT:** No. of productive tillers; **FLL:** Flag leaf length at flowering (cm); **FLW:** Flag leaf width at flowering (cm); **PL:** Panicle length (cm); **TGW:** Thousand grain weight (g); **FYPP:** Fodder yield per plant (g); **HI:** Harvest index (%); **PR:** Photosynthetic rate ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$); **TR:** Transpiration rate ($\text{mmol H}_2\text{O m}^{-2} \text{ s}^{-1}$); **SC:** Stomatal conductance ($\text{mmol H}_2\text{O m}^{-2} \text{ s}^{-1}$); **RWC:** Relative water content (%); **AGR:** Abortive grain rate; **GYPP:** Grain yield per plant (g).

Commented [DR5]: LSD mean Least Significant Different (not CD)

Table 3. Estimates of range of variation and genetic variability parameters for the 16 traits in 50 foxtail millet accessions

S. no	Traits	Range		Mean	Coefficients of variation (%)		Heritability in broad sense (h^2_b) (%)	Genetic advance as per cent of mean (%)
		Min.	Max.		Phenotypic	Genotypic		
1	Days to 50% flowering	48.00	60.50	55.15	4.90	3.75	58.60	5.92
2	Days to maturity	77.00	94.00	88.46	4.11	3.53	73.90	6.25
3	Plant height (cm)	109.50	155.70	140.50	8.08	6.10	57.00	9.49
4	Number of productive tillers	1.50	3.20	2.04	21.02	14.14	45.20	19.60
5	Flag leaf length at flowering (cm)	23.10	39.70	31.44	12.80	10.82	71.50	18.84
6	Flag leaf width at flowering (cm)	1.42	2.86	2.24	16.52	10.91	43.60	14.84
7	Panicle length (cm)	12.90	21.90	18.61	13.28	11.16	70.60	19.30
8	Photosynthetic rate ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$)	13.47	45.97	29.99	22.85	22.52	97.20	45.74
9	Transpiration rate ($\text{mmol H}_2\text{O m}^{-2} \text{ s}^{-1}$)	2.98	6.88	4.64	22.64	19.77	76.30	35.58
10	Stomatal conductance ($\text{mmol H}_2\text{O m}^{-2} \text{ s}^{-1}$)	0.12	0.34	0.212	29.71	25.73	75.00	45.90
11	Relative water content (%)	71.82	90.77	83.96	6.28	5.98	90.50	11.72
12	Abortive grain rate	0.06	0.25	0.16	30.96	23.98	60.00	38.27
13	1000 grain weight (g)	1.90	3.35	2.69	12.40	11.83	91.00	23.25
14	Fodder yield per plant (g)	6.58	17.48	13.79	16.50	12.99	61.90	21.06
15	Harvest index (%)	26.44	42.06	34.88	12.06	10.69	78.60	19.52
16	Grain yield per plant (g)	4.44	11.59	8.29	22.55	19.64	75.90	35.24




Fig.1. Histogram depicting estimates of phenotypic and genotypic coefficient of variation for 16 morpho-physiological and yield attributing traits in 50 foxtail millet germplasm accessions




Fig.2. Histogram depicting estimates of heritability and genetic advance as *per cent* of mean for 16 morpho-physiological and yield attributing traits in 50 foxtail millet germplasm accessions