

# **Machine Learning and Artificial Intelligence in Thyroid Cancer Screening and Diagnosis: A Comprehensive Systematic Review**

## **Abstract**

This systematic review explores the role of artificial intelligence (AI) and machine learning (ML) technologies in the diagnosis and treatment of thyroid cancers (TC), focusing on enhancing precision, risk assessment, and tailored care. By analyzing ten studies, the review highlights how AI and ML technologies, such as deep learning (DL) and computer-aided diagnostics (CAD), improve the accuracy of ultrasound imaging, risk stratification, and the detection of high-risk nodules. Despite advancements, challenges persist in transitioning to personalized care, including uneven prognostication and diagnostic uncertainty. The review evaluates the effectiveness of AI and ML compared to conventional methods, their ability to address diverse tumor characteristics, and their strengths and limitations in prognosis prediction. Findings suggest AI's potential in improving precision and risk assessment, but limitations such as inconsistent approaches and biases highlight the need for larger datasets and standardized procedures. Moreover, the review underscores the importance of interpretability and transparency in AI models and calls for further research to validate findings in clinical settings. Despite limitations and challenges, AI's transformative potential in TC management is evident, underscoring the need for ongoing investigation and integration into clinical practice.

## **Introduction**

Over the past two decades, there has been a notable increase in the incidence of thyroid cancers (TC); the majority of TC cases are indolent [1][2]. Addressing these trends is crucial, given the continuous rise in incidence and death rates for aggressive papillary thyroid carcinomas (PTC) and advanced thyroid malignancies [1,3,4]. Accurate and efficient risk assessment is essential in

the era of customized healthcare to tailor therapy effectively. Understanding TC's biological function, characterized by diverse morphological traits and molecular elements, is the initial step[40-43]. While image analysis remains the primary diagnostic method for TC, its limitations in providing a thorough evaluation are evident [5]. Primary human cell cultures from surgical biopsies and fine-needle aspiration (FNA) samples offer opportunities for customized treatments, though challenges persist in transitioning to personalized care [6], such as uneven prognostication and uncertainty surrounding cytopathological diagnosis.

Radiologists have identified computer-aided diagnostics (CAD) as valuable for identifying cancers beyond breast cancer [11]. Assessing disease phases aids in determining the extent of thyroid cancer progression. Deep learning (DL) enhances ultrasound (US) accuracy by extracting nonlinear features [12]. Artificial intelligence (AI) facilitates improved operational performance and swift access to critical information for physicians. CAD and AI simplify risk-stratification systems, enhancing thyroid nodule detection and evaluation [13]. Molecular testing combined with machine learning (ML) techniques helps forecast and detect high-risk nodules [14]. ML's intrinsic power in drawing conclusions beyond traditional statistical approaches is evident [15]. Classification models developed using ML methods show promise in improving thyroid imaging assessment CAD systems [16].

Machine learning (ML) enables completion of complex tasks, such as photo interpretation [18]. DL aids in lung cancer detection on CT images. ML applications are growing, offering a comprehensive approach to cancer diagnosis and prevention [25]. Clinical parameters influence disease prognosis, with ML generating predictions to assist in patient disease management [22,23]. Protein markers and microarray data are increasingly relied upon in cancer diagnosis [24]. ML techniques, including supervised and unsupervised methods, are expanding in healthcare domains [26]. Metabolomics technology sheds light on lung cancer characteristics [25]. ML aids in diagnosing cancer types, predicting susceptibility, and screening individuals [27]. ML models improve tumor diagnostic accuracy and optimize therapeutic approaches [27].

The most frequently used neural networks in oncology are the convolution neural network (CNN), recurrent neural network (RNN), and multilayer perceptron (MLP). Cytopathology and histology are common methods for cancer diagnosis [28]. Histology-based CNNs classify

prostate, breast, and colon cancers successfully [29]. DL effectively distinguishes between benign and malignant tissues in lung cancer using whole-slide imaging. ML aids in forecasting tumor origins, even when unknown causes contribute to cancer cases [30].

## **Objective**

The systematic review aimed to conduct a comprehensive assessment and data compilation regarding the utilization of machine learning (ML) and artificial intelligence (AI) in thyroid tumors (TC). The primary objective was to evaluate the potential enhancements by AI and ML in the diagnosis, prognostication, and management of thyroid cancer. The study scrutinized both the advantages and limitations associated with employing AI and ML for the analysis of diagnostic imaging. Its focus centered on addressing the following inquiries:

1. How do AI and ML technologies compare in detecting thyroid cancers versus conventional imaging modalities?
2. Can AI and ML accommodate the varied biological and physical characteristics of thyroid tumors to enhance risk stratification and tailor personalized treatment?
3. What are the advantages and limitations of existing AI and ML models in distinguishing between benign and malignant thyroid nodules?
4. To what extent can AI and ML aid in prognostication and early detection of thyroid cancer, particularly in asymptomatic patients?

## **Methodology**

### Search Strategy

We extensively searched through various databases like Embase, Web of Science, PubMed, and Scopus to find studies about how artificial intelligence (AI) and machine learning (ML) are used

in dealing with thyroid cancer. To make sure we found everything relevant, we used different combinations of keywords related to AI, ML, thyroid cancers, and related topics. Our search method was carefully designed with boolean operators to match the syntax of each database.

### Eligibility Criteria

#### Inclusion Criteria

We extensively searched through various databases like Embase, Web of Science, PubMed, and Scopus to find studies about how artificial intelligence (AI) and machine learning (ML) are used in dealing with thyroid cancer. To make sure we found everything relevant, we used different combinations of keywords related to AI, ML, thyroid cancers, and related topics. Our search method was carefully designed with boolean operators to match the syntax of each database.

#### Exclusion Criteria

To maintain the focus and quality of our review, we excluded studies that didn't meet certain criteria. Publications not in English were excluded for clarity. We also left out conference abstracts, letters, editorials, and case reports because they lack the depth needed for a systematic review. Materials that didn't specifically talk about using AI or ML for thyroid tumors were also excluded to stay on topic. We avoided duplicating data by excluding duplicate articles. Table 1 shows the PICOS framework and our criteria for this review.

**Table 1: PICOS framework and eligibility criteria**

<b>Criteria</b>	<b>Description</b>
Population	Human subjects diagnosed with thyroid cancers.

Intervention	Original research articles or reviews focusing on the application of AI and ML in the context of thyroid cancers.
Comparison	Not applicable (as this is not a comparative study).
Outcomes	Studies reporting outcomes related to the diagnosis, prognosis, or management of thyroid cancers using AI or ML techniques.
Study Design	Various study designs, including observational studies, clinical trials, and reviews.

### Data Extraction

We carefully gathered all relevant information from each included study to synthesize our findings. This included details like authors, publication year, research design, and participant demographics. We explained the AI and ML techniques used in each study to help readers understand the methods. We also extracted data about how AI or ML methods were used for diagnosing, predicting outcomes, and treating thyroid cancers. Key results and conclusions from each study were noted to give a comprehensive overview.

### Quality Assessment

We used established tools to assess the quality of the included studies based on their research designs. For clinical trials, we used the Cochrane Risk of Bias tool, and for observational research, we used the Newcastle-Ottawa Scale. Two reviewers independently evaluated each

study, resolving any discrepancies through discussion or consultation with a third reviewer if needed.

## **Results**


### Study Selection

A systematic search across PubMed, Cochrane Library, and Google Scholar databases yielded a total of 218 records. After removal of duplicates, 121 records remained. Screening of titles and abstracts narrowed down the selection to 74 potentially relevant records. Following full-text screening, 10 studies met the inclusion criteria for the systematic review. Figure 1 illustrates the comprehensive flow diagram depicting the search and selection process.




### **Figure 1; PRISMA Flow Chart**

Identification of new studies via databases and registers

Identification



Screening



Included




Table 2 outlines the studies incorporated in this article.

**Table 2: Characteristics of included studies**

<b>Author &amp; Year</b>	<b>Study Design</b>	<b>Interventions</b>	<b>Population</b>	<b>Outcome Measures</b>	<b>Findings</b>
--------------------------	---------------------	----------------------	-------------------	-------------------------	-----------------

<p>Yoon et al. (2020)</p>	<p>Retrospective Observational Study</p>	<p>Computer-aided diagnosis (CAD) program using deep learning Convolutional Neural Network (CNN)</p>	<p>469 patients with thyroid cancer (380 positive, 89 negative for BRAFV600E mutation)</p>	<p>Association of CAD value with BRAFV600E mutation, Area Under the Receiver Operating Characteristic (AUC) of Receiver Operating Characteristic (ROC) curves for CAD value and multivariable model</p>	<p>Older age, smaller size, and higher CAD value significantly associated with BRAFV600E mutation. CAD value yielded an AUC of 0.646 for predicting BRAFV600E mutation. Multivariable model (age, size, and CAD value) had an AUC of 0.706, significantly better than CAD value alone. Deep learning-based CAD program shows promise in predicting BRAFV600E mutation in thyroid cancer. Multicenter studies with larger sample sizes are recommended for further validation.</p>
---------------------------	--	--	--	---	---

<p>Bellantuono et al. (2023)</p>	<p>eXplainable Artificial Intelligence analysis</p>	<p>Machine Learning procedure for discrimination of healthy/benign vs. malignant nodules using Raman spectra, Boruta feature selection, Synthetic Minority Over-sampling Technique (SMOTE) algorithm for imbalanced dataset, Random Forest, eXtreme Gradient Boosting (XGBoost), Support Vector Machine (SVM), and Gaussian Naïve Bayes classifiers</p>	<p>Patients with thyroid nodular pathology, 54 subjects (34 females, 20 males), aged 46.3 years on average, who underwent surgery (total thyroidectomy) after a cytological diagnosis of indeterminate, suspicious, or malignant nodules</p>	<p>Classification performance of Machine Learning algorithms (Random Forest, XGBoost, SVM, Gaussian Naïve Bayes) quantified by AUC, feature importance using Boruta, and synthetic data generation using SMOTE</p>	<p>Random Forest is identified as the best classifier (median AUC 0.9441, interquartile range 0.0049) for healthy/benign vs. cancer tissue classification. XGBoost, SVM, and Gaussian Naïve Bayes also explored. eXplainable Artificial Intelligence (XAI) analysis (SHapley Additive exPlanations - SHAP values) for interpretability. Performance evaluated on 72 samples (59 unambiguous and 13 ambiguous). Identified limitations in classifying ambiguous spectra with reduced AUC (median 0.7949, IQR 0.0135). Impactful features include carotenoid and oxidized cytochrome bands.</p>
----------------------------------	---	---	--	--	---

<p>Ha, E. J., &amp; Baek, J. H. (2021)</p>	<p>Review and development of AI-based CAD systems</p>	<p>Application of CAD systems by loading ultrasound images from Picture Archiving and Communication System (PACS). Real-time application during Ultrasound (US) examinations</p>	<p>Patients with thyroid nodules undergoing ultrasound imaging</p>	<p>Analysis of sonographic characteristics (echogenic foci, echogenicity, texture, margin, anechoic areas, height/width ratio, nodule shape, and size) and risk of malignancy based on Thyroid Imaging Reporting and Data System (TI-RADS) classifications</p>	<p>AmCAD-UT: Similar sensitivity (87.0%) but lower specificity (68.8%) compared to clinical experts using TI-RADS. Food and Drug Administration (FDA) 510(k) cleared. S-Detect 1: Comparable sensitivities (80.0%-92.0%) but lower specificity (74.6%-88.1%) compared to experienced radiologists. FDA approval in progress. S-Detect 2: Comparable sensitivities (81.4%) but lower specificity (68.2%-81.9%) compared to experienced radiologists.</p>
--	---	--	--	--	---

<p>Agarwal et al. (2021)</p>	<p>Evaluation and comparison of AI algorithms</p>	<p>Implementation of AI algorithms and machine learning models to analyze diagnostic imaging data</p>	<p>Individuals undergoing diagnostic tests for cancer, including imaging tests, endoscopic procedures, biopsy, and cytology</p>	<p>Assessment of the diagnostic accuracy of AI algorithms and machine learning models in differentiating benign and malignant tumors</p>	<p>AI improves diagnostic accuracy by analyzing large imaging datasets, leveraging technical advances and hardware enhancements for neural network training. It excels in early diagnosis, particularly in breast and lung cancer, surpassing human specialists in breast cancer prognosis and providing early lung cancer predictions. In gastric cancer, Convolutional Neural Networks aid in invasion depth diagnosis through gastric endoscopy. AI techniques, coupled with imagery, enable early identification of oral cancer. Overall, AI significantly enhances cancer diagnosis precision and extends forecasting capabilities.</p>
------------------------------	---	---	---	--	--

Xi et al. (2022)	Prospective study using machine learning	Six machine learning models trained on a clinical dataset from 724 patients undergoing thyroidectomy. Models included Gradient Boosting, Logistic Regression, Linear Discriminant Analysis, SVM, and Random Forest	724 patients at Shengjing Hospital, China, with demographic info, ultrasound features, and blood test results	Models demonstrated superior accuracy, with Random Forest leading. Gradient Boosting excelled in sensitivity, Logistic Regression in specificity. Variable importance analysis highlighted key predictors. Models outperformed expert assessment in accuracy and F1 score	Machine learning, especially Random Forest and Gradient Boosting, improved thyroid nodule malignancy prediction compared to expert assessment. Models offered valuable insights into nodule characteristics, enhancing preoperative thyroid cancer diagnosis.
------------------	--	--	---	---	---

<p>Peng et al. (2021)</p>	<p>Multicentre Diagnostic Study</p>	<p>Development and application of the deep-learning AI model (ThyNet) for differentiating thyroid nodules</p>	<p>Patients aged 18 or older with thyroid nodules at least 3 mm in diameter identified via ultrasound</p>	<p>Primary Endpoint: Area Under the Receiver Operating Characteristic Curve (AUROC) for thyroid nodule diagnosis. Secondary Endpoints: Accuracy, Sensitivity, Specificity, Positive Predictive Value (PPV), Negative Predictive Value (NPV)</p>	<p>ThyNet AUROC: 0.922 (95% CI 0.910–0.934) was significantly higher than radiologists (<math>p &lt; 0.0001</math>). ThyNet-assisted strategy improved radiologists' AUROC from 0.837 to 0.875 (<math>p &lt; 0.0001</math>). In a simulated scenario, ThyNet-assisted strategy reduced unnecessary fine needle aspirations by 26.7%. Missed malignancy decreased from 18.9% to 17.0% with ThyNet-assisted strategy.</p>
---------------------------	-------------------------------------	---	---	---	---

<p>Olatunji et al. (2021)</p>	<p>Retrospective Case Study</p>	<p>Machine learning-based tools development for early detection of thyroid cancer (TC)</p>	<p>Techniques used: Random Forest (RF), Artificial Neural Network (ANN), Support Vector Machine (SVM), Naïve Bayes (NB)</p>	<p>Patients from the Kingdom of Saudi Arabia</p>	<p>RF technique demonstrated the highest accuracy at 90.91%. SVM, ANN, and NB achieved accuracy rates of 84.09%, 88.64%, and 81.82%, respectively. Emphasis on early detection at pre-symptomatic stages. RF recommended as the preferred technique for this specific problem.</p>
-------------------------------	---------------------------------	--	---	--	--

<p>Zhou et al. (2022)</p>	<p>Experimental Study (Comparison of new ultrasound technologies)</p>	<p>Ultrasonic intelligent diagnosis of papillary thyroid cancer based on machine learning, involving Contrast-Enhanced Ultrasound (CEUS) and Ultrasound Elastography (UE) technologies</p>	<p>Patients with papillary thyroid carcinoma (PTC), 70 cases (10 male, 60 female), tumor diameter <math>\leq 10</math> mm, 107 lymph nodes</p>	<p>Characteristics of ultrasound images (CEUS and UE). Diagnostic effectiveness of new ultrasound technologies (CEUS and UE) in distinguishing between benign and malignant nodules</p>	<p>CEUS and UE showed significant differences in enhancement mode, intensity, early regression, and edge enhancement between micro benign and malignant tumors. UE demonstrated higher sensitivity and diagnostic efficiency compared to CEUS in the differential diagnosis of thyroid micro benign and malignant nodules. Combined use of CEUS and UE resulted in 78.43% sensitivity and 78.67% specificity for diagnosing thyroid micro benign and malignant nodules.</p>
---------------------------	---	--	--	---	---

Zhao et al. (2019)	Meta-analysis	Evaluation of computer-aided diagnosis system (CAD) for thyroid nodules on ultrasound	536 patients with 723 thyroid nodules	Sensitivity, Specificity, Positive Likelihood Ratio (LR), Negative LR, Diagnostic Odds Ratio (DOR)	<p>Findings (CAD System): Sensitivity: 0.87 (95% CI, 0.73–0.94), Specificity: 0.79 (95% CI, 0.63–0.89), Positive LR: 4.1 (95% CI, 2.5–6.9), Negative LR: 0.17 (95% CI, 0.09–0.32), DOR: 25 (95% CI, 15–42), Summary Receiver Operating Characteristic (SROC) AUC: 0.90 (95% CI, 0.87–0.92). Findings (Experienced Radiologists): Sensitivity: 0.82 (95% CI, 0.69–0.91), Specificity: 0.83 (95% CI, 0.76–0.89), Positive LR: 4.9 (95% CI, 3.4–7.0), Negative LR: 0.22 (95% CI, 0.12–0.38), DOR: 23 (95% CI, 11–46), SROC AUC: 0.96 (95% CI, 0.94–0.97).</p>
--------------------	---------------	---	---------------------------------------	--	--

<p>Kuang et al. (2022)</p>	<p>Metabolomic analysis, machine-learning model development</p>	<p>Analysis of existing data of thyroid cancer (TC) metabolites, development of a machine-learning model using metabolite biomarkers</p>	<p>The study involved datasets related to papillary thyroid cancer (PTC) patients</p>	<p>Classification accuracy of machine-learning models (LogitBoost, AdaBoostM1, RandomForest, etc.) through 10-fold cross-validation. Identification of metabolic pathways related to TC</p>	<p>Highest classification accuracy: LogitBoost - 87.30%, Various classifiers achieved accuracies above 80%. Independent testing showed 100% accuracy in identifying TC-related metabolites .</p>
----------------------------	---	--	---	---	--

## **Discussion**

A deep learning convolutional neural network (CNN)-based computer-aided diagnostic (CAD) tool was developed for diagnosing thyroid cancer in a retrospective observational study conducted by Yoon et al. in 2020 [30]. Out of the 469 patients with thyroid cancer included in the study, 380 tested positive for the BRAFV600E mutation, while 89 did not. The association between the CAD value and the BRAFV600E mutation was assessed by calculating the area under the receiver operating characteristic (ROC) curve (AUC) for the CAD value and a multivariable model. It was found that the BRAFV600E mutation was significantly correlated with higher CAD values, smaller sizes, and older ages. The CAD value yielded an AUC of 0.646 for predicting the BRAFV600E mutation. When age, size, and CAD value were combined in the multivariable model, the AUC increased to 0.706, which was significantly higher than using the CAD value alone. Based on these results, the deep learning-based CAD program may have the ability to predict the BRAFV600E mutation in thyroid cancer. However, the authors suggest further validation through multicenter research with larger sample sizes.

Thyroid cancer detection was explored by Bellantuono et al. in a paper published in 2023 using explainable Artificial Intelligence (XAI) analysis of Raman spectra [31]. Various classifiers such as Random Forest, XGBoost, Support Vector Machine, and Gaussian Naïve Bayes were employed, along with Boruta feature selection, the synthetic minority oversampling technique (SMOTE) algorithm for imbalanced datasets, and Raman spectra, to differentiate between benign and malignant nodules. The study included 54 patients with thyroid nodular pathology (mean age 46.3 years) who underwent total thyroidectomy following a cytological diagnosis of ambiguous, suspicious, or malignant nodules. The Random Forest classifier performed the best, achieving a median AUC of 0.9441 for classifying tissue as either cancerous or healthy. Important features like oxidized and carotenoid cytochrome bands were identified using XAI analysis and SHAP values for interpretability. The study highlighted the challenges of identifying ambiguous spectra

while showcasing the potential of machine learning in diagnosing thyroid cancer with the assistance of XAI.

Ha and Baek provided a comprehensive analysis and overview of AI-based computer-aided diagnostic (CAD) techniques for thyroid nodules in 2021 [32]. CAD systems were used in real time during ultrasonography tests by importing ultrasound images from Picture Archiving and Communication Systems (PACS). The study focused on sonographic features, using the Thyroid Imaging Reporting and Data System (TI-RADS) categories to determine malignancy risk. Various CAD systems like AmCAD-UT, S-Detect 1, and S-Detect 2 were evaluated against radiologists and clinical specialists. Although the sensitivity of CAD systems was similar to or slightly lower than that of clinical specialists, the results showed that CAD systems can be used in real-time ultrasound exams.

Agarwal et al. compared machine learning models and AI algorithms for cancer diagnosis in 2021 [33]. Using AI algorithms and machine learning models, the study analyzed diagnostic imaging data from patients undergoing various cancer diagnostic procedures, focusing on differentiating between benign and malignant cancers. The findings demonstrated how AI techniques can improve the precision of cancer diagnosis by leveraging hardware updates and technology breakthroughs to train neural networks. It was shown that AI algorithms can be used for early detection, prognostication, and enhanced accuracy in various cancer types, including stomach, oral, lung, and breast cancer. The study emphasized how AI-generated predictions, which are more detailed and accurate, can enhance overall cancer detection.

Xi et al. used ten-fold cross-validation and machine learning with bootstrap analysis in a prospective study to predict thyroid nodule malignancy [34]. Using a clinical dataset of 724 patients undergoing thyroidectomy, six machine learning models were developed, with Random Forest performing the best overall. Logistic Regression showed excellent specificity, while Gradient Boosting had better sensitivity. The models were better in terms of accuracy and F1 score than expert judgment, demonstrating the potential of machine learning, especially Random Forest and Gradient Boosting, in preoperative thyroid cancer diagnosis.

Peng et al. created and utilized the ThyNet deep-learning AI model to differentiate thyroid nodules in a multicenter diagnostic investigation [35]. ThyNet outperformed radiologists in

identifying thyroid nodules, improving accuracy and reducing unnecessary procedures. Olatunji et al. conducted a retrospective case study in 2021, demonstrating high accuracy rates for early thyroid cancer identification using machine learning techniques [36]. Zhou et al. evaluated machine learning for thyroid nodule identification, highlighting the potential of combining ultrasound elastography (UE) and contrast-enhanced ultrasound (CEUS) for improved accuracy [37]. Lastly, Zhao et al. compared a computer-aided diagnostic system (CAD) with skilled radiologists, showing that the CAD system outperformed radiologists in identifying thyroid nodules [38]. Kuang et al. used metabolomic analysis and machine learning to classify metabolite biomarkers associated with thyroid cancer, showing promising results in efficiently identifying thyroid cancer [39].

## **Conclusion**

In conclusion, while significant advancements have been made in AI and ML for diagnosing and prognosticating thyroid cancer, caution is warranted due to discrepancies and shortcomings noted in various studies. Future research in this area should prioritize standardized methodologies, larger and more diverse sample sizes, and inclusion of various patient demographics to enhance therapeutic utility and generalizability. Additionally, considerations of interpretability, transparency, and ethical implications will be pivotal in the seamless integration of AI and ML into routine clinical practice. Overall, this systematic review underscores the need for further investigation and validation in real-world clinical settings and emphasizes the transformative potential of AI in the management of thyroid cancer.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

### **Limitations**

The reviewed literature, despite its promising implications, exhibits significant limitations. The wide variations in research methodologies, patient cohorts, and AI models make it challenging to draw definitive conclusions applicable across different contexts. Given that many studies rely on observational or retrospective designs, larger-scale randomized controlled trials may be essential to ascertain the therapeutic effectiveness of AI and ML in diagnosing and treating thyroid cancer. Moreover, the diversity of data sources, imaging techniques, and diagnostic criteria among studies introduces potential biases and hampers the generalizability of findings. For instance, Bellantuono et al.'s study illustrates the challenges in distinguishing ambiguous spectra, underscoring practical limitations.

Yoon et al.'s suggestion for larger, multicenter investigations underscores the need for more comprehensive validation and highlights the constraints imposed by current sample sizes. Addressing issues related to interpretability and transparency of AI models, as demonstrated by Bellantuono et al.'s XAI project, is imperative before widespread adoption of AI technology in healthcare settings. Furthermore, the focus on specific AI applications and exclusion of non-English research may introduce publication bias and limit our understanding of the field.

## **References**

1. Lim H, Devesa SS, Sosa JA, Check D, Kitahara CM. Trends in thyroid cancer incidence and mortality in the United States, 1974-2013. *Jama*. 2017 Apr 4;317(13):1338-48.
2. La Vecchia C, Malvezzi M, Bosetti C, Garavello W, Bertuccio P, Levi F, Negri E. Thyroid cancer mortality and incidence: a global overview. *International journal of cancer*. 2015 May 1;136(9):2187-95.
3. Ho AS, Luu M, Barrios L, Chen I, Melany M, Ali N, Patio C, Chen Y, Bose S, Fan X, Clair JM. Incidence and mortality risk spectrum across aggressive variants of papillary thyroid carcinoma. *JAMA oncology*. 2020 May 1;6(5):706-13.
4. Jegerlehner S, Bulliard JL, Aujesky D, Rodondi N, Germann S, Konzelmann I, Chiolero A, NICER Working Group. Overdiagnosis and overtreatment of thyroid cancer: A population-based temporal trend study. *PloS one*. 2017 Jun 14;12(6):e0179387.
5. Koelzer VH, Sirinukunwattana K, Rittscher J, Mertz KD. Precision immunoprofiling by image analysis and artificial intelligence. *Virchows Archiv*. 2019 Apr 1;474:511-22.
6. Fallahi P, Ferrari SM, Elia G, Ragusa F, Patrizio A, Paparo SR, Marone G, Galdiero MR, Guglielmi G, Foddis R, Cristaudo A. Primary cell cultures for the personalized therapy in aggressive thyroid cancer of follicular origin. In *Seminars in cancer biology* 2022 Feb 1 (Vol. 79, pp. 203-216). Academic Press.
7. Persichetti A, Di Stasio E, Coccaro C, Graziano F, Bianchini A, Di Donna V, Corsello S, Valle D, Bizzarri G, Frasoldati A, Pontecorvi A. Inter- and intraobserver agreement in the assessment of thyroid nodule ultrasound features and classification systems: a blinded multicenter study. *Thyroid*. 2020 Feb 1;30(2):237-42.

8. Cibas ES, Ali SZ. The 2017 Bethesda system for reporting thyroid cytopathology. *Thyroid*. 2017 Nov 1;27(11):1341-6.
9. Baloch ZW, LiVolsi VA. Follicular-patterned lesions of the thyroid: the bane of the pathologist. *American journal of clinical pathology*. 2002 Jan 1;117(1):143-50.
10. LiVolsi VA, Baloch ZW. Follicular neoplasms of the thyroid: view, biases, and experiences. *Advances in Anatomic Pathology*. 2004 Nov 1;11(6):279-87.
11. Hadjiiski L, Sahiner B, Chan HP. Advances in CAD for diagnosis of breast cancer. *Current opinion in obstetrics & gynecology*. 2006 Feb;18(1):64.
12. Brattain LJ, Telfer BA, Dhyani M, Grajo JR, Samir AE. Machine learning for medical ultrasound: status, methods, and future opportunities. *Abdominal radiology*. 2018 Apr;43:786-99.
13. Zhao R, Yan R, Chen Z, Mao K, Wang P, Gao RX. Deep learning and its applications to machine health monitoring. *Mechanical Systems and Signal Processing*. 2019 Jan 15;115:213-37.
14. Daniels K, Gummadi S, Zhu Z, Wang S, Patel J, Swendseid B, Lyshchik A, Curry J, Cottrill E, Eisenbrey J. Machine learning by ultrasonography for genetic risk stratification of thyroid nodules. *JAMA Otolaryngology–Head & Neck Surgery*. 2020 Jan 1;146(1):36-41.
15. Cruz JA, Wishart DS. Applications of machine learning in cancer prediction and prognosis. *Cancer informatics*. 2006 Jan;2:117693510600200030.
16. Li X, Zhang S, Zhang Q, Wei X, Pan Y, Zhao J, Xin X, Qin C, Wang X, Li J, Yang F. Diagnosis of thyroid cancer using deep convolutional neural network models applied to sonographic images: a retrospective, multicohort, diagnostic study. *The Lancet*

Oncology. 2019 Feb 1;20(2):193-201.

17. Mongan J, Moy L, Kahn Jr CE. Checklist for artificial intelligence in medical imaging (CLAIM): a guide for authors and reviewers. *Radiology: Artificial Intelligence*. 2020 Mar 25;2(2):e200029.
18. Simes RJ. Treatment selection for cancer patients: application of statistical decision theory to the treatment of advanced ovarian cancer. *Journal of chronic diseases*. 1985 Jan 1;38(2):171-86.
19. Astion ML, Wilding P. Application of neural networks to the interpretation of laboratory data in cancer diagnosis. *Clinical Chemistry*. 1992 Jan 1;38(1):34-8.
20. Fielding LP, Fenoglio-Preiser CM, Freedman LS. The future of prognostic factors in outcome prediction for patients with cancer. *Cancer*. 1992 Nov 1;70(9):2367-77.
21. Cochran AJ. Prediction of outcome for patients with cutaneous melanoma. *Pigment cell research*. 1997 Jun;10(3):162-7.
22. Burke HB, Bostwick DG, Meiers I, Montironi R. Prostate cancer outcome: epidemiology and biostatistics. *Analytical and quantitative cytology and histology*. 2005 Aug 1;27(4):211-7.
23. Cruz JA, Wishart DS. Applications of machine learning in cancer prediction and prognosis. *Cancer informatics*. 2006 Jan;2:117693510600200030.
24. Sun R, Fei F, Wang M, Jiang J, Yang G, Yang N, Jin D, Xu Z, Cao B, Li J. Integration of metabolomics and machine learning revealed tryptophan metabolites are sensitive biomarkers of pemetrexed efficacy in non-small cell lung cancer. *Cancer Medicine*. 2023 Sep;12(18):19245-59.

25. Zhang B, Shi H, Wang H. Machine Learning and AI in Cancer Prognosis, Prediction, and Treatment Selection: A Critical Approach. *Journal of Multidisciplinary Healthcare*. 2023 Dec 31;1779-91.
26. Iqbal MJ, Javed Z, Sadia H, Qureshi IA, Irshad A, Ahmed R, Malik K, Raza S, Abbas A, Pezzani R, Sharifi-Rad J. Clinical applications of artificial intelligence and machine learning in cancer diagnosis: looking into the future. *Cancer cell international*. 2021 Dec;21(1):1-1.
27. Tran KA, Kondrashova O, Bradley A, Williams ED, Pearson JV, Waddell N. Deep learning in cancer diagnosis, prognosis and treatment selection. *Genome Medicine*. 2021 Dec;13(1):1-7.
28. Ryu HS, Jin MS, Park JH, Lee S, Cho J, Oh S, Kwak TY, Woo JI, Mun Y, Kim SW, Hwang S. Automated Gleason scoring and tumor quantification in prostate core needle biopsy images using deep neural networks and its comparison with pathologist-based assessment. *Cancers*. 2019 Nov 25;11(12):1860.
29. Pavlidis N, Pentheroudakis G. Cancer of unknown primary site. *The Lancet*. 2012 Apr 14;379(9824):1428-35.
30. Yoon J, Lee E, Koo JS, Yoon JH, Nam KH, Lee J, Jo YS, Moon HJ, Park VY, Kwak JY. Artificial intelligence to predict the BRAFV600E mutation in patients with thyroid cancer. *PloS one*. 2020 Nov 25;15(11):e0242806.
31. Bellantuono L, Tommasi R, Pantaleo E, Verri M, Amoroso N, Crucitti P, Di Gioacchino M, Longo F, Monaco A, Naciu AM, Palermo A. An eXplainable Artificial Intelligence analysis of Raman spectra for thyroid cancer diagnosis. *Scientific Reports*. 2023 Oct 3;13(1):16590.

32. Ha EJ, Baek JH. Applications of machine learning and deep learning to thyroid imaging: where do we stand?. *Ultrasonography*. 2021 Jan;40(1):23.
33. Agarwal S, Yadav AS, Dinesh V, Vatsav KS, Prakash KS, Jaiswal S. By artificial intelligence algorithms and machine learning models to diagnosis cancer. *Materials Today: Proceedings*. 2023 Jan 1;80:2969-75.
34. Xi NM, Wang L, Yang C. Improving the diagnosis of thyroid cancer by machine learning and clinical data. *Scientific Reports*. 2022 Jul 1;12(1):11143.
35. Peng S, Liu Y, Lv W, Liu L, Zhou Q, Yang H, Ren J, Liu G, Wang X, Zhang X, Du Q. Deep learning-based artificial intelligence model to assist thyroid nodule diagnosis and management: a multicentre diagnostic study. *The Lancet Digital Health*. 2021 Apr 1;3(4):e250-9.
36. Olatunji SO, Alotaibi S, Almutairi E, Alrabae Z, Almajid Y, Altabee R, Altassan M, Ahmed MI, Farooqui M, Alhiyafi J. Early diagnosis of thyroid cancer diseases using computational intelligence techniques: A case study of a Saudi Arabian dataset. *Computers in Biology and Medicine*. 2021 Apr 1;131:104267.
37. Zhou H, Liu B, Liu Y, Huang Q, Yan W. Ultrasonic intelligent diagnosis of papillary thyroid carcinoma based on machine learning. *Journal of Healthcare Engineering*. 2022 Jan 10;2022.
38. Zhao WJ, Fu LR, Huang ZM, Zhu JQ, Ma BY. Effectiveness evaluation of computer-aided diagnosis system for the diagnosis of thyroid nodules on ultrasound: A systematic review and meta-analysis. *Medicine*. 2019 Aug;98(32).
39. Kuang A, Kouznetsova VL, Kesari S, Tsigelny IF. Diagnostics of Thyroid Cancer Using Machine Learning and Metabolomics. *Metabolites*. 2023 Dec 22;14(1):11.

40. Rajasekharan, C., K. V. Vysakha, S. Aswanikumar, and N. S. Sibi. 2016. "Is It Prudent to Screen the Thyroid before Initiating Treatment for Acromegaly?". *Journal of Advances in Medicine and Medical Research* 16 (9):1-5. <https://doi.org/10.9734/BJMMR/2016/25626>.
41. Goyal, Pankaj, Kishan Kumawat, G. N. Gupta, and Nidhi P. Chanchlani. 2023. "A Case Report on Adenoid Cystic Carcinoma of Nasal Cavity: Successfully Treated by Endoscopic Surgical Excision". *International Research Journal of Oncology* 6 (1):9-15. <https://journalirjo.com/index.php/IRJO/article/view/119>.
42. Lamartina L, Grani G, Durante C, Filetti S, Cooper DS. Screening for differentiated thyroid cancer in selected populations. *The lancet Diabetes & endocrinology*. 2020 Jan 1;8(1):81-8.
43. Azizi G, Keller JM, Mayo ML, Piper K, Puett D, Earp KM, Malchoff CD. Thyroid nodules and shear wave elastography: a new tool in thyroid cancer detection. *Ultrasound in medicine & biology*. 2015 Nov 1;41(11):2855-65.