

# Effect of Different Organic Manures along with Microbial Consortia on Growth and Yield of Kalmegh (*Andrographis paniculata* Nees.)

## ABSTRACT

This study aimed to investigate the “Effect of different organic manures along with microbial consortia on growth and yield of Kalmegh (*Andrographis paniculata* Nees)”. Kalmegh has wide range of medicinal and pharmacological applications. The active ingredient andrographolide is important bioprotectant. Hence the need to explore different manures along with microbial consortia to improve its yield is essential. The experiment was laid out in RCBD with thirteen treatments. Different organic manures and combinations with neem cake were taken as treatments with three replications. Significantly higher plant height was observed in treatment - 100% RDF through vermicompost. Maximum plant spread was recorded in the treatment -50% RDF through FYM + 50% RDF through neem cake + microbial consortia. Application of sheep manure + neem cake resulted in higher number of branches plant<sup>-1</sup>, higher number of leaves plant<sup>-1</sup>, highest fresh weight, highest dry weight, highest root to shoot ratio, highest fresh herbage yield and highest B:C ratio. So, sheep manure + neem cake was found best compared to other manures and combinations.

## INTRODUCTION

Kalmegh (*Andrographis paniculata*) belong to the family ‘Acanthaceae’ which is also known as Maha tita and Kalpanath

The plant is also known as the ‘king of bitters’ [2-3] because it is extremely bitter in taste in every part of plant body. On the basis of literature survey, it has been observed that the aerial parts (leaves and stems) of the plants are most commonly used to extract the active phytochemicals, however the whole plant or roots are mentioned to a limited extent [3]. *A. paniculata* has a broad spectrum of pharmacological effects and some of them are extremely beneficial such as Hepatoprotective, antimicrobial, antifungal, antioxidant, anti-inflammatory, antipyretic, anticancer and antidiarrheal effects. (Nyeem *et al.*, 2017)

According to Unani system of medicine it is useful in the treatment of chronic hepatitis. The roots give flavones, apigenin-7, 4'-dio-O-methyl ether, 5-hydroxy-7, 8, 2', 3'-tetra methoxy flavone, andrographin and panicolin and sitosterol [13,14,15,16]. Leaves of Kalmegh contain homoandrographolide, andrographosterol and andrographone. Active compounds of Kalmegh sold in market by different trade names like Actizeet, Bixa botanical,

Life Aveda and other products like Acene-n-pimple cream (Himalaya wellness), Liv 52 (Himalaya Drugs Company), Lerbohep (Lupin Herbal Laboratory), Sage Liverex (Sage Herbals), Purodil syrup (Aimil Pharmaceuticals). Researches conducted in the past have confirmed that *A. paniculata* if properly administrated, has a surprisingly broad range of pharmacological effects. The major source of supply of Kalmegh is Tropical forests.

India is a global leader in the production of medicinal and aromatic plants. The National Medicinal Plants Board and the Food and Agriculture Organization recommended that all the medicinal and aromatic plants to be cultivated organically. Organically grown medicinal and aromatic products are not only readily acceptable in global market but also fetch premium prices than those grown with conventional farming. (Aishwath and Tarafdar) Annually, India's medicinal plants export is worth of Rs.1250 crores. Organically grown medicinal and aromatic plants have an excellent global market and India can exploit this market to its advantage (Rajeswara Rao and Rajput, 2005).

Results on application of organic manures on medicinal and aromatic plants revealed that enhancement of yield and uptake of nutrients. Not only organic manures, biofertilizers (Azotobacter and Azospirillum), arbuscular mycorrhizal (AM) fungi and phosphorus solubilizing bacteria, organic mulches and crop rotations are also play an important role for the enhancement of yield and nutrients utilization.

Influence of organic nutrients on biological yield and bioactive principle of Kalmegh is well pronounced according to different literatures. Organic manures and bio-fertilizers improve soil quality and beneficial soil microorganisms. Keeping the importance of both organic cultivation and beneficial effects of organic manure, the present investigation, study on effect of different organic manures along with microbial consortia on growth and yield of Kalmegh (*Andrographis paniculata* Nees) was carried out.

**Objective of the investigation:** Study on effect of different organic manures along with microbial consortia on growth and yield of *Kalmegh*.

## MATERIAL AND METHODS

- I. **Place of study:** Sanjeevini vatika, Medicinal plants block, Department of Horticulture, College of Agriculture, University of Agricultural Sciences, Gandhi Krishi Vigyana Kendra, Bengaluru.
- II. **Seed collection:** Kalmegh seeds of variety CIM- Megha was collected from CSIR-Central Institute of Medicinal and Aromatic Plants, Bengaluru.
- III. **Seedlings:** The seedlings were raised in the nursery and 30 days old seedlings were transplanted in the experimental plots.
- IV. **Materials:** Measuring scale, electronic weighing scale, hot air oven
- V. **Season:** *kharif* season during the year 2021-2022
- VI. **Location:** The experimental site is situated at 12° 58" North latitude and 77° 55" East longitude with an elevation of about 930 meters above mean sea level.
- VII. **Temperature, Rainfall & RH:** The mean of maximum and minimum temperature recorded during the period of experiment was 27.7° C and 18.5 ° C, respectively. The average of total rainfall received from June to September was 156 mm. The mean maximum and minimum Relative Humidity (RH) of the location was 89.5 per cent and 58.7 per cent, respectively.

Organic manures like Farm Yard Manure (FYM), Poultry manure, Sheep manure, Vermicompost and Neem cake were procured from Zonal Agricultural Research station (ZARS), University of Agricultural Sciences, Gandhi Krishi Vigyana Kendra, Bengaluru. Microbial consortia was purchased from bio-fertilizers production laboratory, Department of Microbiology, University of Agricultural Sciences, Gandhi Krishi Vigyana Kendra, Bengaluru. The above-mentioned microbial consortia consisted the mixture of phosphate solubilising bacteria, trichoderma and pseudomonas.

The experiment was laid-out in Randomized Complete Block Design (RCBD) replicated thrice with thirteen treatments in open field condition. The treatments consisted of combination of different organic manures with and without microbial consortia.

**Treatment details:**

T <sub>1</sub>	Control (No manure)
T <sub>2</sub>	100% RDF through FYM
T <sub>3</sub>	100% RDF through Sheep manure
T <sub>4</sub>	100 % RDF through Poultry manure
T <sub>5</sub>	100% RDF through Vermicompost
T <sub>6</sub>	50% RDF through FYM + 50% RDF through Neem cake
T <sub>7</sub>	50% RDF through Sheep manure + 50% RDF through Neem cake
T <sub>8</sub>	50% RDF through Poultry manure + 50% RDF through neem cake
T <sub>9</sub>	50% RDF through Vermicompost + 50% RDF through Neem cake
T <sub>10</sub>	50% RDF through FYM + 50% RDF through Neem cake 50% + Microbial consortium
T <sub>11</sub>	50% RDF through Sheep manure + 50% RDF through Neem cake + Microbial consortium
T <sub>12</sub>	50% RDF through Poultry manure + 50% RDF through Neem cake + Microbial consortium
T <sub>13</sub>	50% RDF through Vermicompost + 50% RDF through Neem cake + Microbial consortium

**RDF: Recommended Dosage of Fertilizer**

Organic manures were calculated on N equivalent basis (RDF: 75: 75: 50 N, P and K kg/ha) and the spacing followed was 30 cm X 15 cm with the individual plot size of 2.7m X 2.1m.


**RESULTS AND DISCUSSION****Growth parameters:****Plant height (cm)**

The data pertaining to effect of different organic manures with and without microbial consortia on plant height of Kalmegh is presented in Table 1 and depicted in Figure.1

The result with respect to plant height at 90 Days After Transplanting (DAT) was found to be significant. Highest plant height (49.4cm) was recorded in the treatment T<sub>5</sub>

(100% RDF through vermicompost) and least plant height was observed in the treatment T<sub>1</sub> (Control).

The significant increase in plant height was noticed with the application of poultry manure. This might be due to the fact that it is higher in nitrogen content and has instant nutrient releasing property along with rich organic matter content. This might have resulted in recording the higher values of plant height. Similarly, vermicompost was found to be best due to presence of plant growth hormones in it and is rich in humus than other organic manure but slower in decomposition rate when compared poultry manure. Similar results were reported by Kumar and Topal (2015) and Detpiratmongkol *et al.* (2014) in Kalmegh and Thanuja *et al.* (2020) in Ajwain; Ronya *et al.* (2020) in Turmeric; Smitha *et al.* (2019) in sacred basil.



**Fig.1: Effect of different organic manures along with microbial consortia on plant height in Kalmegh**

### Number of branches per plant

The data pertaining to effect of different organic manures on number of branches in Kalmegh is presented in Table 1.

The data recorded on number of branches per plant was found to be significant. The treatment provided with 100% RDF through poultry manure has recorded significantly highest number of branches (11.60) at 60 DAT, which was found to be on par with treatment T<sub>7</sub> (11.13) while the least (5.20) was recorded in the treatment T<sub>1</sub> (Control).

The data with respect to number of branches at 90 DAT was found to be significant. Highest number of branches (14) was recorded in the treatment T<sub>7</sub> (50% RDF through Sheep manure + 50% RDF through neem cake) which was followed by the treatment T<sub>4</sub> (13.67) and least number (10.0) of branches were observed in the treatment T<sub>1</sub> (Control).

### **Plant Spread (cm<sup>2</sup>)**

The perusal of data presented in Table 1 indicate that different organic manures along with and without microbial consortia had significant influence on plant spread.

The result with respect to plant spread at 90 DAT was found to be significant. Highest plant spread (2209.6 cm<sup>2</sup>) was recorded in the treatment T<sub>10</sub>(50% RDF through FYM+ 50% RDF through Neem cake + Microbial consortium) which was on par with treatment T<sub>4</sub>(1475.8 cm<sup>2</sup>) and least plant spread (775.2 cm<sup>2</sup>) was observed in the treatment T<sub>1</sub> (Control).

The significant difference in plant spread was noticed with the application FYM along with neem cake because FYM with combination of neem cake releases nutrients at a slower rate. So, at later stage plants treated with the combination of FYM, neem cake and microbial consortia might have shown better growth. **Similar findings were studied by** Tiwari *et al.* (2012) in Kalmegh, Smitha *et al* (2019) recorded maximum plant spread in sacred basil where organic manures and biofertilizers used in combination as compared to application of biofertilizer or organic manure alone. Singh *et al.* (2012) worked on *Coleus forskohlii* plants treated with Arbuscular Mychorriza, Neem cake and *Pseudomonas florescence* showed significantly increased plant spread.

### **Yield parameters**

#### **Fresh weight of plant (g)**

The fresh weight of plant influenced by different organic manures along with microbial consortia in Kalmegh is shown in the Table 2 and Figure 2.

Different organic manures along with microbial consortia influenced significantly on fresh weight of plant after 90 DAT (at harvest). The highest fresh weight of 56.60g was recorded in plant provided with 50% RDF through Sheep manure + 50% RDF through Neem

cake followed by 54.20 g in plants provided with 50% RDF through Poultry manure + 50% RDF through Neem cake + Microbial consortia and 53.86 g in treatment with 50% RDF through FYM + 50% RDF through Neem cake + Microbial consortia. Kalmegh plant grown without any organic manure or combination of organic manure recorded lowest fresh weight (30.01 g). The data is in conformity with the results of Saeidnejad and Rezvani (2010) in Cumin where highest yield was noticed in plants treated with sheep manure, compost, vermicompost and cow dung manure

### **Dry weight of plant (g)**

Effect of different organic manures along with microbial consortia on dry weight of Kalmegh is presented in Table 2 and depicted in Figure 2.

Different organic manures along with microbial consortia influenced significantly on dry weight of plant. The highest dry weight of 21.49 g was recorded in plant supplied with 50% RDF through Sheep manure + 50% RDF through Neem cake followed by 20.58 g in plants provided with 50% RDF through Poultry manure + 50% RDF through Neem cake + Microbial


consortia) and 20.06 g in the treatment with 50% RDF through FYM + 50% RDF through Neem cake + Microbial consortia. Control treatment plants recorded least dry weight of 11.9 g. The results are in line with the experiments conducted by Saeidnejad and Rezvani (2010) in Cumin.

### **Herbage yield per plot (kg) and per hectare (q)**

Different organic manures along with microbial consortia influenced significantly fresh herbage yield at harvest and is presented in Table 2.

The results indicated that herbage yield per plot and per hectare was significant. The highest fresh herbage yield of 6.60 kg/ plot and 116 q/ ha was recorded in treatment supplied with 50% RDF through Sheep manure + 50% RDF through Neemcake which was found to be on par with treatment supplied with 50% RDF through Poultry manure + 50% RDF through Neem cake + Microbial consortia (6.46 kg/ plot and 113 q/ ha). The least values for fresh herbage was observed in control (3.36 Kg/ plot and 59.3 q/ ha).

This significant difference due to the application of sheep manure along with neem cake and poultry manure in combination with neem cake and microbial consortia might be due to the fact that the sheep manure enhances plant growth hormones, soil fertility and facilitate in availability of plant nutrients throughout the crop growth period whereas in poultry manure + neem cake + microbial consortia combination, poultry manure amends soil, neem cake facilitates plant nutrient uptake and stimulative activity of microflora in the rhizosphere leading to availability of nutrients and vigorous growth and it might have resulted in recording the higher herbage yield. Similarly, Kamal and Yousuf (2012) opined that among the various organic manures, Neem cake followed by poultry manure was superior to cow dung with regards to yield parameter in Turmeric. Wahab *et al.* (2016) indicated that the combination treatment of manure, compost with biofertilizer recorded increase in yield in Fennel crop



**Fig. 2: Effect of different organic manures along with microbial consortia on fresh and dry weight of the plant in kalmegh**

Treatment	Plant Height (cm)	Plant spread (cm <sup>2</sup> )	Number of branches/plant
T <sub>1</sub>	33.7	775.2	10
T <sub>2</sub>	48.2	1160.2	12.4
T <sub>3</sub>	44.8	1086.2	12.4
T <sub>4</sub>	41.9	1475.8	13.67
T <sub>5</sub>	49.4	1955.4	13.6
T <sub>6</sub>	43.2	840.2	11.6
T <sub>7</sub>	43.8	1108.8	14
T <sub>8</sub>	41.1	1046.6	10.8
T <sub>9</sub>	43.3	1372.4	13.6
T <sub>10</sub>	43.9	2209.6	10.8
T <sub>11</sub>	41.0	1137.2	11.33
T <sub>12</sub>	42.0	847.0	13.5
T <sub>13</sub>	45.8	804.2	13.1
<b>'F' test</b>	*	*	*
<b>S Em±</b>	<b>0.27</b>	<b>31.7</b>	<b>0.09</b>
<b>CD (P=0.05)</b>	<b>0.80</b>	<b>92.8</b>	<b>0.28</b>

**Table 1: Influence of different organic**

**anic manures along with microbial consortia on yield parameters of Kalmegh**

Treatment	Fresh weight of plant (g)	Dry weight of plant (g)	Herbage yield / plot (kg)	Herbage yield / hactare (q)
-----------	---------------------------	-------------------------	---------------------------	-----------------------------

**\*Significant at 5% level**

**Table 2: Inf**

**luence of different organic manures along with microbial consortia on yieldparameters of Kalmegh**

T <sub>1</sub>	30.01	11.9	3.36	59.3
T <sub>2</sub>	41.7	14.72	4.24	75.0
T <sub>3</sub>	38.52	14.52	4.34	76.0
T <sub>4</sub>	38.9	14.64	4.39	77.0
T <sub>5</sub>	44.11	17.5	5.05	89.0
T <sub>6</sub>	38.62	16.9	4.8	85.0
T <sub>7</sub>	56.60	21.49	6.6	116.0
T <sub>8</sub>	46.85	17.5	5.3	95.0
T <sub>9</sub>	53.86	20.06	6.28	110.0
T <sub>10</sub>	47.47	17.75	5.47	96.0
T <sub>11</sub>	41.50	15.95	4.85	85.0
T <sub>12</sub>	54.20	20.58	6.46	113.0
T <sub>13</sub>	53.4	19.91	6.22	109.8
<b>'F' test</b>	*	*	*	*
<b>S Em±</b>	<b>0.56</b>	<b>0.20</b>	<b>0.07</b>	<b>1.25</b>
<b>CD (P=0.05)</b>	1.65	<b>0.59</b>	<b>0.20</b>	<b>3.6</b>

results obtained in the experiment, it could be concluded that plants supplied with sheep manure along with neem cake was better to get maximum growth and yield when compared to other treatment combinations and can be suggested to farmers for better growth and yield returns of Kalmegh.

## REFERENCES

1. Chandana, M., Joshi, V., Vijaya, D. and Lakshminarayana, D. (2018). Effect of organic treatments and spacing on growth parameters of Kalmegh (*Andrographis paniculata*) var. Cim-megha. *Journal of Pharmacognosy and Phytochemistry*, 7(6): 1695-1699.
2. Chiramel, T., Bagyaraj, D.J. and Patil, C.S.P.(2006). Response of *Andrographis paniculata* to different arbuscular mycorrhizal fungi. *Journal of Agricultural Technologies*, 2(2): 221-228.
3. Detpiratmongkol, S., Ubolkerd, T. and Yoosukyingstaporn, S.(2014). Effects of chicken, pig and cow manures on growth and yield of Kalmegh (*Andrographis paniculata* Nees). *Journal of Agricultural Science and Technology*, 10(2): 475-482.
4. Gupta, M., Srivastava, P.K., Niranjana, A. and Tewari, S.K.(2016). Use of a bioaugmented organic soil amendment in combination with gypsum for *Withania somnifera* growth on sodic soil. *Pedosphere*, 26(3): 299-309.

5. Kumar, B. and Topal, D., (2015). Effect of Organic Fertilizers on the Growth of Shoot of Kalmegh (*Andrographis Paniculata*). *International Journal of Novel Research in Life Sciences*, 2(2): 9-11.
6. Kumari, A.(2011). Vermicompost and FYM Effects on Biogeochemical Performances of *Andrographis paniculata* Nees. *Asian journal of chemistry*, 23(10): 4641-4644.
7. Nyeem, M.A.B., Mannan, M.A., Nuruzzaman, M., Kamrujjaman, K.M. and Das, S.K. (2017). Indigenous king of bitter (*Andrographis paniculata*): A review. *Journal of Medicinal Plants Studies*, 5(2), pp.318-324
8. Patel, D.H., Hirpara, B.V., Patel, S.A., Panchal, D.B. and Parul, M.(2012). Response of different organic manures and spacing on growth, yield, quality and economics of Kalmegh panchang under loamy sand of inceptisol of Anand. *Asian Journal of Soil Science*, 7(2): 312-314.
9. Premalatha, K., Botlagunta, N., Santhosh, D., Hiremath, C., Verma, R.K., Shanker, K., Sundaresan, V. and Kalra, A.(2021). Enhancement of soil health, germination and crop productivity in *Andrographis paniculata* (Burm. f.) Nees, an important medicinal crop by using a composite bio inoculant. *Journal of plant nutrition.*, 44(15): 2331-2346.
10. Rajeswara Rao, B.R. and Rajput, D.K. (2005). Organic farming in medicinal and aromatic plants. In: *Compilation note of winter school on organic farming - A step towards Eco-farming for sustainable agriculture*, July 4-24, 2005, held at Dept. Agron., College of Agri. ANGRAU, Hyderabad, India. pp. 66-77.
11. Saeidnejad, A.H. and Rezvani moghaddam, P.(2010). Investigation the effect of compost, vermicompost, cow and sheep manures on yield, yield components and essence percentage of cumin (*Cuminum cyminum*). *Journal of horticultural science.*, 24(2): 142-148.
12. Samanhudi, A.Y., Pujiasmanto, B. and Rahayu, M.(2014).Effect of organic manure and arbuscular mycorrhizal fungi on growth and yield of young ginger (*Zingiber officinale* Rosc.). *IOSR Journal of Agriculture and Veterinary Science*, 7(5): 1-5.
13. Tejasree , Kota, Narute T. K., Navale A. M., and Khaire P. B. 2024. "Bioconsortia: A Potential Ecofriendly Bio-Based Product to Enhance the Productivity of the Green

Gram". *Journal of Advances in Biology & Biotechnology* 27 (2):159-70.  
<https://doi.org/10.9734/jabb/2024/v27i2707>.

#### 14. More Citation Formats

15. Li, X., Li, P., Lin, X., Zhang, C., Li, Q., & Gong, Z. (2008). Biodegradation of aged polycyclic aromatic hydrocarbons (PAHs) by microbial consortia in soil and slurry phases. *Journal of hazardous materials*, 150(1), 21-26.

16. Behera, B., Das, T. K., Raj, R., Ghosh, S., Raza, M. B., & Sen, S. (2021). Microbial consortia for sustaining productivity of non-legume crops: prospects and challenges. *Agricultural Research*, 10, 1-14.