

Demonstration on New Rice Variety Co 52 to Improve the Productivity in Cuddalore District of Tamil Nadu.

ABSTRACT

Selection of high yielding variety is an important factor in rice cultivation for a particular location/ environment. Cultivation of newly released rice variety has potential to increase the productivity and needs to be promoted and popularized. In this regard, the present investigation was carried out at farmer's field to improve the rice productivity in Cuddalore district by introducing new rice variety CO 52. The CO 52 blast resistant medium duration high yielding rice variety was released by Tamil Nadu Agricultural University during 2018. A total of seventeen demonstrations with blast resistant newly released high yielding rice variety Co 52 were conducted at farmer's field organized by Vegetable Research Station (TNAU), Palur, Cuddalore District during samba season of 2019-20. An average yield of 6,501 kg ha⁻¹ was recorded in Co 52 demonstration trials with 10.51 per cent increased from the farmers cultivating variety BPT 5204 (5,886.75 kgs/ha). The farmers additional revenue of Rs. 9000- Rs. 10,400/ha was obtained from Co 52 demonstration plots. In this regard one field day and training on improved production technologies were organized at farmer's field to increase the productivity of rice through demonstrations of new rice variety Co 52 with improved production technologies.

Comment [CT2]: (TNAU)

Comment [CT3]: 5887 kg ha⁻¹

INTRODUCTION

Rice (*Oryza sativa* L.) is the world's most important food crop and belongs to Poaceae. The "Global Grain" cultivated widely across the globe feeding millions of human beings. It serves as the staple food for more than half of the world's population [1] (Khan *et al.* 2013). Worldwide, it is grown on an area of 166.1 million

Comment [CT4]: It is called the "Global Grain" and ...

hectares with yield of 745.2 million tonnes. In India, rice ranks second in both area and production, and cultivated over 43.90 million hectares, yielding 114.45 million tonnes with a productivity of 2607 kg/ha [2].

It is grown under diverse soil and climatic conditions; the productivity level of rice is low compared to the productivity levels of many countries in the world. Also about 90 % of the cultivated land belongs to marginal, small and medium farmers which are another constrain in increasing the productivity of rice in the country. Therefore, there is ample scope to increase the productivity of rice in the country. The highest productivity is 6710 kg per ha in China followed by Vietnam (5573 kg /ha), Indonesia (5152 kg/ha), Bangladesh (4375 kg/ha). There are improved technologies and introduction of new high yielding variety which could be adopted to increase the productivity in the country.

Production and productivity of rice was mainly depended on choice of varieties, season and agronomic practices with application of balanced major nutrition[3]. Among the components, selection of varieties play an important role in farming communities. Hence it is imperative to popularize the new high yielding varieties to replace the deteriorating commercial varieties so that overall productivity can be stabilized. Therefore, to meet the immediate needs of the rice farming community, there is a need to popularize the high yielding variety with good agricultural practices to meet the challenges in rice cultivation [4]. Cultivation of newly released blast resistant rice variety has potential to increase the productivity and needs to be promoted and popularized. Keeping in this regard, the present study was conducted at farmers field by introducing the newly released rice variety Co 52.

2. MATERIALS AND METHODS

2.1 Experimental Materials and Site

The experimental materials used in the present demonstrations are newly released rice variety CO 52 (MGR 100) and check variety BPT 5204. The field demonstration trials were conducted at Farmers field in Cuddalore District, Tamil Nadu, India during 2019-20. The soil of the experimental plot was clay loam with pH 7.0-

7.5 and low in organic carbon and total N content but medium in soil available P_2O_5 and K_2O . The climate of research locations is tropical. The maximum temperature recorded was $33.5^\circ C$ in the month of August while the minimum temperature recorded was $20.1^\circ C$ during winter peak (November). Average rainfall of the region is 1000-1100 mm per annum and relative humidity ranges from 45-85 per cent.

Comment [CT5]: Define low

Comment [CT6]: What is medium?

2.2 Experimental Layout and Crop Monitoring

The Co 52 seeds were distributed to farmers at no cost for one acre. The farmers are advised to raise the new variety (Co 52) along with ruling variety BPT 5204 used check variety in samba season of Tamil Nadu. The selected farmers were trained for TNAU Improved production technologies through training programme funded by NADP project and organized by Vegetable Research Station (TNAU), Palur during 2019-20. The seeds of the selected varieties were sown on the nurseries and seedlings (on 22nd to 27th days old seedling) were transplanted to the main field. Spacing was followed in 20×15 cm for both the varieties. All the agronomic and plant protection measures were followed in all the demonstrations and control plots uniformly. The observations were recorded on number of productive tillers per plant and grain yield per hectare ($kg\ ha^{-1}$). Ten to fifteen representative plants in each demonstration plots in all the farmers' field were taken for Co 52 as well as check variety BPT 5204. All the collected data were statistically analyzed by statistical method described by [5].

Comment [CT7]: It would be better describing the method or the authors here.

RESULTS AND DISCUSSION

The performance of CO 52 rice variety at farmers field compared to the farmers cultivating variety BPT 5204 (check variety) are given in Table 1. The data on number of productive tillers per plant revealed that, it was ranged from 17.93 to 23.98. The average of number of tillers in Co 52 demonstrations was 20.58 and the check variety (BPT 5204) was recorded in 18.79. The tillering potential of the variety directly contributes to grain yield. Number of tillers in rice was already reported by [6, 7]. Regarding the grain yield in Co 52 rice demonstration fields, the maximum grain yield of 6670 kg/ha was observed in Cuddalore block farmer's field and minimum yield 6310 kg/ha was recorded in Annagram Block. The average grain yield of all

Comment [CT8]: You can't have .93 or .98 Tillers. Use 18 to 24.

Comment [CT9]: 21

Comment [CT10]: 19

Comment [CT11]: It is difficult to go to references and return to continue to read the article. Best is if names of authors appear here.

Comment [CT12]: Uniformize the citation ($Kg\ ha^{-1}$) across the text.

Comment [CT13]: Idem

demonstration 6501.47 kgs/ha was recorded for Co 52 demonstrations at farmers field and for BPT 5204 variety, it was 5886.75 kgs/ha. It was a 10.51 % increase over the check variety (BPT 5204). These outcomes are somewhat comparable to [8]. The grain yield on rice was already reported in their research papers by [9]. The farmers getting additional revenue of Rs.9,000 to 10,400/ha by cultivating the new high yielding rice variety Co 52 (MGR 100). These findings align with those of [10, 11]. The additional yield and income is due to cultivating new high yielding rice variety along with improved production technologies. Similar kind of front line demonstrations in rice was already reported by Mohammad Hashim *et al.* [12] and Mandavkar *et al.* [13]. The Co 52 rice variety produced higher yield over the check variety (BPT 5204) in all the demonstrations, clearly indicated that showing constant performance in different locations, the Co 52 was easily adopted to new environments and having high stability over the locations in northern district of Tamil Nadu. Any new variety giving stable performance in different locations was good shine for farming [15].

Comment [CT14]: 6501,5 kg ha⁻¹

Comment [CT15]: What is this?

CONCLUSION

Pests and diseases controlled by chemical methods are not being a preferred option in staple food production. Varieties with inbuilt resistant/tolerant to the existing and emerging pests and diseases are the foundation of stress management strategies. Rice blast is an important disease in this crop, which can lead to substantial economic losses, in the absence of an efficient management strategy. Tamil Nadu, an important rice growing state in the country faces several abiotic and biotic stresses and this necessitates location specific rice variety for the zone. The cultivation of improved high yielding varieties like Co 52, along with suitable improved innovative technological interventions can be an important step in this direction. This high yield medium duration rice variety Co 52 with its excellent performance in the demonstrations at Cuddalore District, will play a significant role in improving the productivity, profitability and sustainability of rice cultivation in Tamil Nadu.

REFERENCES

1. Khan AS., Imran M, Ashfaq M. Estimation of Genetic Variability and Correlation for Grain Yield Components in Rice (*Oryza sativa* L.). *American-European J Agric Environ Sci.* 2013;6(2):585–590.
2. Agricultural Statistics at a glance. Government of India, Ministry of Agriculture & Farmers Welfare, Department of Agriculture & Farmers welfare economics & statistics division. 2022.
3. Ganapathy S, Jayakumar J. Evaluation of sugarcane (*Saccharum spp.* hybrids) clones for yield, quality, and its contributing traits. *J. Experimental Agric. Int.* 2023;45(7):113-118.
4. Ganapathy S, Ravichandran V, Jayakumar J. Yield, Quality and disease resistance of sugarcane clones. A Field Evaluation. *J. Experimental Agric. Int.* 2024;46(5):40-46.
5. Panse VG, Sukhatme PV. *Statistical Methods for Agricultural Workers.* ICAR, New Delhi. 1978.
6. Santhiya S, Pushpam R, Subramanian A, John Joel A, Menthol A. Nature of gene action and combining ability effects for grain yield and quality traits in rice (*Oryza sativa* L.). *Electronic J. Plant Breed.* 2024;5(1): 11- 20.
7. Aishwarya Duraiswamy, Sherina Jebakani, Lydia Pramitha K, Ramchander J, Devasena S, Wilson N, Dinesh Kumar P, Ramesh Kumar P. Evaluating the variability parameters among rice (*Oryza sativa* L.) Land races and varieties from Tamil Nadu. *Electronic J. Plant Breed.* 2024;14(2):187- 495.
8. Najeeb S, Sheikh FA, Parry GA, Shikari AB, Zaffar G, Kashup SC, Ganie MA, Shah AB. Farmers participatory selection of new rice varieties to boost production under temperate agro-ecosystems. *J. integrative Agric.* 2018; 17(16): 1307-1314.
9. Vaishnavi Pravin Gupte, Manonmani S, Nivedha, R, Suresh R, Senthil Kumar G, Raveendran M. Genetic diversity studies and identification of donors for lodging resistance in rice (*Oryza sativa* L.). *Electronic J. Plant Breed.* 2023; 14(3): 1158-1166.

10. Singh T, Singh R, Soni RL. Performance of rice variety P 1460 in Front line demonstrations under rainfed conditions in Southern humid region of Rajasthan. Annals of Agricultural Research. New Series. 2012; 33(3): 121-125.
11. Vikram Sigh Gaur, Uttam Kumar Bisen, Naresh Kumar Bisen, Ramakrishna Solanki. Evaluation of Rice varieties under Front Line Demonstrations in the Agro-climatic Zone of the Chattisgarh Plains Madhya Pradesh, under Irrigated condition. J. Exp. Agric. Int. 2024; 46(3): 156-161.
12. Mohammad Hashim, Singh KK, Narendra Kumar, Man Mohan Deo, Dileep Singh, Mukund Kumar. Impact of Front Line Demonstrations in Improving Rice Productivity and Profitability in Jaunpur District of Uttar Pradesh, India. J. Experiment. Agric. International. 2023; 45(11): 74-82.
13. Mandavkar P.M, Sawant PA, Mahadik. Evaluation of Front Line Demonstration Trials on Rice in Raigad District of Maharashtra. J. Extension. Edu. 2012; 20: 4-6.
14. Abu-Elail, F.F.B., El-Taib ABA, Hefny AM. Stability analysis and performance of promising sugarcane varieties for yield and quality traits. Electronic J. Plant Breed. 2021; 12(3): 623- 636.

Table 1. Performance of Co 52 (MGR 100) Rice variety under farmers field Demonstrations at Cuddalore District.

S. No.	Farmers Name & location	No. of tillers / plant			Grain Yield / ha.		
		Co 52	BPT 5204	% increase	Co 52	BPT 5204	% increase
1.	R. Narayanan, Cuddalore Block	21.55	19.34	11.43	6350	5950	6.72
2.	P. Sudhagar Cuddalore Block	18.95	17.21	10.11	6630	6100	8.69
3.	R. Govindhan, Cuddalore Block	23.98	22.33	7.39	6400	5780	10.73
4.	R. Balasubramani, Cuddalore Block	23.08	20.33	13.53	6475	5620	15.21
5.	J. Arunmozhidevan, Kunjipadi Block	21.65	19.25	12.47	6520	5860	11.26

6.	V. Raghavendhar, Cuddalore Block	18.58	17.85	4.09	6670	6250	6.72
7.	V. Mohan Kurinchipadi Block	19.48	18.57	4.90	6470	5750	12.52
8.	V. Sundramurthy, Kurinchipadi Block	17.93	16.71	7.30	6510	5980	8.86
9.	M. Narayanan, Kurinjipadi Block	19.03	17.23	10.45	6450	5680	13.56
10.	C. Prabhu, Annagram Block.	21.22	18.86	12.51	6570	5870	11.93
11.	N. Anbu, Annagram Block	19.35	17.63	9.76	6310	5920	6.59
12.	D. Janakiraman, Annagram Block	23.20	20.75	11.81	6650	5850	13.95
13.	V. Raja, Annagram Block	20.95	19.32	8.44	6580	5850	12.48
14.	Sumathi Nadarajan, Annagram block	19.45	17.42	11.65	6560	5815	12.81
15.	R. Vaithyanathan, Annagram Block	20.53	18.25	12.49	6540	5730	13.74
16.	Kaliyavarathan Kurichipadi Block	21.33	19.81	7.67	6470	6120	5.72
17.	V.Subramani Kurichipadi Block	20.65	18.01	14.66	6350	5950	6.72
	Grand Mean	20.58	18.79	10.05	6501.45	5886.75	10.51
	CD (0.05%)	2.50	2.42	-	577.55	537.41	-
	CV (%)	5.76	6.04	-	6.31	5.72	-

Fig. 1, 2. Performance of Co 52 (MGR 100) Rice variety under farmers field



Fig. 1. Demonstration of Co 52 (MGR 100) at Farmers field



Fig. 2. Demonstration of Co 52 (MGR 100) at Farmers field

UNDER PEER REVIEW