

Original Research Article

Prevalence and Related Risk Factors of *Entamoeba* complex infection among Asymptomatic School Children in the Ho Municipality, South-Eastern Ghana.

ABSTRACT

Background: Amoebiasis is a common cause of gastroenteritis in children, with the *Entamoeba* complex consisting of genetically diverse but morphologically identical species. Amoebic infections are generally subclinical and may be acute or chronic, leading to high morbidity or mortality, especially among children.

Objective: This study was designed to determine burden of *Entamoeba* complex and other enteric parasites among asymptomatic school children within the Ho Municipality of the Volta Region in Southeastern Ghana.

Methodology: This was a cross-sectional study, involving 302 primary school children (aged 4–15 years) in 6 socio-economic settings within the municipality. Single fresh stool specimens were collected from the children and examined microscopically using direct wet mount, iodine preparation and formal ethyl-acetate concentration technique. Structured questionnaire was used to determine demographics and risk factors associated with *Entamoeba* complex infection among the school children.

Results and Discussion: A total of 106 (35.1%) children had gastrointestinal parasitic infection. Of these, seventy-two (23.8%) had *Entamoeba* complex infection with the highest infection (30.4%) recorded among 8–9-year-olds. Children from Peri-Urban communities were more infected with *Entamoeba* complex (32.4%, 33/102) compared to Urban (24.8%, 25/101) and Rural (14.1%, 14/99) respectively. Other parasites detected included; *Giardia lamblia* 21(7.0%), Hookworm 11(2.3%), *Trichuris trichiura* 1(0.3%) and *Ascaris lumbricoides* 1(0.3%). Factors associated with *Entamoeba* complex infection were source of food, fingernails biting/thumbs sucking habits and mother's level of education.

Conclusion: Parasitic infections is prevalent in the study area and we recommend personal and environmental hygiene practices as well as health education for effective elimination of amoebic and other enteric parasitic infections.

Keywords: *Entamoeba complex infections; school children; helminthes; Giardia lamblia.*

1. INTRODUCTION

Amoebiasis is a type of gastroenteritis caused by the parasite *Entamoeba histolytica*, commonly occurring among immigrant populations, travelers, immunocompromised individuals, those residing in mental care homes, prisoners, children in day care centres and their families, and sexually active homosexuals [1-6]. Besides *E. histolytica*, about seven (7) other amoebic organisms- *Entamoeba dispar*, *Entamoeba coli*, *Entamoeba moshkovskii*, *Entamoeba polecki*, *Entamoeba hartmanni*, *Endolimax nana* and *Iodamoeba bütschlii* have been identified [7-10]. That notwithstanding, *Entamoeba polecki*, *Iodamoeba bütschlii*, an additional amoebic organism, *Dientamoeba fragilis*, are occasionally implicated in human diarrhoea cases [11,12]. Moreover, *E. dispar*, *E. moshkovskii*, and the predominantly pathogenic amoeba - *E. histolytica* together constitute what has been known as the Entamoeba complex, as they are morphologically identical to each other, albeit genetically diverse [8,10,11]. However, of these three, only *E. histolytica* is known to be invasive [8, 12].

Amoebiasis is largely transmitted through food or water contaminated with human faeces containing the hardy infective amoebic cyst [13-14]. Hence, improper sanitation and poor personal hygiene practices, which permeate developing countries, are major risk factors of amoebic infection, explaining the disproportionately higher amoebiasis burden in these regions [15-19]. The Sustainable Development Goal (SDG) 6 of the United Nations therefore emphasizes the provision of water, sanitation and hygiene facilities as a means of reducing the burden of amoebiasis and other sanitation-linked [20,21]. Sadly, although more than 90% of the world's population has access to sources of drinking water, poor water quality remains a major challenge, especially in sub-Saharan Africa [19,21]. This has frustrated efforts targeted at reducing the high morbidity and mortality associated with the *Entamoeba* complex and other enteric parasites [18, 22].

Global prevalence of *Entamoeba* complex infection shows that infected asymptomatic individuals make up about 90% of the total infected persons, whereas the remaining are symptomatic [23,24]. Out of the 10% symptomatic individuals, 90% develop intestinal amoebic colitis while the remaining develop extra-intestinal amoebiasis [25]. This accounts for about 50-100 million reported cases of amoebic

colitis or liver abscesses yearly resulting in almost 100,000 deaths [26]. Amoebiasis is next to malaria in relation to mortality from protozoan infection [25, 26]. Clinical presentations of amoebic infection include pain in the abdomen, diarrhoea, dysenteric stools, bloody or mucoid stools, ameboma, toxic mega colon, colonic perforations, peritonitis, amoebic cecitis and appendicitis, cutaneous amoebiasis, rectovaginal amoebic cutis, hemorrhage among other complications [27]. Other clinical manifestations may involve extra-intestinal organs such as the liver. The most common clinical presentation of invasive amoebic infection is the potentially fatal amoebic liver abscess which affects organs such as the kidney, heart, brain, genitourinary tract and pleura pulmonary [28]. Generally, there is a poor understanding of amoebic infections in sub-Saharan Africa[29]. There is a paucity of information on amoebic infections among school children in the Ho Municipality in Southeastern Ghana. Therefore, this study aims to determine the burden of study *Entamoeba* complex and related factors among asymptomatic school children within the municipality.

2. MATERIALS AND METHODS

2.1 Study Area

This study was undertaken within the Ho Municipality, located in the central part of the Volta Region, which lies within the South-Eastern part of Ghana. It is bounded by the Republic of Togo on the east, Adaklu, Agortime-Ziope Districts to the south, and Ho West District to the west and Hohoe Municipality to the north. The municipality has Ho as its capital, occupying a total land mass of 2,361km² [30]. The estimated population of Ho municipality is 180,420, consisting of 84,843 males and 95,577 females, according to the 2021 Ghana Population and Housing Census report. Approximately 125,914 people, made up of 59,254 males and 66,660 females, lived in the urban areas of the Municipality. However, 54,506 people reside in the rural areas of the municipality, of whom 25,589 and 28,917 are males and females, respectively [31].

The study was conducted in six primary schools located in three socio-economic settings classified as: rural (Klefe Achatime and Klefe Demete Evangelical Presbyterian Primary Schools), peri-urban (Sokode Gbogame Municipal Authority and Sokode Gbogame Roman

Catholic Primary Schools), and urban (Ho Bankoe Evangelical Presbyterian Primary and Ho SSNIT Flats Presby Primary Schools) all within the Ho Municipality of the Volta Region (**Figure 1**).




Figure 1: Study Sites

2.2 Study design, population and sampling

This was a cross-sectional study that involved 302 children aged 4-15 years sampled from six primary schools in the municipality from October 2016 to March, 2017. The study involved the administration of standardized questionnaires to collect demographic and socio-economic data on the children and their

parents or legal guardians in English and the local (EWE) language. Briefly, the sex, age, socio-economic settings, source of food, drinking water, fingernails biting /thumb sucking habit, type of toilet facility at home were obtained from the children as well as the educational level of their mothers were obtained. Written Informed Consent was obtained from parents or legal guardians of the school children and assent forms were duly signed by the children before they were enrolled in the study. Interviews were conducted in the local language (Ewe) for pupils who could not communicate effectively in the English language. All interviewers had been trained to avoid interviewer-bias. The minimum sample size was 255, calculated using the Cochran formula [57] at 95% confidence interval (CI), with 5% margin of error with 21% prevalence of *Entamoeba* complex infection in the Ho Municipality of the Volta Region [32,33].

2.3 Stool Specimen Collection

A clean wide-mouthed leak-proof stool containers, screw-capped fitted with scoop and clearly labeled with unique respective specimen number, date and location was provided to each child who had consented to be part of the study. Each child was carefully instructed on how to collect the specimen (to circumvent urine contamination and to fill about a third of the container with fresh stool specimen) in the privacy of the school's toilet facility. Thorough hand washing with soap under running water was ensured to prevent possible infection and re-infection. Single fresh stool specimens were collected from the children each school day, and were immediately transported for laboratory investigation.

2.4 Laboratory investigation

The consistency of the stool was observed and recorded as formed, semi-formed, loose, mucoid, slimy or watery. Evidence of parasitic infection was based on the identification of cysts, trophozoites, ova, larvae or adult worms from direct saline wet mount [34-35], iodine preparation and formol-ethylacetate sedimentation technique [35-37]. Low power 10X and high power 40X objective lenses were used to examine the stool samples for the identification of enteric parasites as described elsewhere [34]. In all, three microscopists examined the slides. Two independently, observed the slides and a third was called on to resolve any discordant results. A high power 40X objective was used to observe the morphological features of any motile organism (protozoan trophozoites) or larvae detected [35].

2.6 Data processing and analysis

Data from interviews and parasitic investigations were entered into Microsoft Excel 2016, and exported into Statistical Products and Services Solutions (SPSS, version 26) for statistical analyses. The data were summarized by determining the frequency distribution of amoebic infection among the sample population. The correlation of amoebic infection between males and females, age groups and risk factors were determined using chi-square test analysis and odds ratios at a 95% confidence interval. A p -value < 0.05 was considered statistically significant.

3. RESULTS

A total of 302 Primary School children aged 4 to 15 years (Mean, 50.3; SD,34.1) in six (6) different communities participated in the study. One hundred and fifty-three (50.7%) were males, whilst the remaining 149 (49.3%) were females indicating an evenly distributed study population in terms of gender. The thirty-one percent of the children who regularly consumed food bought from the street or school canteen, 25.5% bit their fingernails, 18.2% practised open defecation while 23.3% used pipe-borne water as their main source of water supply, were all infected with *Entamoeba* complex. Nearly equal number of males and females were between the ages 12-13 (26.5%) and 10-11 (26.2%) respectively with the least age group being 4-5 (1.3%).

3.1 Parasitological profiles of School Children

Of the 302 stool specimens examined, 106 (35.1%) of them had gastrointestinal parasitic organisms. The overall prevalence of *Entamoeba* complex infection was 72 (23.8%). Four other parasites were identified: *Giardialambliia* 21(7.0%), Hookworm 11(2.3%), *Trichuristrichuira* 1(0.3%) and *Ascaris lumbricoides* 1(0.3%). Of those infected, 94 (31.1%) had protozoan infections, while the remaining 12 (4.0%) had helminth infections. One-hundred and four (34.4%) of those had mixed infections, while the remaining 2 (0.7%) had single infections. Mixed infections involving a combination of protozoan organisms (*Entamoeba* complex and *Giardia lamblia*) were higher (93:30.8%) than co-infections involving *Entamoeba* complex and helminths (83:27.5%). No co-infection involving helminthic parasites was

detected. The prevalence of parasitic infections in the various socio-economic settings was as follows: Urban (25: 24.8%), Peri-Urban (33:32.4%) and Rural (14: 14.1%) (**Figure 2**).




Figure 2. Frequency of gastrointestinal parasites detected by microscopy

3.2 Parasitic Infection among different age groups

Overall, *Entamoeba* complex infection peaked in children aged 8-9 years (30.4%) compared to other age groups (**Table 1**). However, there was no significant association between *Entamoeba* complex infection and age of the participants ($p = 0.071$) (**Table 1**).

Table 1. *Entamoeba* complex infection in Different Age Groups.

Age group (years)	Stool examined	No. infected	X^2	p value	OR	95% CI
4-5	4	1(25)				0.182-2.568
6-7	59	10 (16.9)				1.013-1.292
8-9	69	21(30.4)				1.226-1.499

10-11	79	15(19.0)	15.770	0.071	0.264	1.114-1.380
12-13	80	23(28.8)				1.116-1.359
14-15	11	2(18.2)				1.413-2.041

3.3 Entamoeba complex Infection in Different Socio-economic settings

Children in Peri-Urban areas (32.4%) had more parasitic infections compared to those in the Urban (24.8%) and Rural areas (14.1%) respectively. **(Figure 3).**




Figure 3. *Entamoeba* complex Infection in Different Socio-economic settings

3.4 Parasitic Infection and associated risk factors

Thirty-one percent of children who purchased food from the school canteen or the street were had *Entamoeba* complex infection compared to those who went home during break periods to eat **(Table 2)**. There was significant association ($p = 0.002$) between source of food and *Entamoeba* complex infection among the school children **(Table 2)**. In this study, children who either bite their fingernails or suck their fingers or thumb regularly (25.5%) had *Entamoeba* complex infection. Statistically, there was significant association ($p = 0.022$) between the age of participants and fingernails biting or thumb sucking behaviour

among the children (**Table 3**). Highest rate of *Entamoeba* complex infection was recorded among children whose mothers had secondary education (95.8%) (**Table 4**). The Association between *Entamoeba* complex infection and mother's educational level was statistically significant ($p = 0.007$). An *Entamoeba* complex infection (37.8%) was reported in children who used KVIP although there was no significant association ($p = 0.389$) between *Entamoeba* complex infection and the type of toilet facility used by the children (**Table 4**).

Table 2. Association of *Entamoeba* complex infection and source of food among the children

Variable (Source of food)	Examined (n)	Infected (n) %	X^2	p value	OR	95% CI
Home	126	18(14.3)				0.356-0.418
Street/School canteen	174	54(31.0)	6.249	0.002	0.576	0.469-0.558
Friends	2	0(0.0)				
Total	302	72(23.8)				

Table 3. Association of *Entamoeba* complex infection and fingernails biting/thumbs sucking behaviour among the children

Variable (Biting of fingernails)	Examined (n)	Infected (n) %	X^2	p value	OR	95% CI
Yes	98	25(25.5)				0.375-0.483
No	168	38(22.6)	5.972	0.022	0.556	0.449-0.532
Sometimes	36	9(25.0)				0.352-0.457
Total	302	72(23.8)				

Table 4. *Entamoeba* complex infection and other related risk factors

Variable (Mother's level of education)	Examined (n)	Infected (n) %	X ²	p value	OR	95% CI
BECE	222	28(12.6)				0.414-0.481
SSCE	24	23(95.8)				0.377-0.501
Tertiary	18	17(94.4)	2.776	0.007	0.735	0.272-0.421
No Formal education	38	4(10.5)				0.488-0.717
Total	302	72(23.8)				

Variable (Toilet facility at home)	Examined (n)	Infected (n) %	X ²	p value	OR	95% CI
KVIP	37	14(37.8)				1.062-1.452
Water Closet	97	21(21.6)	4.937	0.389	0.397	1.252-1.480
Pit latrine	48	15(31.3)				1.050-1.366
Open defecation	120	22(18.3)				1.122-1.328
Total	302	72(23.8)				

Variable (Source of drinking water)	Examined (n)	Infected (n) %	X ²	p value	OR	95% CI
Pipe-borne	283	66(23.3)				0.434-0.495
Well/Bore hole	2	2(100)	3.562	0.07	0.953	0.250-0.433

Stream/River	1	0(0.0)	-2.189-3.1805
Sachet	16	4(25.0)	-0.185-1.085
Total	302	72(23.8)	

4. DISCUSSION

Entamoeba complex infection is a common parasitic infection among children in developing nations including Ghana, due to poor personal hygiene practices, sanitation and unsafe water [38--40]. This study determined the prevalence and risk factors associated with *Entamoeba* complex infections in the Ho Municipality in the Volta Region of the Republic of Ghana. In the present study, *Entamoeba* complex was the most common intestinal protozoan parasite identified (23.8%) among all the asymptomatic children sampled in the Ho Municipality of the Volta Region, Ghana. The high prevalence of *Entamoeba* complex infection in the study area could be due to the poor personal and environmental hygiene practices in the study area [39]. Similar studies conducted by Simon-Oke and Ogunleye [41] recorded a high prevalence of (67.6%) 188/278 among primary school children in Akure, Nigeria, whereas Walanaet *al.*, [42] reported a low prevalence of (0.21%) 5/2400 among primary school children in Kumasi, Ghana. Also, Verweijetal., [32] reported a prevalence of (39.8%) 98/246 in the Bawku Municipality of the Upper East Region of Ghana. The asymptomatic children become carriers (cyst excreters) and could serve as points of transmission and infection to other uninfected children [43,44].

The sex of the study participants was not statistically associated with *Entamoeba* complex infection. The *Entamoeba* complex infection was slightly higher in females (24.8%) compared to males (22.9%). Similar study conducted in North East India by Nath et al., [45] revealed that females had higher *Entamoeba* complex infection (15.2%) compared to males (11.5%). Although studies have linked gender-specific roles to *Entamoeba* complex infection, the present study has not established that [45,46].

In the present study, there was a reported high incidence of *Entamoeba* complex infection 31% (54/174) among children who either buy food from the street or school canteen compared to 14.3% (18/126) who

obtained their food from home. The poor personal hygiene of most food vendors, who may be naïve asymptomatic carriers poses a serious health risk to consumers especially children who patronize their foods. Many of such food vendors do not go for routine medical check-ups to ascertain by public health standards if they are medically fit to sell food to the general public especially children. This is corroborated by Ayeh-Kumi *et al.*, [39] who suggested that food vendors in Accra, Ghana, were possible sources of intestinal parasitic infection including *Entamoeba* complex as a result of lack of safe water, improper personal and environmental hygiene.

The study has associated fingernail biting or thumb sucking behaviours of some children with *Entamoeba* complex infection. Nearly a third of the study participants either bite their fingernails or suck their thumbs with associated health risks. This assertion is confirmed by the findings that *Entamoeba* complex cyst can survive days to weeks in moist external environment with the capability of causing infection [47]. Sofiana *et al.*, [48] also observed that fingernails biting was a possible source of *Entamoeba* complex infection since the viable cyst can survive under fingernails for up to 45 minutes. This negative habit could be stopped when parents, legal guardians, and teachers help educate the children and regularly inspect and trim the fingernails of these children.

It is assumed that the level of education of mothers will have a positive impact on the health of their children [49,50]. However, the findings of this study proved otherwise. High incidence of *Entamoeba* complex infection was reported in children whose mothers had secondary (95.8%) and tertiary education (94.4%) compared to those without any formal education (10.5%). Most educated mothers are career women who by the demands of their jobs may not have time to provide home-cooked food for their wards each school day. They may resort to buying food for their wards from the street or giving money to their wards to buy food from school canteen which might be a source of *Entamoeba* complex infection.

Lack of hygienic toilet facilities in the study area was a major risk factor in *Entamoeba* complex infection. *Entamoeba* complex infection was higher in children from the peri-urban (33%) and urban (25%) compared to the rural (14%) areas. This is due to lack of adequate toilet facilities in the peri-urban and urban areas compared to the rural areas. In these socio-economic settings, nearly a third of the school children practised open defecation or defecate in disposable plastic bags and throw away (“fly away”) into

a nearby bush or open space. Others also due to poverty could not pay to use the public toilet facility in the community. The study also revealed that some children defecate in holes dug with simple implements (cutlass or hoe) in the backyard of their homes and covered with soil and this was also commonly practiced in the peri-urban and rural communities. The shallow nature of these holes may be uncovered by free-range-domestic poultry birds or washed into homes or nearby water bodies by runoff when it rains posing a public health risk [20,51].

The level of infection recorded in children who use water closets (WC) may be attributable to the unavailability of water leading to insanitary conditions favourable for *Entamoeba* complex infection. Contaminated water closet flush handles, or seats pose a greater health risk to users. The present study agrees with the findings of Nath et al., [45] which suggested that those who used unhygienic toilet facilities had higher *Entamoeba* complex infection than those who used hygienic toilets. Therefore, proper sanitary conditions including toilet facilities are critical to ensuring the health of children against *Entamoeba* complex infection [20,39,41,51]

Water is very critical to the survival of children and when children are made to depend on unsafe water, they are at a higher risk of deadly diseases including amoebiasis and severe malnutrition. Furthermore, unsafe water and improper sanitation are also connected to stunted growth which leads to irreversible physical and cognitive damage and seriously hinders the children's academic performance [39,48]. Access to safe drinking water and proper sanitation is a human right and safe water and adequate sanitation are critical for the development and well-being of humans [48,50,51].

Safe drinking water is essential to promoting good health and wellness of humans. The present study revealed that *Entamoeba* complex infection was highest (100%) among children who regularly depended on well or bore hole as against sachet (25%) and pipe borne water (23.3%) respectively. The findings of this study agree with the assertion that, water from wells or springs consumed had higher *Entamoeba* complex infection [52-56] compared to other sources.

The present study also revealed the presence of other parasites such as, *Giardia lamblia*, *Trichuris trichuira*, Hookworm and *Ascaris lumbricoides*. Even though the Ghana Health Service and the Ho Municipal Education Directorate embarked on mass administration of Metronidazole and Albendazole to

all primary schools children in December 2015 prior to this study, it is unclear why the prevalence of gastrointestinal parasites (35.1%) and *Entamoeba* complex infection (23.8%) remain high.

LIMITATIONS OF THIS STUDY

Firstly, the study was conducted between October 2016 to March, 2017 and a lot might have changed since then. Secondly, the use of microscopy as the main diagnostic tool, which is the gold standard in resource limited-settings including Ghana could have affected the overall prevalence of the parasites detected.

5. CONCLUSION

This study reported a high prevalence of gastrointestinal parasites among the study participants with *Entamoeba* complex infection being the most common. More females were infected compared to males and children between 8-9 years had the highest rate of infection compared to other age groups. Source of food, fingernail biting or fingers/thumbs sucking habits, level of mothers' education, and socio-economic settings were related risks factors associated with *Entamoeba* complex infection. We recommend a much wider molecular study to differentiate between species or zymodemes of *Entamoeba* complex in the Ho Municipality. The local authorities and other relevant stake holders should intensify their efforts at providing safe water, sanitation and hygiene facilities (WASH) in areas where they are lacking. Also, public education on the prevention of parasitic infections in the municipality should be intensified.

List of abbreviations

E.P: Evangelical Presbyterian, R.C: Roman Catholic, UNICEF: United Nations Children's Education Fund, KVIP: Kumasi Ventilated Improved Pit, WC: Water closet

DATA AVAILABILITY

All data supporting the reported results have been included in the paper.

CONSENT AND ETHICAL APPROVAL

Written Informed Consent was obtained from parents or legal guardians of the school children following an explanation of the objective of the study. Further, assent forms were duly signed by the children before they were enrolled in the study. Laboratory examination of stool specimens was performed free of charge. Study participants who were positive for intestinal parasite were linked to nearby health centre for treatment. The study was conducted in accordance with the Declaration of Helsinki (1964). This study was reviewed and approved by the Ethical and Protocol Review Committee of the College of Health Sciences, University of Ghana, Accra, Ghana (Protocol ID no: CHS-Et/M.1- P 4.11/2016–2017). The Institutional Review Board of the Volta Regional Hospital (now Ho Teaching Hospital) as well as the Ho Municipal Directorate of Ghana Education Service have also approved this study.

REFERENCES

- 1 Skappak C, Akierman S, Belga S, Novak K, Chadee K, Urbanski SJ, Church D, Beck DPL. Invasive amoebiasis: A review of *Entamoeba* infections highlighted with case reports. *Can J Gastroenterol Hepatol*, vol. 28, no.7, pp. 355-359, 2014.
- 2 QuinnTC, StammWE, GoodellSE, MkrtychianE, BenedettiJ, and CoreyL. The polymicrobial origin of intestinal infections in homosexual men. *N Engl J Med*. 309:576–582, 1983.
- 3 BarwickR, UzicaninA, and Lareau S. Outbreak of amebiasis in Tbilisi, Republic of Georgia 1998. *Am J Trop Med Hyg*, vol. 67, pp. 623–631, 2002.
- 4 Braga LL, Gomes ML, and DaSilva MW. Household epidemiology of *Entamoeba histolytica* infection in an urban community in northeastern Brazil. *Am J Trop Med Hyg*, vol. 65, pp. 268–271, 2001.
- 5 Stauffer W, and Ravdin JI. *Entamoeba histolytica*: an update. *Current Opinion in Infectious Diseases*, vol. 16, pp. 479–485, 2003.
- 6 SamieA, Bakri Eand Ra'ed A. *Amoebiasis in the Tropics: Epidemiology and Pathogenesis*. *Current Topics in Tropical Medicine*, Dr. Alfonso Rodriguez-Morales(Ed.), ISBN: 978-953-51-0274-8, InTech, Available from: <http://www.intechopen.com/books/current-topics-in-tropical-medicine/amoebiasis-in-the-tropicsepidemiology-and-pathogenesis>, 2012

- 7 Shimokawa C, Kabir M, Taniuchi M, Mondal D, Kobayashi S, Ali IKM, Sobuz SU et al. *Entamoeba moshkovskii* Is Associated With Diarrhea in Infants and Causes Diarrhea and Colitis in Mice, *The Journal of Infectious Diseases*, vol. 206, no. 5, pp. 744–751, 2012
<https://doi.org/10.1093/infdis/jis414>
- 8 Sardar SK, Ghosal A, Haldar T, Maruf M, Das K, Saito-Nakano Y et al. Prevalence and molecular characterization of *Entamoeba moshkovskii* in diarrheal patients from Eastern India. *PLoS Negl Trop Dis*, vol. 17, no. 5, pp. 1-23.
- 9 Haque RIK, Ali M, Clark CG, and Jr. Petri WA. (1998). A case report of *Entamoeba moshkovskii* infection in a Bangladeshi child. *Parasitol Int*. vol. 47, pp. 201–202, 1998.
- 10 Leber AL, and Novak SM. Intestinal and urogenital amebae, flagellates, and ciliates. *In Manual of clinical microbiology*, 7th ed. p. 1391–1405 ASM Press, Washington, D.C. 1999
- 11 Chan F, Stewart N, Guan M, Robb I, Fuite L, Chan I. et al. Prevalence of *Dientamoeba fragilis* antibodies in children and recognition of a 39 kDa immunodominant protein antigen of the organism. *Eur. J. Clin. Microbiol. Infect. Dis*, vol. 15, pp. 950–954, 1996.
- 12 Cuffari C, Oligny L, and Seidman EG. *Dientamoeba fragilis* masquerading as allergic colitis. *J. Pediatr. Gastroenterol. Nutr*, vol. 26 pp. 16–20, 1998.
- 13 Stanley Jr SL. Amoebiasis. *Lancet*, vol. 361, pp. 1025-1034, 2003.
- 14 American Academy of Paediatrics. Amoebiasis. In: Pickering LK ed. *Redbook 2012 Report of the Committee on Infectious Diseases* 29th ed. Elk Grove Village, IL: American Academy of Pediatrics, pp 222-225, 2012
- 15 Ximénez C, Morán P, Rojas L, Valadez A, and Gómez A. Reassessment of the epidemiology of amoebiasis: State of the art. *Infect Genet Evol* .9:1023–1032, 2009
- 16 Haque R, Mondal D, Duggal P, Kabir M, Roy S, Farr BM, Sack RB et al. *Entamoeba histolytica* infection in children and protection from subsequent amoebiasis. *Infect Immun*, vol. 74, no. 2, pp. 904-909, 2006.
- 17 Chacon-Cruz E. Intestinal protozoa diseases. General hospital Tijuana, Mexico. Pages 23-28, 2009.

- 18 Asemahagn MA. Parasitic Infection and Associated Factors among the Primary School Children in Motta Town, Western Amhara, Ethiopia. *American Journal of Public Health Research*, vol. 2, no. 6, pp. 248-254, 2014.
- 19 United Nations Children's Fund (UNICEF). Annual Report 2015. UNICEF Division of Communication, 3 United Nations Plaza, New York, NY10017, USA. pubdoc@unicef.org www.unicef.org ISBN:978- 92-806-4843-0 Page 25, 2015
- 20 Giribabu D, Bharadwaj P, Sitiraju R, Burra M, Rao PP, and Reddy CS. Combating open defecation through community-led sanitation. *Dr Sulaiman Al Habib Med J*, vol. 1, no. 3-4, pp. 45-51, 2014.
- 21 Saleem M, Burdett T, Heaslip V. Health and social impacts of open defecation on women: a systematic review. *BMC Public Health*, vol. 19, pp.158, 2019.
- 22 Sharif BO, Ali ZR, Mohammed HM. Impact of *Entamoeba histolytica* on the human Body. *Int J Sci and Med Res*, vol. 2, no. 4. pp. 5-8, 2022.
- 23 MacFarlane RC and Singh U, Identification of differentially expressed genes in virulent and nonvirulent *Entamoeba* species: potential implications for amebic pathogenesis. *Infection and Immunity*, vol. 74, no. 1, pp. 340–351, 2006.
- 24 Yanagawa Y, Nagata N, Yagita K et al. Clinical features and gut microbiome of asymptomatic *Entamoeba histolytica* infection. *Clinical Infectious Diseases*, vol. 73, no. 9, pp. e3163– e3171, 2021.
- 25 Ayeh-Kumi PF, Karim IBM, Lockhart LA, Gil Christ CA, Petri WA, Haque R. *Entamoeba histolytica*: Genetic diversity of clinical isolates from Bangladesh as demonstrated by polymorphism in the serine-rich gene. *Exp. Parasitol*, vol.99, pp. 80-88, 2001
- 26 Stauffer W, Ravdin JL. *Entamoeba histolytica*: an update. *Curr Opin Infect Dis*, vol. 16, no. 5, pp. 479-85, 2003.
- 27 De Villiers JP, and Durra G, Amoebic abscess of the brain (Case Report). *Clin. Radiol*, vol. 53, pp. 307-309, 1998.

- 28 Amin OM. Seasonal prevalence of intestinal parasites in the United States during 2000; *Am.J. Trop. Med. Hyg*, vol. 66, pp.799-803, 2002.
- 29 Stauffer W, Abd-Alla M, and Ravdin JI. Prevalence and incidence of *Entamoeba histolytica* infection in South Africa and Egypt. *Arch Med Res*, vol. 37, no. 2, pp. 266–269, 2006.
- 30 <https://www.hma.gov.gh/> Ho Municipal Assembly. Accessed on January 20, 2024 at 5:00GMT
- 31 Ghana Statistical Service (2021). Population and Housing Census. https://census2021.statsghana.gov.ghhttps://statsghana.gov.gh/gssmain/fileUpload/pressrelease/2021%20PHC%20General%20Report%20Vol%203A_Population%20of%20Regions%20and%20Districts_181121.pdf accessed on January 19,2024 at 10:15 am]
- 32 Verweij JJ,Oostvogel F, Brienen EA, Nang-Beifubah A, Ziem A, Polderman AM. Short communication: prevalence of *Entamoeba histolytica* and *Entamoeba dispar* in northern Ghana. *Trop Med Int Health*, vol. 8, pp. 1153 – 1156, 2003.
- 33 Opintan JA, Ayeh-Kumi PF, Newman MJ, Assrim R, Gepi-Attee R, Sevilleja JEAD, and Roche JK. PediatricDiarrhea in Southern Ghana: Etiologyand Association with Intestinal Inflammation and Malnutrition. *American Journal of Tropical Medicine and Hygiene*, vol. 83, no. 4, pp. 936-943, 2010.
- 34 N. Orish, V., Ofori-Amoah, J., HefoumeAmegan-Aho, K., Mac-Ankrah, L., Jamfaru, I., Afeke, I., & K. Adzaku, F. (2017). Low Prevalence of Helminth Infections among Primary School Children in the Volta Region of Ghana. *Asian Journal of Medicine and Health*, vol. 5, no. 3, pp. 1–9.
- 35 Xu B, Gordon CA, Hu W, McManus DP, Chen HG, Gray DJ et al. A novel procedure for precise quantification of *Schistosoma japonicum* eggs in bovine feces. *PLoS Negl Trop Dis*, vol. 6, no. 11, pp. 1-8, 2012.
- 36 Brummaier T, Archasuksan L, Watthanakulpanich D, Paris DH, Utzinger J, McGready R, et al. Improved Detection of Intestinal Helminth Infections with a Formalin Ethyl-Acetate-Based Concentration Technique Compared to a Crude Formalin Concentration Technique. *Tropical Medicine and Infectious Disease*, vol. 6, no. 2, pp. 51, 2021.

- 37 Centers for Disease control and Prevention (CDC) 2016. Laboratory Identification of Parasites of Public Health Concern. Stool Specimens - Specimen Processing. Formalin-Ethyl Acetate Sedimentation Concentration.
<https://www.cdc.gov/dpdx/diagnosticprocedures/stool/specimenproc.html#:~:text=Add%2010%20ml%20of%2010,%C3%97%20g%20for%2010%20minutes>. Accessed on January 20, 2024 at 2:40 GMT]
- 38 United Nations Children's Fund UNICEF. *One Is Too Many: Ending child deaths from pneumonia and diarrhoea*. UNICEF, New York, page 19-24, 2016
- 39 Ayeh-Kumi PF, Quarcoo S, Kwakye-Nuako G, Kretchy JP, Osafo-Kantanka A, and Mortu S. Prevalence of Intestinal Parasitic Infections among Food Vendors in Accra, Ghana. *J Trop Med Parasitol*, vol. 32, no. 1, pp. 1-8, 2009.
- 40 United Nations Children's Fund and World Health Organization. *Progress on Sanitation and Drinking Water: 2015 update and MDG assessment*. WHO, Geneva, July 2015. page 5, 2015
- 41 Simon-Oke IA, Ogunleye E. Prevalence of *Entamoeba histolytica* among primary school children in Akure, Ondo State, Nigeria. *Journal of Public Health and Epidemiology*, vol. 7, no. 11, pp. 346-351, 2015.
- 42 Walana P, Tay SCK, Tetteh P, Ziem JB. Prevalence of Intestinal Protozoan Infestation among Primary School Children in Urban and Peri-Urban Communities in Kumasi, Ghana. *Science Journal of Public Health*, vol. 2, no. 2, pp. 52-57, 2014.
- 43 David LH. Amebiasis. Control of Communicable Diseases Manual 19th ed, *American Public Health Association*, Washington, page 11-15, 2008
- 44 Abozakra R, Mokhles M, and Baraka K. Prevalence and Molecular Differentiation of *Entamoeba histolytica*, *Entamoeba dispar*, *Entamoeba moshkovskii*, and *Entamoeba hartmanni* in Egypt. *Acta Parasit.* vol. 65, pp. 929-935, 2020.
- 45 Nath J, Ghosh SK, Singha B, Paul J. Molecular Epidemiology of Amoebiasis: A Cross-Sectional Study among North East Indian Population. *PLoS Negl Trop Dis*, vol. 9, no. 12, pp. 1-19, 2015.

- 46 Duchêne M. *Entamoeba*. In: Walochnik, J., Duchêne, M. (eds) *Molecular Parasitology*. Springer, Vienna, 2016.
- 47 Surtiastuti HJ, Manan S. Intestinal parasites from fingernails of sidewalk food vendors. *Universa Medicina*, vol.30, no. 2, pp. 120-125, 2011.
- 48 Sofiana L, Sumarni S, and Ipa M. Fingernail biting increase the risk of soil transmitted helminth (STH) infection in elementary school children. *Health Science Journal of Indonesia*, vol. 2, no. 2, pp. 81-86, 2011.
- 49 Al-Mekhlafi AM, Abdul-Ghani R, Al-Eryani SM, Saif-Ali R, and Mahdy MA. School-based prevalence of intestinal parasitic infections and associated risk factors in rural communities of Sana'a, Yemen. *Acta tropica*, 163, 135-141, 2016.
- 50 Al-Zabedi EM, Kaid FA, Sady H, Al-Adhroey AH, Amran AA, Al-Maktari MT. Prevalence and Risk Factors of Iron Deficiency Anemia among Children in Yemen. *American Journal of Health Research*, vol. 2, no. 5, pp. 319-326, 2014.
- 51 Calegar DA, Nunes BC, Monteiro KJL, Santos JPD, Toma HK, Gomes TF, et al. Frequency and molecular characterisation of *Entamoeba histolytica*, *Entamoeba dispar*, *Entamoeba moshkovskii*, and *Entamoeba hartmanni* in the context of water scarcity in northeastern Brazil". *Memórias do Instituto Oswaldo Cruz*, vol. 111, pp. 114-119, 2016.
- 52 Codony F, Pérez LM, Agrados B, Agustí G, Fittipaldi M, and Morató J. Amoeba-Related health risk in drinking water systems: could monitoring of amoebae be a complementary approach to current quality control strategies? *Future microbiology*, vol 7, no. 1, pp. 25-31, 2012.
- 53 Atabati H, Kassiri H, Shamloo E, Akbari M, Atamaleki A, Sahlabadi F, et al. The association between the lack of safe drinking water and sanitation facilities with intestinal *Entamoeba* spp infection risk: A systematic review and meta-analysis. *PloS one*, vol. 15. No. 11, 2020.
- 54 Gomes TS, Vaccaro L, Magnet A, Izquierdo F, Ollero D, Martinez-Fernandez C, et al. Presence and interaction of free-living amoebae and amoeba-resisting bacteria in water from drinking water treatment plants. *Science of the Total Environment*, vol.7, 719, 137080.2020
- 55 Shaheen M, Scott C, Ashbolt NJ. Long-term persistence of infectious

Legionella with free-living amoebae in drinking water biofilms. *International journal of hygiene and environmental health*, vol. 222 no. 4, pp. 678-686, 2019.

56 David A, Holcomb JK, Sumner T, Adriano Z, de Bruijn E, Nalá R, et al. (2020). Human fecal contamination of water, soil, and surfaces in households sharing poor-quality sanitation facilities in Maputo, Mozambique. *International Journal of Hygiene and Environmental Health*, vol. 226, pp. 113496, 2020,

57 Cochran WG. *Sampling techniques* (3rd ed.). New York: John Wiley & Sons. 1977.

UNDER PEER REVIEW