

Original Research Article

Comparison of Physico-Chemical and Sensory Attributes in Plant-Based Meat Analogue Patties and Chicken Patties

ABSTRACT

Aim: This study aims to elucidate the differences between PBMA and chicken meat patties by comparing their physico-chemical and sensory attributes.

Study design and methodology: The plant-based meat analogue (PBMA) market is expanding rapidly alongside the burgeoning alternative protein sector. To enhance marketability, the sensory and textural attributes of PBMA must closely mimic those of traditional meat products. Commercial PBMA patties from three brands and chicken patties from two brands available in the Indian market were analysed. The frozen samples were thawed at refrigeration temperature ($4\pm 1^{\circ}\text{C}$) before laboratory analysis of various physico-chemical and sensory properties.

Results: Qualitative analysis indicated that PBMA patties had lower moisture content but higher fat, crude fiber and total ash content compared to chicken patties. Notably, the cholesterol content of PBMA patties was negligible. Linoleic acid was the most abundant fatty acid in PBMA samples, with significant amounts of oleic and palmitic acids also present. The predominant saturated fatty acid (SFA) in all samples was palmitic acid (C16:0). Sensory evaluation revealed that chicken patties scored higher for overall acceptability than PBMA patties.

Conclusion: Overall, the study demonstrates significant differences in proximate composition, texture and sensory qualities between PBMA and traditional meat patties, highlighting the distinct characteristics of PBMA as an alternative protein source.

Keywords: Plant-based meat analogue patties, proximate composition, total phenolic content, cholesterol content, sensory analysis

1. INTRODUCTION

The rising interest in plant-based meat analogues (PBMA) is driven by concerns related to human health, environmental sustainability and animal welfare. PBMAs are recognized as sustainable alternatives that can significantly reduce global greenhouse gas emissions [1,2]. A survey by [3] revealed that approximately 22% of the world's population adheres to a vegetarian diet. To meet the protein needs of vegetarians, protein-rich plant-based ingredients can be incorporated into meat analogues.

Modern PBMAs are specifically formulated to mimic the sensory attributes and macronutrient profiles of traditional meat using plant proteins (e.g., soy, pea, jackfruit, rice, wheat, mushroom), plant-based fats (e.g., canola, coconut, sesame, mustard, soybean, sunflower oil) and other novel ingredients such as soy leghemoglobin, red-colored vegetable extracts and flavouring agents. Mushroom concentrate, for instance, is used as a substitute for monosodium glutamate (MSG) and hydrolyzed vegetable protein for flavour enhancement [4]. Additionally, essential vitamins and minerals typically found in meat, such as iron, zinc, and vitamin B, are being progressively added to PBMAs to enhance their nutritional value [5].

Comparative analyses between PBMAs and conventional meat products are crucial to scientifically substantiate the health benefits and nutritional quality of PBMAs.

While PBMA might be nutritionally inferior to minimally processed whole plant-based foods, their nutritional profile can be enhanced by incorporating various health-promoting components. Moreover, evaluating the extent to which PBMA meet the nutritional and organoleptic properties of traditional meat products is essential to understand and bridge the existing gaps.

A survey by [6] indicated that a significant portion of the population (53%) consumes PBMA at least once a week, with 35% consuming them sporadically and 12% roughly once every 15 days. Collaboration between the food industry and local municipalities can further promote the development of novel and improved PBMA products[7]. Strategies to increase PBMA consumption should consider dietary habits and lifestyles, focusing on improving sensory properties and providing consumer education[8].

This study aims to analyze the differences between plant-based meat analogues and chicken meat patties by comparing their physico-chemical and sensory characteristics. The goal is to determine how well current market-available PBMA meet the nutritional and organoleptic standards of traditional meat products and to identify potential areas for improvement in PBMA products.

2. MATERIALS AND METHODS

2.1 Sample collection

PBMA patties from three commercial brands and chicken patties from two brands were selected. Eight batches of sample from each of the three PBMA and two chicken patty brands were collected. So, for each parameter eight samples were analysed. These frozen products were thawed at refrigeration temperature ($4\pm 1^{\circ}\text{C}$). Subsequently, various physico-chemical, nutritional and sensory qualities of both PBMA and chicken patties were analysed in the laboratory.

2.2 Proximate composition

The moisture content was estimated by hot air oven, protein using automatic digestion and distillation unit, fat was estimated by ether extraction, crude fibre, total dietary fibre and ash content following standard procedure of [9]. The carbohydrate content was obtained by the differential method.

2.3 Fatty acid profile

Fatty acid methyl esters (FAMES) of the samples were prepared following the method described by [10] for the profiling of the fatty acid constituents. The fatty acid composition of extract was determined by injecting 1 μ L of sample into gas chromatograph on a SPTM-2560 capillary column with an internal diameter of 0.25 mm (100 m \times 0.25 \times 0.2 μ m film thickness). The analysis was performed on a Varian 450 gas chromatograph equipped with a flame ionization detector. Nitrogen was used as carrier gas. The injection port temperature was 220°C, and the detector temperature was 220°C. The oven temperature was increased to 175°C for five min and ramped to 220°C at 15°C/min; it was then held at 220°C for 30 min. A software calculated retention times and peak area percentages. Fatty acids were identified by comparing sample retention times with standard retention times (Supelco 37 component FAME mix, Merck). The results of the fatty acid profile were expressed as relative percentage of the peak areas.

2.4 Cholesterol content

The total cholesterol of the sample was determined by a method described by [11]. 100 μ L of lipid extract (prepared from 2g of sample volume made to 5ml with chloroform) was pipetted and 50 μ L of standard cholesterol solution was added separately into test tubes and evaporated to dryness in a water bath. The dried

residue in each tube was dissolved in 2 ml of chloroform to which 1 ml ZnCl₂ reagent and 1 ml acetyl chloride were added. The samples were then heated in a water bath at 50°C for 10 min. For blank, 2 ml of chloroform to which 1 ml ZnCl₂ reagent and 1 ml acetyl chloride were added. The colour complex formed (pink-red colour) was measured by reading the optical density at 528 nm in a spectrophotometer (UV-1700 PharmaSpec, Shimadzu, Japan), and cholesterol content was expressed as mg per 100g of sample.

$$\text{Cholesterol (mg/100g)} = \frac{\text{OD of unknown} \times \text{Conc. of standard} \times 5 \times 100}{\text{OD of standard} \times 0.1 \times 2 \times 1000}$$

2.5 Total phenolic content

The sum of the phenolic compounds of the samples was determined using the method of Folin and Ciocalteu[12] with modifications. Samples of 0.1 g were homogenized in 20 mL of ethanol and water (1:1). The extraction was kept in a shaking water bath (Kemi water bath incubator shaker, India) at 40°C for 10 min and then centrifuged for 10 min at 5000 rpm in a centrifuge (Eppendorf centrifuge 5430 R, Germany). The filtered extract (1 mL) was mixed with 5 ml Folin-Ciocalteu solution (1ml of Folin-Ciocalteu reagent (Loba Chemie Pvt Ltd., India) in 10 ml water and 4 ml sodium carbonate (75 g/L) (Sigma Aldrich Inc., USA) and incubated in darkness for 30 min. The absorbance was measured spectrophotometrically at a wavelength of 765 nm (UV-1700 PharmaSpec, Shimadzu, Japan). The calibration curve was plotted by mixing 1 ml aliquots of 0.1, 0.5, 1.0, 2.5 and 5.0 mg/ml gallic acid solutions with 5.0 ml of Folin-Ciocalteu reagent (diluted tenfold) and 4.0 ml of sodium carbonate solution (75 g/l). The absorbance was measured after 30 min at 765 nm. The sum of phenolic compounds was expressed as mg/gallic acid equivalents (GAE)/g sample.

2.6 Texture profile

A texture analyzer (TiniusOlsen,HIKF,UnitedKingdom) attached to software, texture expert which calculated the springiness, hardness, chewiness and cohesiveness ratio of the samples. Before the test, the frozen extrudates were cut in a cubical shape with dimensions of 24 mm (width) × 24 mm (length) × 14 mm (height) and thawed in an oven for 30 min at 40°C. The sample was placed on the platform and compressed to 80 per cent of its original height using a 75 mm diameter flat bottom probe. The compression was carried out at a crosshead speed of 0.5 mm per second through a two-cycle sequence to mimic chewing process [13].

2.7 Sensory attributes

For the sensory evaluation, an affective test was carried out at the ICAR-National Meat Research Institute. A total of 50 participants were selected randomly among students and staff at the ICAR-National Meat Research Institute. In this case, consumers had to score how much they liked the different samples on a 8-point hedonic score card (1=extremely undesirable; 8 = extremely desirable). Samples were deep fried for 3 min on the day of the sensory analysis to finish the cook. Then, samples were cut into small pieces of 20 g each and served. All samples were coded with random three-digit numbers and water was provided to clean the palate between samples. The attributes evaluated by the consumers were appearance, flavour, juiciness, texture and overall acceptability.

2.8 Statistical analysis

The experiment was replicated four times, each in duplicate. Data obtained for physico-chemical and sensory parameters was compiled and analyzed using SPSS (version 26.0 for windows, SPSS, Chicago, USA). The data was subjected to analysis

of variance (ANOVA) and Duncan's *post hoc* test for for multiple comparisons among different groups.

3. RESULTS AND DISCUSSION

3.1 Proximate composition

The proximate composition of different PBMA and chicken patties is presented in table 1. Significant differences ($P<0.05$) were observed between PBMA and chicken patties in terms of moisture, protein, fat, crude fiber, total dietary fiber, carbohydrate and total ash content. These variations could be attributed to the heterogeneous nature of the different proteins and ingredients used during formulation.

The moisture content of PBMA and chicken patties differed significantly ($P<0.05$). The results were in accordance with [14] who emphasized that meat analogues displayed moisture content between 46.98 and 53.71 per cent. The moisture content of final product depends on the hydration capacity of basic ingredients, temperature and different cooking conditions.

The protein content varied significantly ($P<0.05$) between PBMA and chicken patties. Among the samples, PBMA patty PP-3 had the highest protein content ($P<0.05$). The protein content in PBMA can be tailored to meet the nutritional needs of specific consumer groups, such as the elderly, pregnant women or children. There was a significant difference ($P<0.05$) in fat content between chicken and PBMA patties, with chicken patties having lower fat content. The addition of fat in PBMA recipes enhances juiciness, tenderness and flavour release [15]. Furthermore, fats help retain volatile flavour compounds, improving the sensory profile of PBMA [16].

Table 1. Values of nutritional parameters analyzed for different PBMA and chicken patties (mean±standard error)

Parameters	CP-1	CP-2	PP-1	PP-2	PP-3
------------	------	------	------	------	------

Moisture (%)	58.39±0.163 ^a	56.46±0.079 ^b	51.46±0.058 ^c	51.17±0.176 ^c	56.77±0.071 ^b
Protein (%)	12.29±0.046 ^b	9.47±0.026 ^d	9.65±0.027 ^c	7.42±0.051 ^e	17.34±0.044 ^a
Fat (%)	10.70±0.087 ^b	10.42±0.058 ^c	9.70±0.026 ^d	12.55±0.066 ^a	8.37±0.034 ^e
Carbohydrate (%)	16.44±0.11 ^c	22.14±0.09 ^b	25.63±0.07 ^a	25.57±0.08 ^a	13.41±0.05 ^d
Crude fibre(%)	0.59±0.02 ^d	0.51±0.02 ^e	1.17±0.02 ^a	0.98±0.02 ^b	0.86±0.02 ^c
Total dietary fibre(%)	1.15±0.01 ^e	1.17±0.01 ^d	1.97±0.01 ^c	2.03±0.01 ^a	1.99±0.01 ^b
Ash (%)	1.60±0.017 ^c	0.99±0.020 ^d	2.38±0.040 ^b	2.32±0.014 ^b	3.25±0.021 ^a
Cholesterol content (mg/100g)	46.21±0.207 ^b	50.64±0.228 ^a	0.00±0.00 ^c	0.00±0.00 ^c	0.00±0.00 ^c

Values reported are mean values and standard errors (n=8). Same superscripts in a row does not differ significantly ($p>0.05$). CP-1: chicken patties-1; CP-2: chicken patties-2; PP-1: PBMA patties -1; PP-2: PBMA patties -2; PP-3: PBMA patties -3

The carbohydrate content of PBMA and chicken patties differed significantly ($P<0.05$). PBMA patty PP-3 had the lowest carbohydrate content among all samples ($P<0.05$). Crude fiber content varied significantly ($P<0.05$) between PBMA and chicken patties, with PBMA patties having higher crude fiber content. Similarly, total dietary fiber (TDF) content was significantly higher ($P<0.05$) in PBMA patties compared to chicken patties. Increased dietary fiber is beneficial for health as it reduces total cholesterol, particularly LDL cholesterol and helps limit glucose absorption [17].

The ash content, representing total mineral content, showed significant differences ($P<0.05$) between PBMA and chicken patties. PBMA patties had higher ash content compared to chicken patties. [18], reported significant difference in the proximate composition among different animal-based and plant based products.

3.2 Cholesterol content

The cholesterol content varied significantly between chicken patties CP-1 and CP-2 ($P<0.05$), with CP-2 having higher cholesterol content. In contrast, PBMA patties had

no detectable cholesterol content. This finding is consistent with [19], who reported zero cholesterol in market-available PBMA products such as the Beyond Burger and Impossible Burger. [20] found that the cholesterol content in chicken nuggets ranged from 125.50 to 32.58 mg per 100 g.

3.3 Fatty acid profile

The fatty acid profile of food products is crucial for understanding their nutritional implications, particularly their effects on cardiovascular health[21]. Substituting polyunsaturated fatty acids (PUFA) for saturated fatty acids (SFA) has been linked to reduced cardiovascular risk, emphasizing the importance of a higher PUFA/SFA ratio in dietary recommendations[22]. Table 2 presents the mean fatty acid compositions of PBMA and chicken patties, highlighting significant differences among samples. In all groups, palmitic acid (C16:0) was the predominant SFA. Notably, PP-3 exhibited the lowest SFA content, whereas CP-2 had the highest. Oleic acid (C18:1n9c) was significantly higher in CP-2, while linoleic acid (C18:2n6c) was notably higher in PP-3. PBMA samples predominantly featured linoleic acid, alongside substantial amounts of oleic and palmitic acids, consistent with common dietary sources of MUFAs and SFAs in human diets.

Oleic acid, a prevalent monounsaturated fatty acid (MUFA) in human diets, has been associated with favorable effects on lipid profiles and metabolic health, potentially mitigating the adverse effects of palmitic acid (C16:0) on insulin sensitivity and mitochondrial function[23].

Table 2. Fatty acid profiles of different PBMA and chicken patties (mean±standard error)

Fatty acids	CP-1	CP-2	PP-1	PP-2	PP-3
C8:0	0.01±0.01 ^d	0.02±0.01 ^d	1.32±0.14 ^a	0.89±0.06 ^b	0.28±0.04 ^c
C10:0	ND	0.01±0.01 ^d	1.14±0.07 ^a	0.89±0.04 ^b	0.24±0.02 ^c

C12:0	0.03±0.01 ^c	0.14±0.03 ^c	10.89±0.28 ^a	9.14±0.17 ^b	0.16±0.02 ^c
C14:0	0.43±0.01 ^d	0.72±0.03 ^c	4.38±0.16 ^a	3.97±0.05 ^b	0.16±0.01 ^e
C16:0	16.22±0.30 ^b	29.84±0.21 ^a	13.12±0.20 ^c	14.22±0.11 ^c	8.25±0.72 ^d
C17:0	2.48±0.04 ^b	2.90±0.15 ^a	0.10±0.02 ^c	0.10±0.03 ^c	0.04±0.01 ^c
C18:0	4.25±0.06 ^{cd}	4.71±0.11 ^{bc}	5.14±0.45 ^{ab}	5.73±0.17 ^a	3.66±0.29 ^d
C18:1n9c	34.13±0.17 ^b	41.38±0.36 ^a	26.01±0.33 ^d	30.06±0.30 ^c	32.41±1.13 ^b
C18:2n6c	38.53±0.27 ^b	17.55±0.40 ^d	35.66±0.55 ^{bc}	32.03±0.32 ^c	51.90±3.17 ^a
C18:3n3	0.93±0.03 ^a	0.20±0.11 ^c	0.48±0.05 ^b	0.52±0.03 ^b	0.51±0.01 ^b
C21:0	ND	0.43±0.13 ^a	0.51±0.16 ^a	0.10±0.09 ^b	ND
C22:0	0.23±0.02 ^c	0.54±0.01 ^b	0.22±0.02 ^c	0.28±0.01 ^c	0.60±0.04 ^a
SFA	23.64±0.35 ^c	38.82±0.12 ^a	36.81±0.49 ^b	35.31±0.21 ^b	13.39±1.13 ^d
MUFA	34.13±0.17 ^b	41.37±0.36 ^a	26.01±0.33 ^d	30.06±0.30 ^c	32.41±1.30 ^b
PUFA	39.46±0.29 ^b	17.75±0.36 ^d	36.14±0.49 ^{bc}	32.56±0.30 ^c	52.40±3.18 ^a
MUFA/SFA	1.44±0.01 ^b	1.07±0.01 ^c	0.71±0.02 ^e	0.85±0.01 ^d	2.46±0.10 ^a
PUFA/SFA	1.67±0.04 ^b	0.46±0.01 ^c	0.98±0.02 ^c	0.92±0.01 ^c	4.11±0.48 ^a
UFA	73.60±0.22 ^b	59.13±0.16 ^d	62.15±0.60 ^c	62.61±0.20 ^c	84.82±1.89 ^a
UFA/SFA	3.12±0.05 ^b	1.52±0.01 ^c	1.69±0.04 ^c	1.77±0.01 ^c	6.57±0.57 ^a

Values reported are mean values and standard errors (n=8). Same superscripts in a row does not differ significantly ($p>0.05$).

MUFA: monounsaturated acids, SFA: saturated fatty acids, UFA = sum of PUFA and MUFA, ND: not detected. CP-1: chicken patties-1; CP-2: chicken patties-2; PP-1: PBMA patties -1; PP-2: PBMA patties -2; PP-3: PBMA patties -3

The presence of specific medium-chain fatty acids (C8:0, C10:0, C12:0, C14:0) in PP-1 and PP-2 suggests the incorporation of coconut oil in these formulations. Conversely, higher levels of palmitic acid (C16:0) and lower levels of linoleic acid (C18:2n6c) in CP-2 likely indicate the use of palm oil, either added during processing or as a frying medium. PP-3, characterised by lower palmitic acid (C16:0) and higher linoleic acid (C18:2n6c), likely reflects the use of soybean oil in its preparation. These findings underscore the variability in fatty acid profiles among different product formulations, influencing their nutritional quality and potential health implications.

3.4 Total phenolic content

The total phenolic content (TPC) of various PBMA and chicken patties was assessed, and the results are summarized in Table 3. Significant differences ($P<0.05$) were observed among all groups, likely attributable to the distinct ingredients used in their formulations. A diet rich in plant polyphenols has been

documented to enhance health and reduce the risk of cardiovascular diseases [21]. Polyphenolic antioxidants mitigate the detrimental effects of reactive oxygen species, which is beneficial for overall health and may help prevent dementia and memory loss [24].

Given the strong antioxidant properties and potential health benefits of phenolic compounds, there is an increasing emphasis on incorporating these compounds into food products to promote health and combat age-related diseases.

As shown in Table 3, the chicken patties exhibited lower total phenolic content compared to the PBMA samples. Specifically, CP-1 and CP-2 had TPC values of 0.934 ± 0.033 mg GAE/g and 0.781 ± 0.014 mg GAE/g, respectively. The PBMA samples, PP-1, PP-2, and PP-3 demonstrated higher TPC values of 1.567 ± 0.020 mg GAE/g, 2.253 ± 0.013 mg GAE/g, and 2.873 ± 0.019 mg GAE/g, respectively. These results highlight a notable gap in phenolic content between meat-based and plant-based patties.

Table 3. Total phenolic content of different PBMA and chicken patties

Parameters	CP-1	CP-2	PP-1	PP-2	PP-3
Total phenolic content (mg GAE /g)	0.934 ± 0.033 _d	0.781 ± 0.014 _e	1.567 ± 0.020 _c	2.253 ± 0.013 _b	2.873 ± 0.019 _{g^a}

(mean±standard error)

Values reported are mean values and standard errors (n=8). Same superscripts in a row does not differ significantly ($p > 0.05$). CP-1: chicken patties-1; CP-2: chicken patties-2; PP-1: PBMA patties -1; PP-2: PBMA patties -2; PP-3: PBMA patties -3

3.6 Texture profile analysis

Texture is a fundamental quality attribute of food products, influencing consumer acceptance and satisfaction. The instrumental texture characteristics between

chicken patties and PBMA patties differed significantly ($P < 0.05$), as displayed in Table 4.

The hardness values of the PBMA patties were significantly higher ($P < 0.05$) than those of the chicken patties. The increased hardness in PBMA patties can be attributed to better cross-linking of the protein, thus resulting in a firmer structure[25]. This trend is consistent with previous findings where meat analogs exhibited distinct textural properties compared to their meat counterparts[26].

Chewiness, which indicates the energy required to chew the food, also varied significantly ($P < 0.05$) among the samples. The differences in chewiness could be influenced by water content and retention during the extrusion process, with higher water retention typically leading to decreased mechanical properties[27, 28].

In terms of cohesiveness, CP-1 and CP-2 had similar cohesiveness ratios. Among the PBMA patties, PP-1 had a significantly lower ($P < 0.05$) cohesiveness ratio, while PP-2 and PP-3 exhibited higher ratios. The higher cohesiveness in certain PBMA patties may be due to the addition of binding agents designed to replicate the meat-like structure.

Springiness, which measures the extent to which a sample returns to its original shape after compression, was consistent among the samples. The springiness values of the chicken patties were consistent with previous studies[29], while the slightly higher values ($P < 0.05$) in PBMA patties may be due to the incorporation of certain binding agents. These findings highlight the significant textural differences between chicken and PBMA patties. Understanding these differences is crucial for food manufacturers aiming to create meat analogs that closely mimic the texture of traditional meat products.

Table 4. Texture profile analysis of different PBMA and chicken patties (mean±standard error)

Parameters	CP-1	CP-2	PP-1	PP-2	PP-3
Hardness (N)	7.52±0.062 ^e	9.30±0.063 ^d	11.80±0.056 ^a	10.28±0.038 ^b	9.47±0.402 ^c
Cohesiveness ratio	0.97±0.008 ^b	0.97±0.004 ^b	0.77±0.004 ^c	1.18±0.108 ^a	1.06±0.006 ^{ab}
Chewiness (N cm)	6.60±0.093 ^d	8.27±0.026 ^a	7.43±0.028 ^c	6.63±0.034 ^d	7.63±0.069 ^b
Springiness (cm)	0.97±0.004 ^c	0.97±0.006 ^c	1.04±0.003 ^a	1.02±0.004 ^b	1.04±0.002 ^a

Values reported are mean values and standard errors (n=8). Same superscripts in a row does not differ significantly ($p>0.05$). CP-1: chicken patties-1; CP-2: chicken patties-2; PP-1: PBMA patties -1; PP-2: PBMA patties -2; PP-3: PBMA patties -3

3.7 Sensory analysis

Sensory attributes play a crucial role in determining consumer acceptance of food products. Table 5 presents the sensory evaluation scores of chicken patties (CP) and PBMA patties, assessed based on appearance, flavour, juiciness, texture, and overall acceptability.

The appearance values varied significantly ($P<0.05$) between chicken and PBMA patties, as shown in Table 5. CP-1 had the highest appearance score, while PP-3 had the lowest. The appearance scores of CP-2, PP-1, and PP-2 did not differ significantly ($P>0.05$), likely due to similar enrobing techniques used in both chicken and PBMA patties. The appearance of PBMA patties was designed to mimic traditional meat products, with ingredients such as beetroot juice used to replicate the reddish color of meat patties [30, 31].

Flavour scores showed significant variation ($P<0.05$) between chicken and PBMA patties. CP-1 had the highest flavour score, indicating a preference for the flavour of chicken patties. PBMA patties, particularly PP-3, received lower flavour scores. The

slight bitterness and less appealing flavour of PBMA patties highlight the challenge in replicating the taste of chicken patties.

Juiciness scores also differed significantly ($P < 0.05$) between the samples. CP-1 and CP-2 had similar juiciness scores, while PBMA patties PP-2 and PP-3 did not differ significantly ($P > 0.05$) in their juiciness scores. The juiciness of PBMA products may be influenced by the moisture content of the initial formulation and the cooking method employed.

Texture or tenderness scores varied significantly ($P < 0.05$) between the samples. CP-2 and PP-2 had the highest texture scores, indicating a preference for their tenderness. In contrast, PP-3 had the lowest texture score. The texture of PBMA patties is affected by the use of binding agents and the specific mechanical energy applied during processing.

Overall acceptability scores showed significant variation ($P < 0.05$) between chicken and PBMA patties. CP-1 had the highest overall acceptability score, followed by CP-2. Among the PBMA patties, PP-1 and PP-2 had lower acceptability scores, while PP-3 had the lowest score.

These results highlight the necessity of optimising sensory attributes-particularly appearance, flavour and texture, in the development of PBMA products to enhance consumer acceptance. Although PBMA patties demonstrate potential, there remains a notable gap in matching the sensory qualities of traditional chicken patties.




Fig.1. Radar plot for sensory scores of different PBMA and chicken patties

Table 5. Sensory score values of different PBMA and chicken patties (mean±standard error)

Parameters	CP-1	CP-2	PP-1	PP-2	PP-3
Appearance and colour	7.19±0.091 ^a	7.00±0.00 ^b	7.00±0.00 ^b	7.00±0.00 ^b	6.13±0.081 ^c
Flavour	7.00±0.00 ^a	6.38±0.081 ^b	6.13±0.081 ^c	6.00±0.00 ^c	5.00±0.00 ^d
Juiciness	6.88±0.081 ^a	7.00±0.00 ^a	6.31±0.091 ^b	6.00±0.00 ^c	6.00±0.00 ^c
Texture/tenderness	6.63±0.081 ^b	7.00±0.00 ^a	6.00±0.00 ^c	7.00±0.00 ^a	5.00±0.00 ^d
Overall acceptability	7.25±0.094 ^a	7.00±0.00 ^b	6.00±0.00 ^c	6.13±0.081 ^c	5.00±0.00 ^d

Based on eight point hedonic scale (1=extremely undesirable; 8 = extremely desirable). Values reported are mean values and standard errors (n=8). Same superscripts in a row does not differ significantly ($p>0.05$). CP-1: chicken patties-1; CP-2: chicken patties-2; PP-1: PBMA patties -1; PP-2: PBMA patties -2; PP-3: PBMA patties -3

4. CONCLUSION

This study highlights the significant differences between plant-based meat analog (PBMA) patties and traditional chicken patties in terms of proximate composition, texture, and sensory qualities. The proximate composition analysis revealed notable variations in moisture, protein, fat, crude fiber, total dietary fiber, carbohydrate and total ash content between PBMA and chicken patties. These differences underline

the current gaps in nutritional and sensory attributes that need to be addressed to enhance the acceptance of PBMA products.

The texture and sensory evaluations showed that PBMA patties, while promising, still fall short in replicating the sensory qualities of chicken patties, particularly in appearance and flavour. This gap can be bridged by exploring more suitable colouring and flavouring ingredients that can better mimic the sensory profiles of meat. Additionally, understanding the structure formation mechanism during extrusion and shear processes is crucial. Advances in these areas will enable the development of PBMA products with improved textural and sensory characteristics.

To produce meat substitutes with superior resource efficiency and the desired nutritional and sensory qualities, there is a need for continued development of analytical techniques and structural procedures. Enhancing the extraction and detailed characterization of unique protein fractions will provide better insights into the functionality of raw materials, leading to improved PBMA products. Physicochemical, thermal, chemical or enzymatic treatments of legume and oilseed meals, concentrates or isolates can yield products with qualities suitable for food applications. Further scientific research is essential to substantiate the nutritional quality and health benefits of PBMA compared to conventional meat products.

REFERENCES

1. Springmann M, Clark M, Mason-D'Croz D, Wiebe K, Bodirsky BL and Lassaletta L. Options for keeping the food system within environmental limits. *Nature*. 2018;562: 519–525.

2. Willett W, Rockstrom J, Loken B, Springmann M, Lang T, Vermeulen S. Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. *Lancet*. 2019;393: 447–492.
3. Leahy E, Lyons S and Tol RS. An estimate of the number of vegetarians in the world (No. 340). 2010; ESRI working paper.
4. Bakhsh A, Lee SJ, Lee EY, Hwang YH and Joo ST. Traditional plant-based meat alternatives, current, and future perspective: a review. *Journal of Agriculture and Life Sciences*. 2021;55(1): 1-10.
5. Curtain F and Grafenauer S. Plant-based meat substitutes in the flexitarian age: An audit of products on supermarket shelves. *Nutrients Journal*. 2019;11(11): 2603.
6. Rizzo G, Testa R, Dudinskaya EC, Mandolesi S, Solfanelli F, Zanolli R, Schifani G and Migliore G. Understanding the consumption of plant-based meat alternatives and the role of health-related aspects. A study of the Italian market. *International Journal of Gastronomy and Food Science*. 2023;32, p.100690.
7. Westling M, Wennstrom S and Ostrom A. Public meals as a platform for culinary action? Tweens' and teens' acceptance of a new plant-based food. *International Journal of Gastronomy and Food Science*. 2022;27: p.100485.
8. Perez-Cueto FJ, Rini L, Faber I, Rasmussen MA, Bechtold KB, Schouteten JJ and De Steur H. How barriers towards plant-based food consumption differ according to dietary lifestyle: Findings from a consumer survey in 10 EU countries. *International Journal of Gastronomy and Food Science*. 2022;29, p.100587.

9. AOAC International. Official Methods of Analysis. 2016; Twentieth ed. Rockville, Maryland, USA.
10. Wang, JUN Wu W, Wang X, Wang MIN and Wu F. An effective GC method for the determination of the fatty acid composition in silkworm pupae oil using a two-step methylation process. Journal of Serbian Chemical Society. 2015;80(1): 9-20.
11. Hanel HK and Dam H. Determination of small amounts of total cholesterol by Tschugaeff reaction with a note on determination of lanosterol. Acta Chemica Scandinavica. 1955;9: 677.
12. Szpicer A, Onopiuk A, Barczak M and Kurek M. The optimization of a gluten-free and soy-free plant-based meat analogue recipe enriched with anthocyanins microcapsules. LWT-Food Science Technology. 2022;168, p.113849.
13. Tuccillo F, Kantanen K, Wang Y, Diaz JMR, Pulkkinen M, Edelmann M, Knaapila A, Jouppila K, Piironen V, Lampi AM and Sandell M. The flavor of faba bean ingredients and extrudates: Chemical and sensory properties. Food Research International. 2022;162, p.112036.
14. Chiang JH, Tay W, Ong DSM, Liebl D, Ng CP and Henry CJ. Physicochemical, textural, and structural characteristics of wheat gluten-soy protein composited meat analogues prepared with the mechanical elongation method. Food Structure. 2021;28. p.100183.
15. Egbert Rand Borders C. Achieving success with meat analogs. Food Technology Journal. 2006;60:28-34.

16. Kyriakopoulou K, Dekkers B and van der Goot AJ. Plant-based meat analogues. In: Sustainable meat production and processing (pp. 103-126). 2019; Academic Press.
17. Dhingra D, Michael M, Rajput H and Patil RT. Dietary fibre in foods: a review. *Journal of Food Science Technology*. 2012;49 (3): 255–266.
18. Katidi A, Xypolitaki K, Vlassopoulos A and Kapsokefalou M. Nutritional Quality of Plant-Based Meat and Dairy Imitation Products and Comparison with Animal-Based Counterparts. *Nutrients*. 2023;15(2), p.401.
19. Bohrer B M. An investigation of the formulation and nutritional composition of modern meat analogue products. *Food Science and Human Wellness*. 2019;8(4): 320-329.
20. El-Anany A M, Ali R F and Elanany A M. Nutritional and quality characteristics of chicken nuggets incorporated with different levels of frozen white cauliflower. *Italian Journal of Food Science*. 2020;32(1).
21. Martinez E and Ramos-Escudero F. Valorization of flours from cocoa, sinami and sachainchi by-products for the reformulation of Peruvian traditional flatbread (*'Pan Chapla'*). *International Journal of Gastronomy and Food Science*. 2024;p.100930.
22. Ozdemir VF, Kocyigit R, Yanar M, Aydin R, Diler A, Palangi V and Lackner M. An investigation of slaughter weight and muscle type effects on carcass fatty acid profiles and meat textural characteristics of young Holstein Friesian bulls. *Heliyon*. 2024;10(6).
23. Shramko VS, Polonskaya YV, Kashtanova EV, Stakhneva EM and Ragino YI. The short overview on the relevance of fatty acids for human cardiovascular disorders. *Biomolecules*. 2020;10(8), p.1127.

24. Huang D. Dietary antioxidants and health promotion. *Antioxidants*. 2018;7 (1): 9.
25. Zhang J, Liu L, Zhu S and Wang Q. Texturisation behaviour of peanut–soy bean/wheat protein mixtures during high moisture extrusion cooking. *International Journal of Food Science and Technology*. 2018;53(11): 2535-2541.
26. Sun YT and Ruiz-Carrascal J. Home made vegan nuggets with texturized soy protein and tempeh as compared to chicken-based ones: texture, consumer perception and environmental impact. *International Journal of Gastronomy and Food Science*. 2023;33, p.100748.
27. Lin S, Huff H E and Hsieh F. Texture and chemical characteristics of soy protein meat analog extruded at high moisture. *Journal of Food Science*. 2000;65(2): 264–269.
28. Palanisamy M, Franke K, Berger RG, Heinz V and Topfl S. High moisture extrusion of lupin protein: Influence of extrusion parameters on extruder responses and product properties. *Journal of the Science Food Agriculture*. 2019;99(5): 2175-2185.
29. Wan Rosli WI, Solihah MA, Aishah M, Nik Fakurudin NA and Mohsin SSJ. Colour, textural properties, cooking characteristics and fibre content of chicken patty added with oyster mushroom (*Pleurotus sajor-caju*). *International Food Research Journal*. 2011;18(2).
30. Kyriakopoulou K, Keppler JK and van der Goot AJ. Functionality of ingredients and additives in plant-based meat analogues. *Foods*. 2021;10(3), p.600.

31. Botella-Martinez C, Viuda-Martos M, Fernandez-Lopez JA, Perez-Alvarez JA and Fernandez-Lopez J. Development of plant-based burgers using gelled emulsions as fat source and beetroot juice as colorant: Effects on chemical, physicochemical, appearance and sensory characteristics. *LWT-Food Science and Technology*. 2022; 172, p.114193.

UNDER PEER REVIEW