

Integrated effect of *Pseudomonas* sp. and bioresources against *Alternaria* leaf spot disease of brinjal (*Solanum melongena* L.)

ABSTRACT

This study was conducted during the *Kharif* season 2022-2023 at the Central Research farm, Department of Plant Pathology, SHUATS, Allahabad, U.P to conclude the Integrated effect of *Pseudomonas* sp. and bioresources against *Alternaria* leaf spot disease of brinjal (*Solanum melongena* L.). Seedling treatment with bioagent viz., *Pseudomonas* sp. at 0.25% and bioresources applied as a foliar spray viz., Neem extract + Clove extract(5 + 5%), Neem oil + Clove oil(5+5%), Neem extract (10%), Clove extract(10%), Neem oil(0.2%), Clove oil(0.2%) and Bavistin(0.1%, treated check) for their effectiveness to manage *Alternaria* leaf spot of brinjal caused by *Alternaria alternata*. All the treatments were significantly reduced the severity of the disease and increased growth parameters. Among all the treatments *Pseudomonas* sp. + Neem extract + Clove extract(32.37%) was most effective followed by *Pseudomonas* sp. + Neem oil + Clove oil (34.39%) were significantly superior over other treatments in reducing the leaf spot infection of brinjal. The highest cost benefit ratio was obtained in the treatment *Pseudomonas* sp. + Neem extract + Clove extract(1:6.72). While other treatments also showed significantly effective/efficacy to check the disease intensity (PDI) and yield over control in the field condition.

Key words: Brinjal, *Pseudomonas* sp, Seedling treatment, Bioresources, Foliar spray, *Alternaria alternata*.

Comment [MM1]: Delete sp and insert 'sp'

Comment [MM2]: Insert hyphen (-) between bio and resources

Comment [MM3]: Separate neem and oil

Comment [MM4]: Delete sp and insert 'sp'

Comment [MM5]: Delete sp and insert 'sp'

Comment [MM6]: Delete over and insert to

Comment [MM7]: Delete sp and insert 'sp'

Comment [MM8]: Delete ly and use significant

Comment [MM9]: Delete effective and use efficacy

INTRODUCTION

Brinjal or Eggplant (*Solanum melongena* L.) is an important vegetable crop grown in varied parts of the world and is the second major vegetable crop next to potato in India. It is considered as the “King of Vegetables”. It is a perennial, ~~herbaceous, erect or semi-spreading plant~~ ~~plant which is herbaceous, erect or semi-spreading, and that~~ is usually grown as a seasonal crop in almost all the seasons (Shrivastava and Butani, 1998).

Comment [MM10]: Use the re-cast version of the sentence

Brinjal fruits are an excellent source of starch, proteins, minerals, vitamins, dietary fibers and low-fat content. It is one of the richest sources of antioxidants mainly ascorbic acid which has been reported to successfully suppress the development and growth of tumors, inhibit inflammation and cardiovascular diseases. The higher ascorbic acid content of the fruits not only helps in better retention of colour and ~~flavour but also affects fruit ripening and stress resistance~~ ~~flavour~~ (Kumar and Arumugam, 2013).

Comment [MM11]: Delete the cancelled flavour so that the sentence be complete.

Total production of Brinjal is about 32 million tonnes in the world where ~~in~~ India is world's second largest producer after China (Choudhary and Gaur, 2009). In India total brinjal cultivation area (730.4 ha), production (12800.8MT) and productivity (17.5 t/ha). Uttar Pradesh covered the cultivated area of 8.01 thousand hectares with an annual production of 275.40 thousand MT and productivity of 34.40MT per hectare and in Andhra Pradesh total cultivated area is 14.60 ha) and production (380.03 MT) (Horticulture Statistics Division, 2018).

Comment [MM12]: Delete in

Brinjal is also a popular vegetable in China, Japan, Egypt, Italy, USA, Syria, Philippines, Thailand, Indonesia, France and Turkey (Beura et al., 2008). The major brinjal producing states in India are Andhra Pradesh, Maharashtra, Karnataka, Orissa, Madhya Pradesh and West Bengal.

The disease first makes its appearance in young seedlings. It attacks leaves and then spreads to fruits which subsequently rot and become unfit for consumption (Bochalya et al., 2012)^[7]. Genus *Alternaria* belongs to deuteromycetes having different species, which are destructive plant pathogen to the families such as Solanaceae, cucurbitaceae, brassicaceae. Species of the *Alternaria* genus are cosmopolitan, surviving both as saprophytes as well as weak parasites. In several cases, small dark-coloured spots are ~~also~~ formed on tender twigs. *Alternaria* ~~sp.~~ is one of the most important and frequently occurring disease of the crop nation and worldwide. *Alternaria* leaf spot was a minor disease, but due to climatic changes, it emerged moderately to severe form ~~now~~ ~~days~~. It was reported up to 25% yield losses due to leaf spot of brinjal (Balai and Ahir, 2013).

Comment [MM13]: Separate dark and coloured with hyphen (-)

Comment [MM14]: Delete also

Comment [MM15]: Delete sp and insert 'sp'

Comment [MM16]: Add ly to frequent

Comment [MM17]: Delete now a days and insert nowadays

Pseudomonas fluorescens @2.5% was found the most effective treatment which gave recorded minimum disease intensity (%) and yield (q/ha), as compared to other treatments (Kakraliya et al.,

2017). It has been previously reported that the active ingredients of neem constitute mostly of terpenoids, eg, Nimbin, Nimbicidine, Azadirachtin etc (Brahmachari, 2004). It has been concluded that *Azadirachtaindica*, a common medicinal plant could be exploited as the source of a potent biocide that have immense fungi toxic effect to fungal pathogen. The anti-fungal action of garlic clove is due to the compound allicin. It has strong anti-microbial and anti-fungal activities. Thus, inhibition of fungi observed in this study may be related to allicin or ajoene which crubs the performance of some enzymes that are important to fungi (Kutawaet al., 2018).

Comment [MM18]: Insert hyphen (-) between anti and fungal

Comment [MM19]: Insert hyphen (-) between anti- microbial

Comment [MM20]: Insert hyphen (-) between anti and fungal

Vermi-compost consistently promote biological activity which can cause plant to germinate, to produce flower and yield better than in commercial containing media, independent of nutrient availability (Aranconet al., 2004).

Comment [MM21]: Use vemi-compost and not vermicompost

Bio-agent and bio-resources belonging to various groups recommended for the management of Alternaria leaf spot of brinjal. Generally, farmers are using only the chemicals for managing the disease, but it has negative impact on the environment and causes resistance in the pathogen.

Comment [MM22]: Insert hyphen between Bio and agent.

Comment [MM23]: Insert hyphen between bio and resources

Comment [MM24]: Add causes between and resistance

Comment [MM25]: Change resistant to resistance

Materials and Methods

The present investigation on “Integrated effect of *Pseudomonas* sp. and bioresources against Alternaria leaf spot of brinjal (*Solanum melongena* L.)” an experiment was carried out at Central Research Field of Sam Higginbottom University of Agricultural, Technology and Sciences, Prayagraj during kharif season 2022-2023. Field experiment was laid out in Randomised block design with three replications. The materials and methods adopted during the studies are described here in this chapter.

Comment [MM26]: Delete an experiment

Identification

The species identification was confirmed by National Fungal Culture Collection of India (NFFCI), Pune. And identified as *Alternaria alternata* (Fr.) Keissler.

The periphery of the colony was olive-green, with a black centre and dull white spots. The growth of the fungus was smooth. The conidiophores were short, arising singly. In shape, the conidia were obclavate but few were oval or pyriform with a rather short, broadly rounded base with 3-8 transverse septa and several longitudinal septa with variable size and shape.



Plate 1. Microscopic view of

Alternaria alternata under 40X

Application of treatments:

Seedlings of brinjal are treated with *Pseudomonas* sp. in 1 lit of water for 30mins @ 2.5% Application of vermicompost before flowering @ 5 ton/ha and with some of neem and clove products namely, neem oil @ 0.2%, neem extract 10%, clove oil @ 0.2%, clove extract @ 10%, neem oil + clove oil @ 0.1%, neem extract + clove extract @ 5% + 5% and Bavistin @ 0.1% were sprayed as foliar spray. As the water and oil do not mix with each other, an emulsifier was used to mix them. An emulsifier creates even consistency between oils and water. Potassium silicate is a good oil emulsifier and is approved for organic use as a foliar spray. Potassium silicate @ 0.2gm was used for mixing oil and water. The subsequent spray was given at 15 days intervals.

Comment [MM27]: Delete sp and insert "sp"

Comment [MM28]: Separate for and organic

Observations:

Comment [MM29]: It is observations (check spelling)

Observations were recorded on plant height (cm), number of branches at 30, 60 and 90 DAT and percentage of disease intensity recorded at 60, 75, 90 DAT.

Disease severity was measured by using 0-5 scale with modification described by **Jaimanet al., (2013)**.

Sum of all disease ratings

$$\text{Disease intensity (\%)} = \frac{\text{Total number of rating} \times \text{Maximum disease grade}}{\text{Total number of rating} \times \text{Maximum disease grade}} \times 100$$

List 1. Disease rating scale (Wheeler, 1969)

Disease rating	Percent leaf area covered
0	Free from infection
1	One or two necrotic spots on few lower leaves of plant, covering nearly 1-10% surface area of plant.
2	A few isolated spots on leaves covering 11-25% surface area of plant.
3	Many spots coalesced on the leaves covering 26-50% of the surface area of the plant.
4	Concentric rings on the stem petiole, fruit covering 51-75% leaf area of plant.
5	Whole plant blighted leaf and fruits starting to fall covering more than 75% leaf area of the plant.

Comment [MM30]: Use covering not coving

Benefit cost ratio:

Gross returns will be calculated by multiplying total yield with the market price of the produce.

Cost of cultivation and cost of treatment imposition will be deducted from the gross returns, to

find out net returns and cost benefit ratio by the following formula below:

$$\frac{\text{Net returns}}{\text{Total Cost of Cultivation}}$$

$$B : C = \frac{\text{Net returns}}{\text{Total Cost of Cultivation}}$$

Total Cost of Cultivation

Where, B: C = Benefit Cost Ratio

Comment [MM31]: Add the and below to the sentence

Formatted: Font color: Text 1

Formatted: Justified, Indent: Left: 0", Right: 0.14", Space After: 3.05 pt, Line spacing: Multiple 1.46 li

RESULT AND DISCUSSION

Table 1. Effect of selected treatments on *Alternariaalternata* disease intensity (%) of brinjal at 60, 75, 90 DAT.

	Treatments	Disease intensity (%) Mean of three replications		
		60DAT	75 DAT	90 DAT
T0	Control	18.67	31.57	41.43
T1	<i>Pseudomonas</i> sp. + Neem oil @0.2%	15.20	27.38	37.56
T2	<i>Pseudomonas</i> sp. + Neem extract @10%	14.10	25.42	36.24
T3	<i>Pseudomonas</i> sp. + Clove oil @0.2%	17.45	28.44	39.42
T4	<i>Pseudomonas</i> sp. + Clove extract @10%	14.96	26.20	36.61
T5	<i>Pseudomonas</i> sp. + Neem oil @0.1%+ clove oil @0.1%	13.45	24.48	34.39
T6	Bavistin (treated check)	11.08	22.34	32.66
T7	<i>Pseudomonas</i> sp. + Neem extract @5%+ clove extract @5%	12.88	22.53	32.37
CD(5%)		0.77	1.32	1.05
SE d±		0.37	0.65	0.35
CV		3.46	3.64	1.98

Comment [MM32]: Replace all spwith "sp"

Comment [MM33]: Replace all spwith "sp"

Comment [MM34]: Replace all spwith "sp"

Comment [MM35]: Replace all spwith "sp"

Comment [MM36]: Replace all spwith "sp"

Table 2. Effect of treatments on yield (q/ha)

Treatments	Yield q/ha	Cost of yield Rs./q	Total cost of yield (Rs.)	Total cost cultivation (Rs.)	C:B ratio
T ₀ -Control	152.38	2000	304000	65400	1: 4.64
T ₁ -Neem oil	185.46	2000	370000	67380	1: 5.49
T ₂ -Neem extract	208.23	2000	416000	66400	1: 6.26
T ₃ -Clove oil	174.34	2000	348000	67650	1:5.14
T ₄ -Clove extract	198.56	2000	396000	67000	1;5.90
T ₅ -Neem oil + Clove oil	216.02	2000	432000	67600	1:6.39
T ₆ -Bavistin	233.00	2000	466000	66540	1;7.0
T ₇ -Neem extract+Clove Extract	225.03	2000	450000	66900	1:6.72

Fig 1. Effect of treatments on Disease intensity(%).





Fig 2. Effect of treatments on Yield (q/ha)



The data presented in table 1 and depicted in fig 1 reveals the response of selected treatments on Alternaria leaf spot disease intensity (%) of brinjal at 60, 75 and 90 DAT under field conditions.

Disease intensity (%) at 60, 75, 90 DAT was significantly reduced in the treatment T₆-Bavistin (11.08%, 22.34%, 32.66%) followed by T₇-*Pseudomonas* sp. + Neem extract + Clove extract (12.88, 22.53, 32.37) as compared to other treatments including control.

The data presented in table 2 and depicted in fig 2 reveals the response of selected treatments on

Alternaria leaf spot disease intensity (%) of brinjal at 60, 75 and 90 DAT under field conditions.

Highest yield was obtained in the treatment T₆-Bavistin (236.16q/ha) followed by *Pseudomonas* sp. (229.16q/ha) as compared to other treatments including control.

CONCLUSION

Pseudomonas sp. + Neem extract + Clove extract applied as foliar spray thrice at 15 days interval against Alternaria leaf spot of brinjal significantly reduced disease intensity (%), increased plant height, number of branches, yield and highest benefit cost ratio. Results shows that all the treatments are statistically significant over control and *Pseudomonas* sp. + Neem extract + Clove extract was superior and significant in reduction percent of Alternaria leaf spot of brinjal as compared to control.

REFERENCES

Arancon, N. Q., Edwards, C. A., Atiyeh, R. M. and Metzger, J. D. (2004). Effects of vermicomposts produced from food waste on greenhouse peppers. *Bioresources and Technology*, **93**: 139-144.

Balai, L. and Ahir, R. (2013). Survey and Occurrence of Leaf Spot of Brinjal Caused by *Alternariaalternata*(Fr.) Keisslerin Jaipur District. *Advances in Life Sciences*, **2**(1): 71-72.

Beura, S. K., Mahanta, I. C. and Mohapatra, K. B. (2008). Economics and chemical control of Phomopsis twig blight and fruit rot of brinjal. *Journal of Mycopathological Research*, **46**: 73-76.

Bochalya, S., Shkhawat, S., Singh, R. and Chohan, P. K. (2012). Studies on age of fruit, age of inoculum and inoculum load on development of Alternaria fruit rot of brinjal caused by *Alternariaalternata*. *Annals of Agriculture BioResearch*, **18**: 388-390.

Brahmachari, G. (2004). Neem-An Omnipotent Plant: A Retrospection. *Chemistry and Chemical Biology*, **5**: 408-421.

Choudhary, B. and Gaur, K. (2009). The Development and Regulation of BT Brinjal in India. *International service for the Acquisition of Agri-Biotech Application*, **38**.

Comment [MM37]: Replace all spwith "sp"

Comment [MM38]: Replace all spwith "sp"

Comment [MM39]: Replace all spwith "sp"

Jaiman, R. K., Patel, N. R., Patel, K. D., Agalodiya, A. V. and Patel, P. K. (2013).

Management of Ramularia blight in fennel. *International Journal of Seed Spices*, **3**(1): 50-51.

Kakraliya, S. S., Choskit, D., Pandit, D. and Abrol, S. (2017). Effect of Bio-Agents, Neem

Leaf Extract and Fungicides Against Alternaria Leaf Blight of Wheat (*Triticum aestivum* L.).

Natural Products Chemistry & Research. **5**: 8.

Kumar, R. S. and Arumugam, T. (2013). Phenotypic evaluation of indigenous brinjal types

suitable for rainfed conditions of South India (Tamil Nadu). *African Journal of Biotechnology*,

12(27): 4338-4342.

Kutawa, A. B., Danladi, M. D. and Haruna, A. (2018). Antifungal activity of Garlic (*Allium*

sativum) extract on some selected fungi. *Journal of Medicinal Herbs and Ethnomedicine*. **4**: 12-

Shrivastava, K. P. and Butani, D. K. (1998). Pest management in vegetable, Part-1. *Research*

periodical and book publishing house, Texas, USA, 294.

Wheeler, B. E. J. (1969). An introduction to plant disease. John Wiley, London, U. K. 301.

