

1 **UROGENITAL SCHISTOSOMIASIS AND ASSOCIATED RISK FACTORS AMONG**  
2 **WOMEN IN THE THREE(3) SENATORIAL ZONES OF ANAMBRA STATE**

17

---

**ABSTRACT****Background**

Urogenital Schistosomiasis, a trematode infection (*Schistosoma haematobium*) is endemic to Nigeria. The disease results in urogenital consequences such as cancer and infertility among others.

**Aim of the study**

This study determined the status of urogenital schistosomiasis among women in the three senatorial zones of Anambra State, Nigeria.

**Methodology**

This is a cross-sectional study involving 500 women randomly selected in some selected communities of the 3 senatorial zones of Anambra State. The study was conducted between October 2023 and March 2024, in six communities namely Omogho, Oraifite, Agulu, Achalla, Nsugbe and Awkuzu. 500 urine samples were collected from the 500 women who consented, the urine samples were checked for haematuria using combi 9 dip stick, they were centrifuged for 10 minutes at 1500rpm, the deposit was viewed under x40 microscope objective to detect *S. haematobium* eggs. Data on socio-demographic characteristics and risk factors were obtained through a well-structured questionnaire. Statistical Package for the Social Sciences (SPSS) version 25 was used for analysis, with statistical significance established at p-values less than 0.05.

**Results**

Urogenital schistosomiasis was found in 56(11.2%) of women in the three senatorial zones. Anambra south senatorial zone had the highest prevalence of 22(12.9%), while Anambra Central had the least prevalence 17(9.9%) each. Women of age group 16-20 years had the highest 30(34.5%) urogenital Schistosomiasis infection, and the highest haematuria 11(12.6%), the difference in infection rate according to age is statistically significant  $p < 0.05$ ,  $p = 0.000$ . Women with low educational level had more infection, difference in infection according to educational level is statistically significant  $p < 0.05$ . Women who had water contact through swimming or bathing in the infected water bodies had the highest prevalence 50 (12.4%), other risk factors include nearness to stream, use of infested water bodies as main water source 51(12.8%).

**Conclusion**

The present study indicated that urogenital Schistosomiasis is endemic in Anambra State, MDA should extend to everybody in endemic communities. Continuous health education should be implemented.

**Keywords:** PHCs, CHEW, Female, genital, Schistosomiasis.

20  
21  
22

23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56**INTRODUCTION**


Urogenital schistosomiasis is a significant public health challenge, especially in sub-Saharan Africa caused by *Schistosoma haematobium*. This parasitic disease spreads through contact with fresh water contaminated by snails carrying infective larvae known as cercariae and results in various health problems [2]. Nigeria shoulders a substantial burden of Urogenital Schistosomiasis within sub-Saharan Africa[3]. Around 101 million individuals in Nigeria are at risk due to the endemic spread of the infection across the 36 states [4], mostly school-aged children and adolescents [5][6]

The prevalence of Urogenital Schistosomiasis in Nigeria varies from region to region, with some areas having much higher prevalence [7][8][9][10] than others. One major risk factor for Urogenital Schistosomiasis transmission is engaging in activities which has to do with coming in contact with water bodies where infected snails are present [4]. sociocultural and behavioural factors such as cultural barriers [11][12], misinformation [13], misconceptions and poor health-seeking behaviours [14] perpetuate transmission. Environmental factors ,for urogenital Schistosomiasis epidemiology and distribution include temperature, altitude, rainfall, and land use cover. whereas rainfall provides temporary snail habitats [15], water level, temperature and height, influence the survival and reproduction of intermediate host snails. Socio-demographic factors such as age [16][17] and gender may also affect the prevalence of infection. Children and young adults at school have an increased risk of infection because they interact with water frequently [18]. Additionally, the internal migration of those who have been displaced by flooding, unrest, and insurgencies may aid in the disease's spread [7].

Haematuria, or blood in the urine, and chronic inflammation brought on by eggs stuck in the tissues of the pelvic organs—the bladder, lower uterus, cervix, vagina, prostate gland, and seminal vesicles—are among its characteristic symptoms, others include vaginal discharge, pelvic and abdominal pain, post coital bleeding, and pathologies such as bladder and cervical cancers[19][20][21]. In Anambra State few studies were carried out on urogenital schistosomiasis on women hence this study evaluates the status and risk factors of urogenital schistosomiasis among women in the three senatorial zones of Anambra State.

**MATERIAL AND METHODS****2.1 Study area**

This study was carried out in Anambra state. Anambra state is located in latitude 6.2758 N and longitude 7.0068E, it has a population of 4182,032 according to the 2006 Nigeria census[22]. It has an area of 1774 square meters. Anambra state has a tropical wet and dry or savanna climate with yearly temperature of 28.99c (84.18F).It has about 212.36mm of rain and 243.38 rainy days annually.(weather and climate. Com),(FIG. 1, Map of Anambra State) showing the study areas. The study areas are bounded by streams and rivers where indigenes do their daily chores like bathing and washing. These fresh water bodies provide suitable habitats to snails which are intermediate hosts of Schistosomes.



57  
 58 fig 1 : Map of Anambra State Showing the Selected Local Government Area for the Study (Source: Geography Information  
 System Laboratory, Department of Estate Survey and Geo informatics, Nnamdi Azikiwe University,  
 2023)

61 **2.2 Study Design**

62 The study was carried out at the 3 senatorial zones of Anambra State. Two LGAs were selected from each senatorial  
 63 zone, and one community selected from each LGA. In Anambra North Senatorial Zone of Anambra State, (Oyi and Anambra  
 64 East), in Anambra South senatorial zone, (Orumba North and Ekwusigo) while in Anambra central senatorial zone,  
 65 (Anaocha and Awka North LGAs) were selected. Previous studies implicated these communities as endemic areas for  
 66 urogenital schistosomiasis [27][41][23]. The study was a cross-sectional study conducted from October 2023 to March 2024.  
 67

68 **2.3 Study population**

69 The study population consisted of 500 adolescents and women aged between 16 and 50 years who consented and were  
 70 residents of Agulu-Anaocha; Achalla –Awka North; Oraifite –Ekwusigo; Omogho –Orumba North; Nsugbe- Anambra East  
 71 and Awkuzu-Oyi.  
 72 communities.

73

74 **2.4 Inclusion Criteria**75 Inclusion criteria, all women that volunteered, all women from ages 16-50year and all women who have lived in the  
76 community for at least 10 years.

77

78 **2.5 Exclusion criteria**79 Women who are not aged 16- 50 years, women who have not lived in the community for 10years,and women who did not  
80 consent

81

82

83 **2.6 Sample Size Estimation**84 The sample size for this study was calculated in accordance with Yamane's formula [22]. The formula used for the  
85 calculation is  $n = \frac{N}{1 + N(e^2)}$ 

86 where n =sample size

87 N=total population:

88 From 1991 population census , Number of females in Omogho is 1664:

89 Number of females in Nsugbe 8,314

90 Achalla Females 7017

91 Agulu 25737

92 Oraifite 13,552

93 Awkuzu 14431

94 Total population =70715

95 e=error term at 95% confidence interval which is 0.05

96  $n = \frac{N}{1 + N(e^2)}$ 

97 n= 398 approximately The sample size for the study was 500 individuals

98

99

99 **2.7 Sample collection and parasite egg determination**

100 Participants were provided with 20ml sterile containers strictly labelled with specific identification numbers to avoid mix up.

101 Ten ml mid -stream urine samples were collected from each participant between 10 am and 2.00 pm ,a period known for

102 maximum Schistosome egg extraction[42]. Reagent strip (Meditest Combi 9 test strip, manufactured by Macherey-Nagel

103 GmbH and Company, Germany) was immersed in the urine sample within 60 seconds to check for haematuria. The samples

104 were preserved with 10% formaldehyde, chilled on ice packs and transported to the laboratory of Parasitology and

105 Entomology Department of Nnamdi Azikiwe University. The urine samples were centrifuged at 1500rpm for 10 minutes in a

106 model benchtop centrifuge and sediments were examined at x 40 magnification. [23] *S. haematobium* eggs were identified

107 and confirmed by Prof. CA Ekwunife before they were recorded.

108

109

109 **2.8 Associated Risk factors Data collection**

110 During sample collection, questionnaires were administered too for information on risk factors, participants were given

111 structured questionnaires, the information on their activities was matched with the observed prevalence in order to

112 determine the factor that predisposes them to higher infection.

113

114

114 **2.9 Data Analysis**

115 Data from the study was summarized using tables. Chi- square(x) a test was used to compare the prevalence of

116 urogenital Schistosomiasis with respect to age, senatorial zones etc, the Statistical Package for the Social Sciences

117 (SPSS) version 25 was used for analysis, with statistical significance established at p-values less than 0.05.

118

119

120

121

122

123

124

125

126

127 **3.0 RESULTS**128 **3.1 Demographic data of the sampled women**

129 Out of 500 women that participated in the study, the highest number 204(40.8%),was from the age range 46-50years while  
 130 the least number 5(1.0%) represented the age range of 21-25years(Table1).Other factors are seen in Table 1.

131 **Table 1: Demographic data of respondents:**

<b>Age group</b>	<b>No. examined</b>	<b>%</b>
16-20	87	17.4
21-25	5	1.0
26-30	23	4.6
31-35	20	4.0
36-40	38	7.6
41-45	123	24.6
46-50	204	40.8
Total	500	100%
<b>Location by senatorial zones</b>		
Anambra South	170	34.0
Anambra North	158	31.6
Anambra Central	172	34.4
Total	500	100%
<b>Location by study sites</b>		
Omogho	88	17.6
Oraifite	82	16.4
Nsugbe	78	15.6
Awkuzu	80	16.0
Agulu	73	14.6
Achalla	99	19.8
Total	500	100%
<b>Marital status</b>		
Married	401	80.2
Widowed	46	9.2
Single	103	20.6
Total	500	100%
<b>Education status</b>		
Informal	82	16.4
FSLC	193	38.6
WAEC	177	35.4
Higher school	48	9.6
Total	500	100%
<b>Occupation</b>		
Farming/Fishing	313	62.6
Student	85	17.0
Trading	102	20.4
Total	500	100%

132

133

134

135

136

137

138

139

### 140 3.2 Infection rate of urogenital schistosomiasis by age

141 Prevalence and intensity of *schistosoma haematobium* by age is shown on table 2. The overall *schistosoma haematobium*  
 142 prevalence among the studied individuals was 56(11.2%). However the highest *schistosoma haematobium* prevalence is  
 143 recorded among the age group of 16-20 years which was 30(34.5%), 21-25 age range had no prevalence of *S. haematobium*  
 144 0(0.0%). the difference in infection rate according to age is statistically significant ( $p < 0.05$ ),  $p = 0.000$  as shown on (Table 2).

145 **Table 2 : Infection rate of urogenital Schistosomiasis by age**

Age group (yrs)	Total No. examined	S. haema. (%)	Microhaematuria(%)
16-20	87	30 (34.5)	11(12.6
21-25	5	0 (0)	0(0)
26-30	23	3 (13.0)	2 (8.7)
31-35	20	2 (10.0)	0 (0)
36-40	38	5 (13.2)	0(0)
41-45	123	5 (4.1)	5(4.1)
46-50	204	11 (5.4)	6(2.9)
Total	500	56	24

146

147

### 148 3.3 Infection rate of urogenital schistosomiasis by Location by senatorial zones

149 On prevalence by senatorial zones where Anambra South had the highest prevalence of 22(12.9%) and highest intensity,  
 150 while Anambra Central had lowest prevalence of 17(9.9), Difference due to senatorial zones was not statistically  
 151 significant. ( $p < 0.05$ ).  $p = 0.65$

152 **Table 3a: Infection rate of urogenital Schistosomiasis by Location by senatorial zones**

Location	Toatal No examined	S. haem.+ve (%)	Micro haematuria (%)
Anambra South	170	22(12.9)	12(7.1)
Anambra North	158	17(10.8)	5(3.2)
Anambra Central	172	17(9.9)	7(4.1)
Total	500	56	24 (14.4)

153

154

### 155 3.4 Infection rate of urogenital Schistosomiasis by study sites

156 Prevalence by location by study sites had the highest prevalence of 16(18.2%) at Omogho, while Oraifite had lowest  
 157 prevalence of 6(7.3%), Difference due to study sites was not statistically significant. ( $p < 0.05$ )  $p = 0.178$  see table 3b.

158 **Table 3b Infection rate of urogenital Schistosomiasis by study sites**

Location by sites	Total No examined	S. haem.(%)	Microhaematuria (%)
Omogho	88	16 (18.2)	10(11.3)
Oraifite	82	6(7.3)	0(0)
Nsugbe	78	11(14.1)	2(2.6)
Awkuzu	80	6(7.5)	5(6.3)
Agulu	73	7(9.6)	5(6.8)
Achalla	99	10(10.1)	2 (2.0)
	500	56	24

159

160

161

162

163

164 **3.5: Infection rate of urogenital schistosomiasis by Educational status**

165 Women with Informal education had the highest infection rate 24(29.3) and highest microhaematuria 10(12.2), while WAEC  
 166 had the lowest infection rate.The difference in infection rate due to education was statistically significant.( $p < 0.05$ ),  $p = 0.000$ .  
 167 see table 4

168 **Table 4: Infection rate of urogenital Schistosomiasis by Education status**

Education status	Total No examined	<i>S. haematobium.</i> (%)	Microhaematuria (%)
Informal	82	24(29.3)	10(12.2)
FSLC	193	19(9.8)	10(5.2)
WAEC	177	9(5.1)	4(2.3)
Higher School	48	4(8.3)	0(0)
Total	500	56	24

169

170

171 **3.6: Infection rate of urogenital schistosomiasis by Marital Status**

172 Widows had the highest percentage of infection rate 12(26.1%) and haematuria 7(15.2%).While married women had the  
 173 lowest 30(7.3%) infection rate the difference is statistically significant.  $P = 0.001$ , ( $p < 0.05$ ).see table 5

174 **Table 5 Infection rate of urogenital Schistosomiasis by Marital Status**

Marital Status	Total No. examined	<i>S. haematobium.</i> (%)	Microhaematuria (%)
Married	351	30 (7.3)	3(0.9)
Widowed	46	12 (26.1)	7(15.2)
Single	103	14 (13.6)	14 (13.6)
Total	500	56	24

175

176 **3.7: Infection rate of urogenital Schistosomiasis by Occupation**

177 Women who engage in Farming and fishing had the highest infection rate 44(14.1%) while traders had the lowest infection  
 178 rate 4(3.9%), the difference is statistically significant.( $p < 0.05$ ), $p = 0.016$ .see table 6.

179 **Table 6: Infection rate of urogenital Schistosomiasis by Occupation**

Occupation	Total No. examined	S.haem. (%)	Microhaematuria (%)
Farming/ Fishing	313	44 (14.1)	15(4.8)
Student	85	8(9.4)	4(4.7)
Trading	102	4(3.9)	5(4.9)
	500	56	24

180

181

182

183

184

185

186

187

188

189

190

191 **3.8: Relationship between prevalence and risk factors**

192 The significant risk factors associated with *S. haematobium* infection include frequent contact with infected freshwater  
 193 bodies (rivers/streams), washing/swimming and fishing. As expected, these results indicate a link between water contact  
 194 and infection prevalence. Women who use stream/river as water source had highest infection 12.8% ,Women who swim,  
 195 bath and wash in these water-bodies had highest infection rate 50(12.4%), those living near water body had higher infection  
 196 12.1%,While those that use rain water had lowest infection rate 4.6%.Indiscriminate activities of defecating and urinating  
 197 into water bodies posed a risk for perpetuating transmission of infection. See Table 7

198 **Table 7: Relationship between prevalence and Risk factors**

Source of domestic water source	Yes	No infected	%	NO	No infected	%
River/stream/lake	391	50	12.8	109	6	5.5
Borehole	53	3	5.6	447	53	11.8
Rain water	48	2	4.16	452	54	11.9
Water activity						
Swimming/bath/wash	402	50	12.4	98	6	6.1
paddy farming	447	54	12.1	53	2	3.8
Fishing	471	55	11.7	29	1	3.4
<b>Indiscriminate activity in the water</b>						
Urinating/daefecating	323	54	16.7	177	2	1.1
<b>Water proximity</b>						
Near(<1km)	423	51	12.1	77	5	6.5
Far(>1km)	47	4	8.5	453	52	11.5

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215 **Discussion**

216 The overall prevalence of urogenital Schistosomiasis among women studied in Anambra State was 56(11.2%). This  
217 prevalence of infection was generally lower as compared to what was reported in the study conducted in the Mwanga  
218 district, Kilimanjaro region northern Tanzania among reproductive women where the prevalence was 36% [24] and in Volta  
219 basin of Ghana prevalence was 24.8% [25]. However, the prevalence reported in this study is more or less similar to what  
220 was reported in a study conducted in Sengerema and Misungwi district north-west of Tanzania where the prevalence was  
221 5% [26]. This rate is notably higher compared to recent studies by [27] in Anambra State and indicates the need for revised  
222 intervention strategies. This incidence may have been influenced by the omission of adult women from mass drug  
223 administrations (MDAs) programs, which primarily target elementary school-aged children. According to earlier research  
224 [28], this selective approach impedes control efforts since sick adults act as community reservoirs for the disease. In  
225 comparison to other Nigerian regions where MDAs have effectively covered school-aged children (5–14 years), the overall  
226 prevalence of Urogenital Schistosomiasis in this study is lower than in the following studies- [29][30][31].

227 From this study women aged 16–20 years had the highest prevalence, which may be attributed to their increased  
228 involvement in water-related activities that bring them in contact with infested water bodies, this observation is in accord  
229 with the studies of [32][9][33][23][34]. It is a general characteristic of helminths infections, urogenital schistosomiasis being  
230 one of them, that the prevalence of infection varies significantly from one place to another according to variation in exposure  
231 pattern even in places close to one another [26]. In this study, it was observed that Anambra South senatorial zone had  
232 relatively higher prevalence of infection as compared to other zones. The observed relatively higher prevalence of infection  
233 in the zone is likely due to higher level of exposure to cercarial infested water bodies as a result of high engagement in  
234 activities such as farming and fishing in the rivers, however this difference in infection rate according to location was not  
235 statistically significant  $p > 0.05$ ,  $p = 0.172$ .

236 This study showed that urogenital schistosomiasis was more prevalent among women who had no formal education and  
237 those with primary level of education compared to those who had secondary school and higher school level of  
238 education. ( $p < 0.05$ ). This observation is in line with other studies by [35]. It has been suggested that, education may affect  
239 attitudes and behaviours with individuals with low educational status being more likely to cross a stream or river barefooted  
240 than their more educated counterparts, [36] made similar observation. From this study it was observed that self-awareness  
241 of the disease may account for the relatively lower risk of the disease among women with secondary and higher school  
242 level of education as was observed by [37]. This mandates the need for health education in endemic communities to lower  
243 the overall risk of acquiring the disease by raising community's level of awareness about the disease.

244 From this study high prevalence 50(12.4%) was observed among women with the habit of swimming in rivers, streams and  
245 lakes. This indicates that long duration of hours of water contact was considered as an important risk factor for exposure to  
246 urogenital Schistosomiasis rather than frequency of water contact, this is in line with study of [23] in Anambra State and  
247 [38]. This study, showed that *Schistosoma* infection was found in a large proportion of women 50(12.8%) whose source of  
248 water for domestic use was stream/river/lake. This could be explained on the premises that majority of rural areas from  
249 which the women came from, had no access to protected water sources. They therefore stand the huge risk of *Schistosoma*  
250 infection from their exposure to cercariae infected streams and rivers when accessing these sources to obtain water for  
251 domestic use, this was also observed by [38]. Similar findings were made by [39] who observed that 58.9% of respondents  
252 whose source of water for washing was streams and 42.1% whose source of water for bathing was river had high prevalence  
253 of urogenital Schistosomiasis. [40] also made same observation. It was also observed in this study that defecating and  
254 urinating into water bodies enhance transmission of the infection by infesting the waters with the ova of *Schistosoma*  
255 *haematobium* and *S. mansoni*, these are then ingested by the intermediate snail hosts to perpetuate infection. This is in line  
256 with the studies of [23] and [34] in Anambra State Nigeria.

257 This study has some limitations that need to be considered for a comprehensive understanding of its findings. Firstly, the  
258 study's geographical scope was limited to only six communities of Anambra State although these represent the three  
259 senatorial zones of the state, they may not fully represent the urogenital schistosomiasis situation in other regions of

260 Anambra State, due to the focal nature of the disease. Secondly, the exclusion of males in the study could have resulted in  
261 underestimating the prevalence of urogenital schistosomiasis in these communities. To accomplish total eradication, a more  
262 thorough, all-encompassing MDA program to all sectors of the endemic community, including all males and females, is  
263 advised. This holistic approach will stop the problem of reinfection and transmission of urogenital Schistosomiasis and assist  
264 policymakers and healthcare providers in making decisions about resource allocation.

265

### 266 **Conclusion**

267 The study demonstrates that, despite a notable rise in the praziquantel treatment coverage index, schistosomiasis remains  
268 endemic in Anambra State.

269 This presents a significant health hazard, as the infected women who are not included in the MDAs could play a leading  
270 role in the spread of the disease. However, when compared to findings from earlier research, the observed decline in the  
271 disease's prevalence in the majority of the study locations indicates that recent efforts to combat the illness have been  
272 successful.

273

276

277

283

### 284 **Ethical Approval and Consent to participate**

285 Ethical approval was obtained from Nnamdi Azikiwe University Research and Ethics Committee.  
286 (NAUTH/CS/66/VOL.16/VER.3/231/2022/138).Permission for access to communities was obtained from directors of health  
287 departments of local government areas of the different towns, with introductory letter from the head of department of  
288 Parasitology and Entomology Nnamdi Azikiwe University. key stakeholders including traditional rulers and town union  
289 leaders were sensitized during advocacy visits to the community. Informed consent of participants enrolled in the study was  
290 duly obtained and confidentiality was maintained. Participation was voluntary and participants were free to withdraw from  
291 the study at any time without obligations

292

293

297

298

## 298 **REFERENCES**

- 299 1. Klohe K, Koudou BG, Fenwick A, Fleming F, Garba A, Gouvras A, (2021). A systematic literature review of  
300 schistosomiasis in urban and peri-urban settings. *PLoS Neglected Tropical Diseases*. 15(2):e0008995.
- 301 2. Peletu BJ, Ofoezie IE, Ikwuka AO. Attitude, knowledge, perception, behavioural, cultural and religious practices  
302 influencing transmission of urogenital schistosomiasis in Owena, Kajola and Baiken communities bordering  
303 Owena Reservoir/Dam, Ondo East Local Government Area, Ondo State, Nigeria. *European Journal Med Health*  
304 *Sci*. 2023;5(1):23–30.
- 305 3. Sibanda T, Makwikwi T.(2023).The prevalence and burden of urogenital schistosomiasis: a case study of Mount  
306 Darwin and Makoni districts in Zimbabwe. *All Life*.16(1):1–12

307

308

309

- 310 4. Ezeh CO, Onyekwelu KC, Akinwale OP, Shan L, Wei H. (2019). Urinary schistosomiasis in Nigeria: a 50-year  
311 review of prevalence, distribution and disease burden. *Parasite*. 26:19.  
312
- 313 5. Ekwunife C, Nwaorgu O, Ukonze C, Ukaga C, Ezeunala M. (2014). Urinary schistosomiasis and health education  
314 in Anambra state of eastern Nigeria. *International journal of Infectious Diseases*. 21:89.  
315
- 316 6. Balogun JB, Adewale B, Balogun SU, Lawan A, Haladu IS, Dogara MM. (2022) Prevalence and associated risk  
317 factors of urinary schistosomiasis among primary school pupils in the Jidawa and Zobiya communities of Jigawa  
318 State, Nigeria. *Annual Global Health*. 88(1):71.  
319
- 320 7. Enabulele EE, Platt RN, Adeyemi E, Agbosua E, Aisien MSO, Ajakaye OG. (2021). Urogenital schistosomiasis in  
321 Nigeria post receipt of the largest single praziquantel donation in Africa. *Acta Tropics*. 219:105916.  
322
- 323 8. Njoku JC, Ajayi JA, Pitman SL, Dakul D, Njoku OM. (2014). Urogenital schistosomiasis in females from some  
324 suburban communities of Jos, north-central Nigeria. *IOSR Journal of Pharmaceutical Biology Science*. 9(33):69–  
325 79.  
326
- 327 9. Abubakar B, Abubakar A, Moi I, Gagman H, Aminu MU, Katagum Y. (2022). Urinary schistosomiasis and associated  
328 risk factors among primary school students in the Zaki Local Government Area, Bauchi State, Nigeria. *Dr Sulaiman  
329 Al Habib Medical Journal*. 4:196–204.  
330
- 331 10. Opara KN, Akomalafe RT, Udoidung NI, Afia UU, Yaro CA, Basse BE. (2022). Urogenital schistosomiasis among  
332 primary school children in rural communities in Obudu, Southern Nigeria. *International Journal of Maternal Child  
333 Health AIDS*. 10(1):70–80.  
334
- 335 11. Nwoke BEB, Dozie INS, Nwoke EA, Anosike JC. (2004). Human schistosomiasis and Nigerian environment and  
336 climate change. *Journal Biology Resource of Biotechnology*. 2(1):1–11.  
337
- 338 12. Bolaji OS, Adeyeba OA, Ojurongbe O, Odewale G, Ukaga CN. (2014). Water contact activities and socio-cultural  
339 factors on urinary schistosomiasis in rural area of Osun State, Nigeria. *International Journal Res Applied Natural  
340 Social Sciences*. 2(4):101–6.  
341
- 342 13. Antwi S, Aboah KE, Sarpong CK. (2014). The unacknowledged impact of urinary schistosomiasis in children: 5  
343 cases from Kumasi, Ghana. *Ghana Medical Journal*. 48(4):228–33.  
344
- 345 14. Sacolo H, Chimabari M, Kalinda C. (2018). Knowledge, attitude and practices on schistosomiasis in sub-saharan  
346 Africa: a systematic review. *BMC Infect Dis*. 18:46.  
347  
348  
349
- 350 15. Banhela N, Taylor M, Zulu S, Strabo L, Kjetland E, Gundersen S. (2017). Environmental factors influencing the  
351 distribution and prevalence of *Schistosoma haematobium* in school attenders of iLembe and uThungulu health  
352 districts, KwaZulu Natal Province, South Africa. *South African Journal of Infectious Disease*. 32(4):132–7.  
353
- 354 16. Onyekwere AM, Rey O, Nwanchor MC, Alo M, Angora EK, Allienne JF. (2022). Prevalence and risk factors  
355 associated with urogenital schistosomiasis among primary school pupils in Nigeria. *Parasite Epidemiology  
356 Control*. 2022;18:e00255.  
357
- 358 17. Sumbele IUN, Tabi DB, Teh RN, Njunda AL. (2021). Urogenital schistosomiasis burden in school-aged children in  
359 Tiko, Cameroon: a cross-sectional study on prevalence, intensity, knowledge and risk factors. *Tropical Medical  
360 Health*. 49(1):75.  
361  
362
- 363 18. Ayabina DV, Clark J, Bayley H, Lamberton PL, Toor J, Hollingsworth TD. (2021). Gender-related differences in  
364 prevalence, intensity and associated risk factors of *Schistosoma* infections in Africa: a systematic review and meta-  
365 analysis. *PLoS Neglected Tropical Disease*. 15(11):e0009083.  
366
- 367 19. Ekwunife, C.A., Okafor, F.C. and Nwaorgu, O.C. (2009). Ultrasonographic screening of urinary Schistosomiasis  
368 infected patients in Agulu community, Anambra state, southeast Nigeria. *International Archives of Medicine*, 2, 34.  
369

- 370 20. Mazigo HD, Samson A, Lambert VJ, Kosia AL, Ngoma DD, Murphy R, et al. Female genital schistosomiasis is a  
371 sexually transmitted disease: gaps in healthcare workers' knowledge about female genital schistosomiasis in  
372 Tanzania. *PLoS Global Public Health*. 2022;2(3):e0000059.  
373
- 374 21. Masong MC, Wepnje GB, Marlene NT, Gamba V, Mengue MT, Kouokam E.(2021) Female genital schistosomiasis  
375 (FGS) in Cameroon: a formative epidemiological and socioeconomic investigation in eleven rural fishing  
376 communities. *PLoS Global Public Health*.1(10):e0000007  
377
- 378 22. . National Population Commission. Population of Anambra State. [www.npc.org.ng](http://www.npc.org.ng)2006. Accessed 13 December  
379 2022.  
380
- 381 23. Ekwunife CA,and Okafor F C (2004)Schistosomiasis infection in primary schools in Agulu town of Anambra  
382 state,Nigeria.*Animal Research International* vol.1(3)pp.203-204.  
383
- 384 24. Poggensee G, Kiwelu I, Weger V.(2000). Female genital schistosomiasis of the lower genital tract: prevalence and  
385 disease-associated morbidity in Northern Tanzania. *Journal of Infectious Diseases* 181:1210–3. [PubMed] [Google  
386 Scholar  
387
- 388 25. Hotez PJ, Harrison W, Fenwick A, Bustinduy AL, Ducker C, Mbabazi PS (2019). Female genital schistosomiasis  
389 and HIV/AIDS: reversing the neglect of girls and women. *PLoS Neglected Tropical Diseases*. 13(4):e0007025.  
390
- 391 26. Aribodor OB, Okaka CE, Sam-Wobo SO, Okpala BC, Aribodor DN, Obikwelu EM.(2021) Urinary schistosomiasis  
392 and primary evidence of female genital schistosomiasis among pupils in Nsugbe community, Anambra State,  
393 Nigeria. *Nigerian Journal of Parasitology*. 42(2):394–402.  
394
- 395 27. Ozougwu JC, Imakwu CA, Nwachukwu I, Okeke OP, Uzochukwu CU, *Asian. Journal of Parasitology* 6(3):29–36.  
396
- 397 28. Inobaya MT, Olveda RM, Chau TN, Olveda DU, Ross AG. (2014).Prevention and control of schistosomiasis: a  
398 current perspective. *Res Rep Tropical Medicine*. 5:65–75.  
399
- 400 29. World Health Organization.(2003). *Manual of basic techniques for a health laboratory*. World Health Organization:  
401 Geneva. Second edition. 223–249.  
402
- 403 30. Ojo JA, Adedokun SA, Akindele AA, Olorunfemi AB, Otutu OA, Ojurongbe TA (2021). Prevalence of urogenital and  
404 intestinal schistosomiasis among school children in South-West Nigeria. *PLoS Neglected Tropical Disease*.  
405 15(7):e0009628.  
406
- 407 31. Azoro AV, Onyeanula NI, Egeruo AS, Dike MN, Mbagwu CB.(2022). Endemicity of urinary schistosomiasis in Ihube  
408 Okigwe Imo state. *International Journal of Veterinary Science and Animal Husbandry*. 7(1):23–5.  
409
- 410 32. Zida A, Briegel J, Kabré I, Sawadogo MP, Sangaré I, Bamba S (2016). Epidemiological and clinical aspects of  
411 urogenital schistosomiasis in women, in Burkina Faso, West Africa. *Infectious Disease of Poverty*. 5(1):1–10.  
412
- 413 33. Abdulkareem BO, Habeeb KO, Kazeem A, Adam AO, Samuel UU.(2018). Urogenital schistosomiasis among  
414 schoolchildren and the associated risk factors in selected rural communities of Kwara State, Nigeria. *Journal of*  
415 *Tropical Medicine*.:6913918.  
416
- 417 34. Okoye EP, Ekwunife CA,Onyido AE, Obijiofor EC,Nzekwu CI Nnatuanya IO, OkekeUM, Ude EA (2024).Prevalence  
418 of urogenital schistosomiasis among school age children in Riverine area of Anambra West LGA, Anambra State.  
419 *South Asian journal of parasitology* vol7 page 98-109 Article no SAJP.114878.  
420
- 421 35. Ajibola O, Gulumbe BH, Eze AA, Obishakin E(2018). Tools for detection of schistosomiasis in resource-limited  
422 settings. *Medical Science*. 6(2):39  
423
- 424 36. Ugbomoiko U. S., Ofoezie I. E., Okoye I. C., Heukelbach J.(2010).Factors associated with urinary schistosomiasis  
425 in two peri-urban communities in South-Western Nigeria. *Annals of Tropical Medicine and Parasitology*.  
426 104(5):409–419. doi: 10.1179/136485910X12743554760469  
427
- 428  
429  
430

- 431 37. Wepnje G. B., Anchang-Kimbi J. K., Ndassi V. D., Lehman L. G., Kimbi H. K.(2019). Schistosoma haematobium  
432 infection status and its associated risk factors among pregnant women in Munyenge, south west region, Cameroon  
433 following scale-up of communal piped water sources from 2014 to 2017: a cross-sectional study. BMC Public  
434 Health. 19(1):p. 392. doi: 10.1186/s12889-019-6659-7. [PMC free article  
435
- 436 38. Opara KN, Akomalafe RT, Udoidung NI, Afia UU, Yaro CA, Bassey BE.(2022). Urogenital schistosomiasis among  
437 primary school children in rural communities in Obudu, Southern Nigeria. International Journal of Maternal Child  
438 Health AIDS. 10(1):70–80.  
439
- 440 39. Global Network Neglected Tropical Diseases.(2015). Government of Nigeria Releases New Data on the Prevalence  
441 of Schistosomiasis and Intestinal Worms. SABIN: Vaccine Institute, United States of America. Available from:  
442 Accessed 23 March 2023.  
443
- 444 40. Enabulele EE, Platt RN, Adeyemi E, Agbosua E, Aisien MSO, Ajakaye OG.(2021). Urogenital schistosomiasis in  
445 Nigeria post receipt of the largest single praziquantel donation in Africa. Acta Tropica. 219:105916.  
446  
447
- 448 41. Ndukwe YE, Obiezue RN, Aguzie IO, Anunobi JT, Okafor FC. (2019).Mapping of urinary schistosomiasis in  
449 Anambra state, Nigeria. Ann Glob Health.85(1):52  
450
- 451 42. Yamane T. Statistics: An Introductory Analysis. 2nd ed New York: Harper and Row; 1967. [Google Scholar.  
452  
453
- 454 43. Kloos H, Rodrigues, Rodrigues J,Pereira W. (2006). Combined methods for the study of water contact behavior in a  
455 rural schistosomiasis- endemic area in Brazil. Acta tropica 97(1)31-41.