

Modulatory Effects Of Natron On Biochemical Indices And Cardiac Muscle Gene Expression In Postpartum Albino Rats

ABSTRACT

Peripartum cardiomyopathy is a major public health problem that causes significant morbidity, mortality and huge economic burden among peripartum women. The proposal that natron consumption during postpartum period is involved in the development of peripartum cardiomyopathy requires further scientific elucidation. The current research investigated the effects of natron administration on antioxidant status of cardiac muscles, serum cardiac troponins among postpartum albino rats. The effects of natron administration on gene expressing cardiac troponins were also investigated. Colorimetric, sandwich microplate immunoenzymometric assay was used for troponins estimation. Real Time PCR was used for the gene expression studies. Natron administration reduces oxidative stress at lower dose, while it induces oxidative stress at higher concentration. Also natron administration up regulate cardiac troponins genes and STAT3 genes. Therefore, the current study revealed that natron played a protective role at a low concentration. However, it can cause oxidative stress in the cardiac muscles at higher doses. Thus, oxidative stress in cardiac muscles and modulation of cardiac muscle genes is likely the mechanism by which high natron intake causes PPCM. Further research is recommended in humans. Postpartum women should avoid high intake of natron.

Key words: Peripartum cardiomyopathy, Oxidative stress markers, Biochemical indices, Natron, Albino rats

INTRODUCTION

Cardiovascular diseases such as peripartum cardiomyopathy are the most alarming and leading causes of morbidity and mortality among postpartum women. These are diseases caused by additive and interactive effects of inherited and environmental factors. Substantial evidences show that oxidative stress due to the disturbance in the balance between the productions of reactive oxidants (free radical) such as reactive oxygen species (ROS), generated from consumption of exogenous substances such as natron, and antioxidant defense play a vital role in

biomolecules in postpartum period are very rare and have not been reported in northern Nigeria to date. It is on this basis the present study was designed to investigate the modulatory effect of natron on biochemical indices and cardiac muscle gene expression in postpartum albino rats.

MATERIALS AND METHODS

Chemicals and Reagents

Analytical grade chemicals and reagents were used for this study.

Source of Natron (Kanwa)

The natron was purchased from the Central Market, Sokoto, Nigeria.

Natron (Kanwa): Purchase And Preparation

The natron was made into powder form using mortar and pestle, weighed and stored at room temperature before used.

Natron solutions were freshly prepared, each corresponding to the respective dosage for each group of rats using 1 mL of distilled water.

Experimental Animals

Were purchased and kept in the animal house of the Faculty of Pharmaceutical Science, UsmanDanfodiyo University Sokoto. A total of 28 adult rats, 20 females and 8 males, averagely 11 weeks old and 160g weight was used for the study. They were allowed access to clean water and food *ad libitum*. Thereafter were allowed to acclimatise for a week before the commencement of the experiment. The rats were handled according to rules and regulations of animal's ethics.

Breeding

The rats were grouped into 4, each containing 5 females and 2 males in a cage. They were allowed to breed for the period of 6 weeks. After conception, the females were isolated till they deliver.

Natron Administration

The postpartum rats were randomly regrouped into 4, each containing 5 rats respectively. Natron preparations was administered orally daily for 28 days as shown below:

- **Group 1:** 1.0 mL of distilled water (control)
- **Group 2:** 100 mg/kg body weight of natron.
- **Group 3:** 200 mg/kg body weight of natron.
- **Group 4:** 300 mg/kg body weight of natron.

Sample Collection And Preparation

Blood

The animals were allowed to fast overnight and blood samples was collected from the animals through cardiac puncture into clean labelled test tubes after being anaesthetized with chloroform. The blood was allowed to clot and then centrifuged at 4000 rpm for 10 minutes. The serum was used for the analysis of some cardiac biomarkers.

Cardiac muscle samples

The animals were dissected, and the respective cardiac tissues were removed instantly using a scalpel. 20mg of tissue was homogenized in 0.3mL binding buffer 4(BB4) for gene expression analysis. 50mg of tissues was homogenized in 1mL of extraction solution for antioxidant analysis. The homogenate was centrifuged at $10,000 \times g$ for 15minutes at 4°C.

The supernatant was aspirated into clean sample tubes and stored at -20°C until analysis.

Analyses of Cardiac Muscle Antioxidant Status

Malondialdehyde level was estimated in the cardiac muscle homogenate via the method of Shah and Walker's (1989). The activities of superoxide dismutase (SOD) was assayed using the method of Malstrom *et al.* (1975). Glutathione peroxidase (GPx) activity was measured using enzymatic method described by Ursini *et al.* (1985). Catalase activity was measured using the method of Johansson *et al.* (1988).

Estimation of the Serum Levels of Cardiac Function

The levels of cardiac function proteins: Troponins I, C and T were estimated using Pars BiochemELISA kits following the method of Apple *et al.* (1999).

Gene Expression Studies Using Real Time PCR

- Total RNA was isolated from the cardiac muscles of postpartum Albino rats using transgen[®] isolation kit (Shanghai ZJ Bio-Tech Co., Ltd., Shanghai, China) following the manufacturer's guidelines.
- The RNA was quantified using Nanodrop spectrophotometer (Bioevopeak Nucleic acid analyzer, SP-MUV200F, Shandong, China).
- cDNA synthesis and Real time PCR was performed according to TransScript[®] Green one-step qRT-PCR Supermix kit protocol (AQ211-01, TransGen Biotech, China) on a Rotor-Gene Q 5plex HRM qPCR machine (QIAGEN, Germany).
- Ct values was normalized to house-keeping gene using the $\Delta\Delta C_t$ method.

DATA ANALYSIS

The experimental data were statistically analyzed using SPSS version 25.0 (Inc., Chicago, IL, USA) for windows and expressed as mean \pm standard error of mean. A one-way analysis of

variance (ANOVA) was used to analyze the data followed by a Tukey's *post hoc* test to compare the mean differences among the various groups. The level of significance was set at $p \leq 0.05$.

The Delta-Delta Ct algorithm(Livak method) was used to calculate the log-fold change between Ct values of target genes and reference gene (GAPDH) expressed.

RESULTS

Table 1 Shows cardiac malondialdehyde (MDA) concentration and antioxidant enzyme activity in postpartum albino rats administered natron. There was significant increase ($P < 0.05$) in MDA concentration in group 3 and 4 and decrease in group 2 compared to controls. There was significant decrease ($P < 0.05$) in GPX and SOD activity in group 3 and 4 and increase in group 2 compared with controls. There was significant increase ($P < 0.05$) in CAT activity in all groups administered natron compared to **controls**. Table 2 Shows serum troponin I,C and T concentration in postpartum albino rats administered natron. There was significant decrease in troponin I,C and T in groups administered natron compared to **controls**. Figure 1 Shows the effect of Natron Administration on expression of the TNNT2 gene of postpartum Albino Rats. There was **increase** expression of TNNT2 gene with increase natron concentration. Figure 2 Shows the effect of Natron Administration on expression of the TNNI3 gene of postpartum Albino Rats. There was increase expression of TNNI3 gene with **increase** natron concentration. Figure 3 Shows the effect of Natron Administration on expression of the STAT3 gene of postpartum Albino Rats. There was **increase** expression of STAT3 gene with **increase** natron concentration. Figure 4 Shows the effect of Natron Administration on expression of the TNF- α gene of postpartum Albino Rats. There was **decrease** expression of TNF- α gene in group 3 and 4 compared to **controls**.

Table 1: Cardiac Malondialdehyde (MDA) Concentration **And** Antioxidant Enzyme Activity In Postpartum Rats Administered Natron

Parameter	Group 1(n=5)	Group 2(n=5)	Group 3(n=5)	Group 4(n=5)
MDA(nmol/mL)	0.93±0.05 ^a	0.53±0.04 ^c	0.99±0.07 ^b	1.20±0.08 ^b
GPX(U/mL)	13.61±0.28 ^a	15.40±0.29 ^c	6.81±0.59 ^b	4.59±0.39 ^b
CAT(U/mL)	137.68±12.3 ^a	256.04±25.28 ^b	175.80±16.34 ^b	169.38±17.22 ^b
SOD(U/mL)	0.62±0.05 ^a	0.49±0.06 ^b	0.35±0.04 ^b	0.32±0.04 ^b

Values are mean ± SEM. Value with different superscripts on the same row differ significantly(p<0.05). Group 1: postpartum + 1.0 mL DH₂O (control); Group 2: postpartum + administered 100mg/kg body weight of natron; Group 3: postpartum + administered 200mg/kg body weight of natron; Group 4: postpartum + administered 300mg/kg body weight of natron; MDA: malondialdehyde; GPX: Glutathione peroxidase; CAT: Catalase; SOD: Superoxide dismutase. n : number of participant.

Table 2: Serum Troponin I, C And T Concentration In Postpartum Albino Rats Administered Natron

Parameter	Group 1(n=5)	Group 2(n=5)	Group 3(n=5)	Group 4(n=5)
Troponin I(µg/L)	3.86±0.09 ^a	3.75±0.08 ^b	3.60±0.12 ^c	3.76±0.15 ^b
Troponin C(nm/L)	3.79±0.20 ^a	3.48±0.19 ^b	3.91±0.11 ^c	3.47±0.17 ^b
Troponin T(µg/L)	3.53±0.11 ^a	3.46±0.07 ^b	3.26±0.14 ^b	3.60±0.07 ^c

Values are mean ± SEM. Values bearing different superscripts on the same row differ significantly(p<0.05). Group 1: postpartum + 1.0 mL DH₂O (control); Group 2: postpartum + administered 100mg/kg body weight of natron;

Group 3: postpartum + administered 200mg/kg body weight of natron; Group 3: postpartum + administered 300mg/kg body weight of natron; n : number of participant.

UNDER PEER REVIEW




Figure 1: Effect of natron administration on the expression of TNNT2 gene of postpartum albino rats. Postpartum rats were administered with natron at 100, 200 and 300 mg/kg body weight for 28 days. The Delta-Delta Ct algorithm (Livak method) was used to calculate the log-fold change between Ct values of target genes and reference gene (GAPDH) expressed.




Figure 2: Effect of natron administration on the expression of TNNI3 gene of postpartum albino rats. Postpartum rats were administered with natron at 100, 200 and 300 mg/kg body weight for 28 days. The Delta-Delta Ct algorithm (Livak method) was used to calculate the log-fold change between Ct values of target genes and reference gene (GAPDH) expressed.




Figure 3: Effect of natron administration on the expression of STAT3 gene of postpartum albino rats. Postpartum rats were administered with natron at 100, 200 and 300 mg/kg body weight for 28 days. The Delta-Delta Ct algorithm (Livak method) was used to calculate the log-fold change between Ct values of target genes and reference gene (GAPDH) expressed.




Figure 4: Effect of natron administration on the expression of TNF- α gene of postpartum albino rats. Postpartum rats were administered with natron at 100, 200 and 300 mg/kg body weight for 28 days. The Delta-Delta Ct algorithm (Livak method) was used to calculate the log-fold change between Ct values of target genes and reference gene (GAPDH) expressed.

DISCUSSION

The observed decrease in MDA level at 100 mg/kg body weight of natron compared to control is an indication that ROS generated during post-delivery is counteracted by natron, probably due to the presence of some antioxidant minerals. This suggests that natron at a lower dose could have therapeutic benefits. The increased MDA level at higher doses suggests that the consumption of natron at these doses may be detrimental to the health. Natron at a higher dose is known to induce oxidative stress, which may lead to lipid peroxidation of the cardiomyocytes membranes because MDA is an index of lipid peroxidation. This may be responsible for high level of MDA found in this study at higher dose of natron administration. Ajiboye *et al.* (2015); Imafidion *et al.*

(2016) and Farias *et al.* (2017) have earlier reported increased levels of MDA in albino rats administered with natron. Also studies have shown that high sodium present in natron may be associated with an increased generation of ROS in the cardiac muscles (Imafidionet *al.*, 2016; Kutshiket *al.*, 2018; Bello *et al.*, 2023).

The observed significant decrease in GPX and SOD activity might be due to the overwhelming effect of their activities to mitigate oxidative stress associated with ROS generated from natron and post-delivery OS. It might also be due to the presence of molecules in natron that inhibits its activity or due to disruption of the enzyme's structure by ROS. The increased activity of GPx following administration of 100 mg/kg body weight of natron is an indication of the protective role of natron to counteract the post-delivery OS probably due to the presence of selenium (GPx cofactor) in natron. This may also be responsible for the observed decrease in MDA level at the same concentration. However, the observed increase in MDA level at higher doses of 200 and 300 mg/kg body weight of natron may be due to an increase in ROS generation by natron, which overwhelms the antioxidant enzymes' capacity to neutralize ROS. Overall, the decrease in antioxidant enzyme activities in the animals at a higher dose of natron in this study is consistent with earlier study, which showed a reduction in antioxidant status after natron administration (Saidu *et al.*, 2020; Bello *et al.*, 2023). The observed increase in CAT with an increase in natron concentration might be due to the body compensatory response to mop up an increase in free radicals induced by natron administration. Another possible explanation may be due to the direct effects of natron on the genes of the enzyme. This may suggests the protective role of natron, which stimulates the synthesis of the enzyme. Similar findings were reported by (Saidu *et al.*, 2020; Bello *et al.*, 2023).

The observed decrease levels in troponin I, C and T in groups administered natron compared to **controls** may be due to increased turnover of the troponin molecules in cardiac muscle as a result of oxidative stress induced by natron. **TNNT2, TNNI3** and STAT3 gene expression studies **show increase** expression, TNF- α **shows** decrease expression. This indicates that natron administration affects the gene expression of cardiac troponins in cardiac muscle. The study also indicates that natron consumption down-regulates expression of the inflammatory cytokine TNF- α . These findings may suggest that the observed increase in the expression of TNNT2, TNNI3 and STAT3 genes could be a compensatory response to cardiac demand of cardiac proteins (troponins).

Conclusion

The current study revealed that natron played a protective role at a low concentration. However, it can cause oxidative stress in the cardiac muscles at higher doses. Thus, oxidative stress in cardiac muscles and modulation of cardiac muscle genes **is** likely the **mechanism** by which high natron intake causes PPCM.

Recommendations

1. Further research is recommended in higher animals such as humans.
2. Postpartum women should avoid high dose intake of natron.

REFERENCES:

Agnieszka, Jablonska., Branislav, Zagraban., Christoph, Neumayer., and Wolf, Eilenberg. (2021). Polymorphisms in the IL-6 and TNF- α gene are associated with an increased risk of abdominal aortic aneurysm. *International Journal of Cardiology*, **329**: 192-197.

- Ajiboye, TO., Komolafe, YO., Yakubu, MT., and Ogunbode, SM. (2015). Effects of trona on the redox status of cellular system of male rats. *Toxicology and Industria. Health*, **31(2)**: 1–9.
- Apple, FS., Christenson, RH., Valdes, RJ., Andriak, AB., Duh, S., Feng, YJ., Saeed, AJ., Johnson, NJ., Koplen, B., Mascotti, K., and Wu, A. (1999). Simultaneous rapid measurement of whole blood myoglobin, creatine kinase MB and cardiac troponin I by the Triage cardiac panel for detection of cardiac infarction. *Clinical Chemistry***45**: 199–205.
- Avalle, L., Camporeale, A., Morciano, G., Carocchia, N., Ghetti, E., Orecchia, V., Viavattene, D., Giorgi, C., Pinton, P., and Poli, V. (2018). Stat3 localizes, acting as a gatekeeper for mitochondrion Ca^{2+} fluxes and apoptotic responses. *Cell Death and Differentiation*,**98**: 42–50.
- Avasthi, AS., Jawaid, SA., Jain, S., Bhatnagar, M., Purkayastha, S., and Ghosal, S. (2014). Free radical scavenging and antioxidant impact of Indian medicinal plants extracts on H_2O_2 mediated oxidative stress on human erythrocyte. *American Journal of physical and clinical therapy*, **2(9)**: 1052–1069.
- Bello, HA., Saidu, Y., Abubakar, M.K., Wali, U., Abbas, A.Y., and Muhammad S.A. (2023). Correlation between Oxidative Stress Markers and Biochemical Indices of Cardiac Function of Postpartum Albino Rats Administered Natron. *International Journal of Veterinary Science*, **12(3)**: 347- 352.
- Bononi, A., Giorgi, C., Patergnani, S., Larson, D., Verbruggen, K., Tanji, M., Pellegrini, L., Signorato, V., Olivetto, F., and Pastorino, S. (2017). Bap1 regulates ip3r3-mediated Ca^{2+} flux to mitochondria suppressing cell transformation. *Nature*, **546**: 549–553.

- Christopher, N. Toepfer., Giuliana, G. Repetti., Amanda, C. Garfinkel., and Jourdan, F. Ewoldt. (2020). Genetic Studies of Hypertrophic Cardiomyopathy in Singaporeans Identify Variants in *TNNI3* and *TNNT2* That Are Common in Chinese Patients. *Genomic and Precision Medicine*, **13**: 424–434.
- Chinedum, OK., Ndukaku, OY., Emmanuel, OI., and Ndubuisi OT. (2014). Effect of potash on the hearts of rabbits. International Organization of Scientific Research (IOSR), *Journal of Pharmacy and Biological Sciences*, **9**: 33-38.
- Dambazau, SM., Saidu, Y., Bilbis, LS., Idris, A., Dogara, HI., and Umar, UA. (2016). Lipid profiles and liver function parameters of postpartum rats administered dry lake salt (kanwa). *Journal of Scientific Research and Reports*, **10**: 1-8.
- Fangzhou, C. Y., and Zhao, Huizhao, Chen. (2018). MicroRNA-98 reduces amyloid β -protein production and improves oxidative stress and mitochondrial dysfunction through the Notch signaling pathway via HEY3 in Alzheimer's disease mice. *International Journal of Molecular Medicine*, **4**: 91-102.
- Farias, J. G., Molina, V. M., Carrasco, R. A., Zepeda, A. B., Figueroa, E., Letelier, P., and Castillo, R. L. (2017). Antioxidant therapeutic strategies for cardiovascular conditions associated with oxidative stress. *Nutrients*, **9**: 1-23.
- Gomes, AV., and Potter, JD. (2004). Cellular and molecular aspects of familial hypertrophic cardiomyopathy caused by mutations in the cardiac troponin I gene. *Molecular and Cellular Biochemistry*, **263(1-2)**: 99-114.

- Guadagnin, E., Mazala, D., and Chen, Y.W. (2018). Stat3 in skeletal muscle function and disorders. *International Journal of Molecular Science*, **19**: 2265.
- Gupta, D. and Wenger, N.K. (2018). Peripartumcardiomyopathy: Status 2018. *Wiley Clinical Cardiology*, **41**: 217-219.
- Hughes, K., and Watson, C.J. (2018). The multifaceted role of stat3 in mammary gland involution and breast cancer. *International Journal of Molecular Science*, **19**: 1695.
- Imafidion, K. E., Egberanmwun, I. D., and Omoregie, I. P. (2016). Toxicological and biochemical investigations in rats administered “kaun” (trona), a natural food additive used in Nigeria. *Journal of Nutrition and Intermediary Metabolism*, **6**: 22-25.
- Kankara, SS., Ibrahim, MH., Mustafa, M., and Go, R. (2015). Ethnobotanical survey of medicinal plants used for traditional maternal health care in Katsina State, Nigeria. *South African Journal of Botany*, **97**: 165–175.
- Kasembeli, M.M., Bharadwaj, U., Robinson, P., and Twardy, D.J. (2018). Contribution of stat3 to inflammatory and fibrotic diseases and prospects for its targeting for treatment. *International Journal for Molecular Science*, **19**: 2299.
- Kaur, S., and Mondal, P. (2014). Study of total phenolic and flavonoid content, antioxidant activity and antimicrobial properties of medicinal plants. *Journal of Microbiology Experiment*, (1): 1–6.
- Kong, YM., Baqar, S., Jerums, G. and Ekinici, E. I. (2016). Sodium and its role in cardiovascular disease - the debate continue. *Frontiers in Endocrinology* **7**: 164.

- Kutshik, R. J., Idogun, F. O., Ujah, F. E. and Gotom, S. S. (2018). Comparative investigation of the effects of kanwa (trona) and tokansenyi (plant potash) on liver and kidney of albino rats. *International Journal of Biological and Chemical Sciences*, **12**: 422-430.
- Laudisi, F., Cherubini, F., Monteleone, G., and Stolfi, C. (2018). Stat3 interactors as potential therapeutic targets for cancer treatment. *International Journal of Molecular Science*, **19**: 1787.
- Leila, Saremi., Marziyeh, Shafizadeh., Emran Esmaeilzadeh., and Mohammad, Ebrahim, Ghaffari. (2021). Assessment of IL-10, IL-1 β and TNF- α gene polymorphisms and health controls. *Molecular Biology Reports*, **48 (3)**: 2285-2290.
- Malena, P. P., Polyxeni, G., Aggeliki, G., and Iakovos, A. (2019). A case report of recessive restrictive cardiomyopathy caused by a novel mutation in cardiac troponin I (TNNI3). *BMC Medical Genetics* **20(1)**: 98-112 .
- Masayasu, M., and Hiroshi, Y. (1979). A simplified assay method of superoxide dismutase activity for clinical use. *International Journal of Clinical Chemistry*, **92(3)**: 337-342.
- Megan, Beetch., Cayla, Boycott., Sadaf, Harandi-Zadeh., Tony, Yang., and Benjamin, J. E. M. (2021). Pterostilbene leads to DNMT3B-mediated DNA methylation and silencing of OCT1- targeted oncogenes in breast cancer cells. *The Journal of Nutritional Biochemistry*, **98**: 108815.
- Mohammed, CB., Chahid, B., Fatima, B., Fatima-Zohra, S., Meriem, B., and Farid, C. (2014). Antioxidant activity, phenolic and flavonoid content in leaves, flowers, stems and seeds of mallow (*Malvasylvestris*L.) from North Western of Algeria. *African Journal of Biotechnology*, **13(3)**: 486–491.

- Moliterno, A.R., Ginzburg, Y.Z., and Hoffman, R. (2021). Clinical insights into the origin of thrombosis in myeloproliferative neoplasms. *Blood*,**137(9)**: 1145-1153.
- Muhammad, AS. Saidu, Y., Bilbis, LS., Onu, A. Isezuo., SA. and Sahabi, S. (2014). Effect of dried lake salt (kanwa) on lipid profile and heart histology of female Albino rats. *Nigerian Journal Basic Applied Science*, **22(3&4)**: 73–78.
- Nour, Hamad., Karem, H. Alzoubi., Samer, F. Swedan., Omar, F. Khabour., and Khalid EL-Salem. (2021). Association between tumor necrosis factor alpha and lymphotoxin alpha gene polymorphisms and migraine occurrence among Jordanians. *Neurological Sciences*, **42 (9)**: 3625- 3630.
- Ojiako, O. A., Chikezie, P. C., and Ogbuji, AC. (2015). Radical scavenging potentials of single and combinatorial herbal formulations in vitro. *Journal Traditional and Complimentary Medicine*,**2**:1–7.
- Padmanabhan, P. and Jangle, SN. (2012). Evaluation of DPPH radical scavenging activity and reducing power of four selected medicinal plants and their combinations. *International Journal Pharmaceutical Science, Drug and Disease*, **4(2)**: 143–146.
- Philip, DA., Julie, AA., Robert, DB., Peter, MAC., Bartolome, RC., Nicholas, JC., Courtney, C. and Ian, JD. (2018). Cardiac troponin I and cardiovascular risk in patients with chronic obstructive pulmonary disease. *Journal of the American College of Cardiology*,**72**: 126-137.
- Rahana, Y. P., Hendrik, J. H., Michelle, M., and Albert, V. R. (2020). Increased consumption precedes contractile dysfunction in hypertrophic cardiomyopathy caused by pathogenic TNNT2 gene variants. *Journal of American Heart Association*,**9 (8)**: 015316.

- Rincon, M., and Pereira, F.V. (2018). A new perspective: Mitochondrial stat3 as a regulator for lymphocyte function. *International Journal of Molecular Science*, **19**: 1656.
- Ronner, L., Podoltsev, N., Gotlib, J., Heaney, M.L., Kuykendall, A.T., and Connell, C.O. (2020). Persistent leukocytosis in polycythemia vera is associated with disease evolution but not thrombosis. *Blood*, **135 (19)**: 1696-1703.
- Saidu, H., Kabir, A., Ndice, N., Yau, JA. and Abdullahi, UM. (2018a). Prevalence and characteristics of peripartum cardiomyopathy among women with cardiac failure referred for echocardiography in a tertiary hospital in northern Nigeria. *Journal of Biosciences and Medicines*, **6**: 95-104.
- Saidu, Y., Usman, M. J., Isa, S. A., Isezuo, S. A., Bilbis, L. S., Sahabi, S. M., Bello, S. and Muhammad, S. A. (2018b). Biochemical and histological changes in the heart of post-partum rats exposed to natron. *Indian Heart Journal*, **70**: 886-892.
- Saidu, Y., Olaosebikan, R. O., Isa, S. A., Malami, I. and Muhammad, S. A., (2020). Effect of natron administration on the antioxidant status and lipid profile of rats. *Journal of Food Science*, **85**: 4033-4038.
- Shah, JK. and Walker's, AM. (1989). Quantitative determination of MDA. *Scientific journal of Biochemistry and Biophysics*, **11**: 207-11.
- Spitz, D. R., and Oberley, L. W., (1989). An assay for superoxide dismutase activity in mammalian tissue homogenates. *Analytical Biochemistry*, **179(1)**: 8-18.

Su, Y.L., Banerjee, S., White, S.V., and Kortylewski, M. (2018). Stat3 in tumor-associated myeloid cells: Multitasking to disrupt immunity. *International Journal Molecular Science*,**19**: 1803.

Wang, Li. Shi., and Qinghai, Li. Kai. (2019). Oral administration of Ginkgolide B alleviates hypoxia induced neuronal damage in rat hippocampus by inhibiting oxidative stress and apoptosis. *Iranian Journal of Basic Medical Sciences*, **22(2)**: 140-145.

Wen, Liu., Zongkai, Wei., Yanfen, Zhang., Yan, Liu., Ruocen, Bai., Chunyan, Ma., Jun, Yang., and Dandan, S. (2021). Identification of three novel pathogenic mutations in sarcomere genes associated with familial hypertrophic cardiomyopathy based on multiomics study. *International journal of Clinical Chemistry*,**520**: 43-52.