

Study of Genetic Variability, Heritability and Genetic Advance in maize (*Zea mays* L.) inbred lines

ABSTRACT

The present experiment was conducted to study genetic parameters for ten traits viz., days to 50% anthesis and days to 50% silking, plant height, ear height, ear length, ear girth, kernel rows, kernels per row, 100 grain weight and total grain yield per hectare during kharif and rabi seasons of 2023-24. Using an Alpha Lattice Design, 150 inbred lines were planted in two replications. Analysis of variance revealed significant differences across all traits with phenotypic coefficients of variation (PCV) generally higher than genotypic coefficients of variation (GCV). The highest PCV and GCV were observed for ear height (kharif: 27.59, 23.48; rabi: 25.03, 20.94) and total grain yield (kharif: 51.26, 47.60; rabi: 38.85, 38.65). Broad-sense heritability was highest for total grain yield (Kharif-86.34, rabi-88.97) in the both seasons. The high heritability of these traits indicates their potential for successful transmission to offspring, enabling efficient selection. High heritability combined with high genetic advance was observed for ear height and total grain yield in both seasons, suggesting these traits as promising targets for improvement through selection driven by additive genetic factors.

Key words: Heritability, Variability, Maize, yield.

INTRODUCTION

Maize (*Zea mays* L.) is one of the most versatile emerging crops with wide range adaptability for diverse agro-climatic conditions. Globally, maize is known as the "queen of cereals" due to its highest genetic yield potential among cereals. It is widely used for feed and starch-making, and is grown in at least 171 countries across the globe on an area of 201.98 Mha. Among major crops, it has the highest global production of 1162 Mt, with average productivity of 5755 kg/ha [1]. "In India, Maize ranks as the third most important cereal crop, followed by rice and wheat, and contributes approximately 10% to the total food grain production in the country. India was the fifth largest producer of maize globally, with a share of 2.59% in world production" [2]. "It is the primary energy source and a major staple food crop for humans, and feed and fodder crop for livestock, in many countries across Africa, Latin America and Asia, including India. Apart from food, feed and fodder-based industries, it also has multi-faceted uses as raw material in industries such as biofuel and starch. Thus, it occupies a prominent position in global agriculture and trade" (Yathish et al. [3]).

"Heterosis largely depends on the genetic divergence between the parental lines, and selecting genetically diverse parents can be achieved using biometrical methods. In maize breeding programs, the primary goal is to enhance economically important traits while preserving a sufficient level of variability" [4]. To increase the genetic diversity of germplasm, understanding the existing genetic variability within the material is crucial. Genetic variability, which represents heritable differences among genotypes or cultivars, is essential for the success of a long-term breeding program. Zhang *et al.* [5] demonstrated "the importance of quantifying genetic variability among maize

cultivars grown in an area before initiation of breeding programme. Genetic variability is essential for systematically enhancing such complex traits, which can be assessed through phenotypic and genotypic coefficients of variation". Keeping this in view, the present study was conducted to determine the genetic variability, heritability and genetic advance in maize inbred lines which is of great importance for the plant breeders to initiate the breeding programme.

MATERIALS AND METHODS

The present study was undertaken at Winter Nursery Centre, ICAR- Indian Institute of Maize Research, Rajendra Nagar, Hyderabad during kharif and rabi seasons of 2023-24. The experimental material consists of 150 maize inbred lines and these were sown in two replications with Alpha Lattice Design. Each genotype was planted in a single row, 3 meters' length in each replication, with a spacing of 60 cm between rows and 20 cm within rows. The crop was cultivated with the recommended agricultural practices.

Observations were recorded for ten quantitative traits viz., Days to 50 percent tasseling, Days to 50 percent silking, Plant height (cm), Ear placement height (cm), ear length (cm), ear girth (cm), Number of kernel rows per cob, Number of kernels per row, Grain yield per hectare(kg/ha), and 100-grain weight(grams).

Heritability in the broad sense (h^2_{bs}) and genetic advance were calculated using the formula provided by Johnson *et al.* [6]. The genotypic and phenotypic coefficients of variation were determined according to the methods of Burton [7]. The estimates of PCV and GCV were categorized as low (0-10%), moderate (10-20%), and high (>20%) based on the classification by Sivasubramanian and Madhavamenon[8]. The analysis was done using R 4.4.1-win software.

RESULTS AND DISCUSSION

Analysis of variance

The analysis of variance revealed significant differences among the genotypes for all the traits viz., days to 50% anthesis and days to 50% silking, plant height, ear height, ear length, ear girth, kernel rows, kernels per row, 100 grain weight and total grain yield per hectare in both seasons (Table 1), indicating the presence of genotypic variation among the evaluated genotypes. There is an opportunity to select desirable genotypes with enhanced yield component traits, which may perform well and result in increased yield. These findings are in consistent with the results of previous researchers such as Chandel *et al.*[9], Sinha and Thakur [10], Meena *et al.* [11].

Variability and Heritability

The variability of ten quantitative traits in 150 maize inbred lines was assessed using phenotypic coefficient of variation (PCV), genotypic coefficient of variation (GCV), heritability and genetic advance as a percentage of the mean (Table 2).

In this study, the Phenotypic Coefficient of Variation (PCV) was higher than the Genotypic Coefficient of Variation (GCV) for most of the quantitative traits (Table 2) in both the kharif and rabi seasons of 2023-24. Specifically, the difference between PCV and GCV was more pronounced for ear length, kernels per row, and test weight in kharif 2023, and for plant height, ear height, ear length, and kernels per row in rabi 2023-24. This suggests that environmental factors had a greater impact on the phenotypic expression of these traits compared to others.

UNDER PEER REVIEW

Table 1. Analysis of Variance for Yield and Yield-Contributing Traits in Maize

S.No.	Character/trait	Mean sum of squares							
		Replication (Df:1)		Genotypes (Df:149)		Blocks (Df:14)		Error (Df:135)	
		Kharif 2023	Rabi 2023-24	Kharif 2023	Rabi 2023-24	Kharif 2023	Rabi 2023-24	Kharif 2023	Rabi 2023-24
1	Daysto50percent Anthesis	38.88	6.45	30.80**	28.80**	4.20	7.18	5.70	6.27
2	Daysto50percent silking	19.76	29.45	31.85**	26.70**	5.10	4.48	9.32	7.32
3	Plantheight(cm)	82.16	122.24	616.9**	611.77**	136.81	210.63	128.07	172.90
4	Earplacementheight(cm)	1.37	17.76	223.94**	197.72**	50.88	40.11	34.21	34.92
5	Ear length(cm)	6.39	3.91	5.88**	5.44**	2.35	1.99	2.07	2.38
6	Ear girth(cm)	2.67	0.05	0.27**	0.2**	0.165	0.07	0.11	0.08
7	KernelrowsPer Ear	0.17	2.08	4.19**	4.00**	0.86	1.33	1.26	1.24
8	Number of Kernel Per Row	19.25	8.67	29.00**	26.02**	8.82	5.23	7.95	8.34
9	100grainweight(g)	73.01	8.33	26.16**	24.85**	9.58	6.21	9.60	5.16
10	Total Grainyield Per Hectare	27343	46995	525865**	347401**	52156	15339	37130	20273

**Significantat1%level

*Significantat 5%level

S. No.	Character	Mean		Range				Coefficient of Variation				Heritability (broad sense) $h^2(bs)$ (%)		GAM	
				Minimum		Maximum		PCV (%)		GCV (%)					
		K	R	K	R	K	R	K	R	K	R	K	R	K	R
1	DA	60.88	72.00	50.00	59.00	75.00	80.00	7.00	5.81	5.83	4.66	69.40	64.00	9.98	7.69
2	DS	62.88	73.97	51.00	59.00	76.00	82.00	7.18	5.57	5.38	4.20	56.21	56.94	8.31	6.50
3	PH	126.06	122.92	62.00	182.00	192.66	73.00	15.31	16.11	12.39	12.05	65.44	55.93	20.64	18.56
4	EH	41.29	43.08	15.00	19.00	72.66	78.00	27.59	25.03	23.48	20.94	72.43	69.98	41.17	36.08
5	EL	11.06	11.31	5.66	6.60	16.50	18.06	18.06	17.49	12.43	10.93	47.42	39.09	17.64	14.08
6	EG	2.63	2.81	1.26	1.46	3.73	3.93	16.87	13.61	10.76	8.64	40.70	40.27	14.14	11.29
7	KR	11.75	11.53	8.00	8.00	17.33	18.00	14.00	14.04	10.37	10.16	54.85	52.39	15.82	15.15
8	KPR	20.25	18.85	8.00	10.00	32.33	35.33	21.25	21.98	15.99	15.76	56.61	51.43	24.78	23.29
9	TWT	20.93	19.70	10.00	12.00	36.00	32.00	20.20	19.66	13.74	15.92	46.30	65.58	19.26	26.50
10	GYHA	1036.00	1103.45	76.19	180.56	2661.90	2380.95	51.26	38.85	47.60	36.65	86.34	88.97	91.00	71.21

Table 2. Genetic Variability Parameters for Yield and Yield-Contributing Traits in Maize

PCV=Phenotypic coefficient of variation

GCV=Genotypic coefficient of variation

DA: Day to 50 percent Anthesis, DS: Day to 50 percent Silking, PH: Plant height, EH: Ear Placement Height, EL: Ear Length, EG: Ear Girth, KR: Kernel Rows Per Ear, KPR: Kernels Per Row, TWT: 100 Grain Weight, GYHA: Total Grain Yield Per Hectare.

K: Kharif 2023, R: Rabi 2023-2024.

The highest phenotypic and genotypic coefficients of variation (PCV and GCV) were observed for ear height (kharif: 27.59, 23.48; rabi: 25.03, 20.94) and total grain yield (kharif: 51.26, 47.60; rabi: 38.85, 38.65), which indicates the presence of genetic variability in the studied material. High PCV and GCV values for grain yield and ear height in maize were also reported by Singh *et al.* [12], Abiramiet *al.* [13], and Bharathi *et al.* [14]

Moderate GCV and PCV values were observed for the traits of plant height (GCV: kharif-12.39, rabi-12.05; PCV: kharif-15.31, rabi-16.11), which align with similar findings reported by Pradhan *et al.* [15]. The traits of ear length (GCV: kharif-12.43, rabi-10.93; PCV: kharif-18.06, rabi-17.49) and kernel rows per ear (GCV: kharif-10.37, rabi-10.16; PCV: kharif-14.00, rabi-14.04) also exhibited moderate GCV and PCV values in both seasons. These results are consistent with the experimental findings of Barathiet *al.* [14].

Low GCV and PCV were observed for the traits of days to 50% anthesis (GCV: kharif-5.83, rabi-4.66; PCV: kharif-7.00, rabi-5.81) and days to 50% silking (GCV: kharif-5.38, rabi-4.2; PCV: kharif-7.18, rabi-5.57) in both seasons. The minimal difference between genotypic and phenotypic coefficients of variation suggests that environmental factors had little effect on the expression of these traits. These findings are consistent with the experimental results of Barathiet *al.* [14] and Ellandulaet *al.* [16].

Higher broad-sense heritability of traits indicates that a larger proportion of their variation is passed on to offspring. In our study, broad-sense heritability showed the highest values for total grain yield (kharif-86.34, rabi-88.97), followed by ear height (kharif-72.43, rabi-69.98) and Days to 50% anthesis (kharif-69.4, rabi-64) in the both seasons. Additionally, plant height (65.44) in the kharif season and test weight (65.18) in rabi also showing the high heritability. The traits exhibited high heritability in this study, indicated the minimal environmental influence and suggesting that phenotypic selection would be effective for their improvement. Higher heritability indicates the presence of additive gene action, making selection effective in early generations. Similar findings have been reported by Vashisthaet *al.* [17] for traits such as 100 grain weight, days to silking, cob length, plant height, and ear height in maize. Magar *et al.* [18] also reported high heritability for grain yield per plant and 100 grain weight, consistent with our findings. Rafiq *et al.* [19] similarly found high heritability for plant height, ear height, grain yield per plant, grains per row, ear length, and 100 grain weight. Ellandulaet *al.* [17], who also reported high heritability for ear height, days to 50% anthesis, and test weight which were aligned with our results.

Moderate heritability was observed for days to 50% silking (kharif-56.21, rabi-56.94) in both seasons which aligns with findings reported by Ellandulaet *al.* [16] and Gokulakrishnanet *al.* [20]. Similarly moderate heritability was observed for ear length (kharif-47.42 rabi-39.09) in both seasons which were consistent with the experimental findings of Thakur *et al.* [21]. Moderate heritability for Kernel rows (kharif-54.85, rabi-52.39), and kernels per row (kharif-56.61, rabi-51.43) in both kharif and rabi seasons showed results similar to those reported by Suresh *et al.* [22], suggesting

a significant influence from environmental factors. These findings indicate the presence of non-additive gene actions, which may limit the potential for improvement through selection in these traits.

Genetic advance as a percentage of the mean indicated that these traits are predominantly influenced by additive genetic factors. Utilizing heritability estimates alongside genetic advance enhances the reliability and utility of selection procedures. In the present study, genetic advance was calculated and expressed as a percentage of the mean for all traits (Table 2). High genetic advances were observed for total grain yield (kharif-91,rabi-71.21), followed by ear height (kharif-41.17,rabi-36.08) and number of kernels per row (kharif-24.78,rabi-23.29) in both the seasons. Bharathi *et al.* [14] reported high genetic advance values for total grain yield and ear height. The high genetic advance values for kernels per row observed in the study align with the findings from Thakur *et al.* [21] and Wedwessen *et al.* [23]. Additionally, plant height (20.64) in the kharif season also showed the high genetic advance.

High heritability combined with high genetic advance was observed for traits like ear height and total grain yield in both the kharif and rabi seasons. Additionally, high heritability coupled with high genetic advance was noted for test weight in rabi season only. These findings suggest that these traits are promising targets for improvement through selection, likely driven by additive genetic factors. Kinfe and Tsehaye [24] and Bhiusa *et al.* [25] reported high genetic advance along with high heritability for grain yield. Similarly, Bekele and Rao [26] documented high genetic advance and higher heritability for 100-seed weight, which aligns with the findings of this study. The results emphasize that, these traits can be enhanced either through selection methods or by progeny selection techniques.

The minimal genetic advance observed for days to 50% anthesis (kharif-9.98, rabi-7.69) and days to 50% silking (kharif-8.31, rabi-6.50) suggests that these traits are primarily governed by non-additive gene action. Ellandula *et al.* [16] and Gokulakrishnan *et al.* [20] reported low genetic advance values for days to 50% silking, and Khan *et al.* [27] for 50% anthesis which were in consistent with the present study. This indicates that these traits are highly influenced by environmental factors, making selection strategies less effective for their improvement.

CONCLUSION

The analysis of variance indicated the significant variation among the inbred lines, with phenotypic coefficients of variation (PCV) exceeding genotypic coefficients of variation (GCV) for all traits. The highest PCV and GCV were observed for ear height and total grain yield during both seasons, demonstrating substantial variability among the genotypes. This suggests that direct selection will be effective for improving these traits. High heritability combined with high genetic advance for ear height and total grain yield suggests these traits are governed by additive gene action, making them heritable and fixable, and thus, phenotypic selection for these traits will be beneficial.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

REFERENCES

1. FAOSTAT(2020) Crops and livestock products. Available at <https://www.fao.org/faostat/en/#data/QCL>
2. APEDA, 2022. https://apeda.gov.in/apedawebsite/SubHead_Products/Maize.htm
3. Yathish KR, Karjagi CG, Gangoliya SS, Gadag RN, Mallikarjuna MG, Sekhar JC, Das AK, Soujanya PL, Kumar R, Singh A, Singh SB. Development of low-phytate maize inbred lines through marker-assisted introgression of Ipa1. *Crop and Pasture Science*. 2023 Mar 14;74(9):843-55.
4. Hallauer AR, Sears JH. Changes in Quantitative Traits Associated with Inbreeding in a Synthetic Variety of Maize 1. *Crop Science*. 1973 May;13(3):327-30.
5. Zhang ZX, Zheng YL, Li JS, Liu JL. Allozyme polymorphism and relationships to quantitative traits: diversity of 10 local varieties. *Maize Gen. Coop.* 1995. *Newsl.*, 69: 138-142.
6. Johnson HW, Robinson HF, Comstock RE. Estimates of genetic and environmental variability in soybeans.
7. Burton, G.W., 1952. Quantitative inheritance in grasses. *Proceedings of the 6th International Grassland Congress*. Pennsylvania State College, USA, 23 August 1952. 277-283.
8. Sivasubramanian, P and Menon, P.M. 1973. Inheritance of short stature in rice. *Madras Agricultural Journal*. 60: 1129-1133.

9. Chandel U, Kumar D, Guleria SK. Combining ability effects and heterotic grouping in newly developed early maturing yellow maize (*Zea mays* L.) inbreds under sub-tropical conditions. *Electronic Journal of Plant Breeding*. 2019 Sep 30;10(3):1049-59.
10. Sinha SK, Thakur D. Estimation of Combining Ability in Maize (*Zea mays* L.) using Line x Tester Analysis *International Journal of tropical Agriculture*. 2017;35(4):1035-1046.
11. Meena MK, Singh R, Meena HP. Genetic variability, heritability and genetic advance studies in newly developed maize genotypes (*Zea mays* L.). *The Bioscan*. 2016;11(3):1787-91.
12. Singh N, Singh J, Kaur L, Sodhi NS, Gill BS. Morphological, thermal and rheological properties of starches from different botanical sources. *Food chemistry*. 2003 May 1;81(2):219-31.
13. Abirami, S, Vanniarajan C. and Arumugachamy, S. 2005. Genetic variability studies in maize (*Zea mays* L.) germplasm.
14. Bharathi P, Ravikesavan R, Yuvaraja A, Boopathi NM, Iyanar K. Genetic variability and correlation in maize inbred lines under irrigated and moisture stress condition. *Electronic Journal of Plant Breeding*. 2021;12(3):928-33.
15. Pradhan P, Thapa B, Ghosh A, Subba V, Sahu CR, Kundagrami S. Genetic variability assessment on yield attributing traits in maize (*Zea mays* L.) inbred lines. *The Pharma Innovation Journal*. 2022;11(5):1642-5.
16. Ellandula A, Kumar Singh S, Narayan A, Karla U, Sagar US. Exploring genetic variability and association in QPM maize (*Zea mays*) inbred lines. *Electronic Journal of Plant Breeding*. 2023 Oct 3;14(3):1222-9.
17. Anshuman Vashistha AV, Dixit NN, Dipika D, Sharma SK, Marker S. Studies on heritability and genetic advance estimates in maize genotypes. 2013;4(2):165- 168.
18. Magar BT, Acharya S, Gyawali B, Timilsena K, Upadhyaya J, Shrestha J. Genetic variability and trait association in maize (*Zea mays* L.) varieties for growth and yield traits. *Heliyon*. 2021 Sep 1;7(9).
19. Rafiq CM, Rafique M, Hussain A, Altaf M. Studies on heritability, correlation and path analysis in maize (*Zea mays* L.). *Journal of Agricultural Research (03681157)*. 2010 Mar 1;48(1).
20. Gokulakrishnan M, Patel JM, Patel RM, Chaudhary DR, Vaghela GK. Genetic variability of quantitative characters over diverse environments in maize inbred lines. *Electronic Journal of plant breeding*. 2021;12(4):1236-43.
21. Thakur D.K, Sinha S.K, Mehta N.M and Prabha N. Studies of genetic variability, heritability and genetic advance for grain yield and its contributing traits in maize (*Zea mays* L.) hybrids. *PharmaInnov.J*.2023. 12(7),pp.754757.

22. Suresh D, Kachapur RM, Talekar SC, Gurumurthy R. Genetic variability and trait association studies in maize (*Zea mays* L.). *Journal of Farm Sciences*. 2021 Jun 29;34(02):122-5.
23. Wedwessen T, Zeleke H. Genetic variability, heritability and genetic advance estimates in maize (*Zea mays* L.) inbred lines. *International Journal of Novel Research in Life Sciences*. 2020;7(2):10-6.
24. Kinfe H, Tsehaye Y. Studies of heritability, genetic parameters, correlation and path coefficient in elite maize hybrids. *Academic research journal of agricultural science and research*. 2015;3(10):296-303.
25. Bhusal T, Lal GM, Marker S, Synrem GJ. Genetic variability and traits association in maize (*Zea mays* L.) genotypes. *Annals of Plants and Soil Research*. 2017;19(1):59-65.
26. Bekele A, Rao TN. Estimates of heritability, genetic advance and correlation study for yield and its attributes in maize (*Zea mays* L.). *Journal of Plant Sciences*. 2014;2(1):1-4.
27. Khan S, Mahmud F. Genetic variability and character association of yield components in maize (*Zea mays* L.). *American Journal of Plant Sciences*. 2021 Nov 4;12(11):1691-704.