

**Integrated Plant Nutrient Management on Bulb Quality and Nutrient Uptake of  
Aggregatum Onion (*Allium cepa* var. *aggregatum*)**

**ABSTRACT**

A field experiment was conducted in the Department of Horticulture, Pandit Jawaharlal Nehru College of Agriculture and Research Institute, Karaikal, U.T. of Puducherry, during the summer season of 2023 in aggregatum onion [*Allium cepa* var. *aggregatum*] cv 'Perambalur Local' with the objective of assessing the impact of soil fertility and integrated plant nutrient management on quality of onion bulbs and the plant response to nutrient uptake. **The experiment was a 6 x 3 factorial laid out in a randomized complete block design and replicated two times. The treatments consisted of six fertilizer types and three bioenhancers levels with two replications resulting in eighteen treatment combinations.** The study materials included vermicompost, poultry manure, panchagavya, jeevamirtham, as well as recommended farmyard manure (FYM) and N, P, K fertilizers. One of the notable findings was that the application of RDFYM + 75% N + RDP + RDK + 25% N through poultry manure coupled with the application of jeevamirtham at 500 L ha<sup>-1</sup> as soil drench thrice during irrigation (at planting, 20 and 45 DAP), resulted in the highest Total Soluble Solids (TSS) and ascorbic acid content, measuring 12.64 °Brix and 15.81 mg 100 g<sup>-1</sup> respectively. Moreover, this treatment also exhibited the highest uptake of N, P and K by onion bulbs, recording 15.27 kg ha<sup>-1</sup>, 3.38 kg ha<sup>-1</sup> and 15.27 kg ha<sup>-1</sup> respectively (F<sub>3</sub>B<sub>2</sub>). Furthermore, post-harvest soil analysis indicated that the treatment RDFYM + 50% N + RDP + RDK + 50% N through poultry manure, combined with jeevamirtham at 500 L ha<sup>-1</sup> as a soil drench thrice during irrigation (at planting, 20 and 45 DAP), showed the highest levels of available N, P and K. The recorded values were 179.60 kg ha<sup>-1</sup> of N, 12.66 kg ha<sup>-1</sup> of P and 171.58 kg ha<sup>-1</sup> of K (F<sub>5</sub>B<sub>2</sub>). **In short-duration aggregatum onions, using a combination of jeevamirtham and poultry manure boosts nutrient availability, which in turn enhances yield, nutrient quality and nutrient uptake. This approach is recommended for optimal plant health and productivity.**

**Keywords:** Aggregatum onion; Soil fertility; Integrated plant nutrients; Poultry manure; Jeevamirtham.

## 1. INTRODUCTION

“Aggregatum onion (*Allium cepa* var. *aggregatum*) is one of the important commercial bulbous vegetable crops having multivariuous uses especially in the South Indian delicacies. Though India is the second largest producer of onion in the world, there is a need for increasing as well as sustaining the productivity of this commercially important vegetable to meet the ever-growing demand. Onion is popularly designated as “Queen of the kitchen” due to its rich flavour, fragrance and unique taste, which is attributable to a volatile compound “allyl-propyl disulphide” contained in it” (Yadav *et al.*, 2015). “Onion is fairly good in its nutrient status containing carbohydrate (11.0 g), proteins (1.2 g), fiber (0.6 g), moisture (86.8 g) and several vitamins like vitamin A (0.01 mg), vitamin C (11 mg), thiamine (0.08 mg), riboflavin (0.01 mg) and niacin (0.2 mg). It also contains minerals like phosphorus (39 mg), calcium (27 mg), sodium (1.0 mg) and potassium (157 mg) in 100 g of bulb” (Sharma *et al.*, 2018). “Application of chemical fertilizer alone is reported to have resulted in increased crop yield in the initial years but adversely affected the sustainability at a later stage. The cost of chemical fertilizers is also increasing day by day. Reducing the dependence on chemical fertilizers without compromising the sustainability in production is considered as a vital issue in modern agriculture, which could be achieved only through integrated plant nutrient supply system (IPNS). Integrated nutrient management serve as the effective source of manuring in obtaining sustainable productivity without any detrimental effects on soil in an eco-friendly manner. Besides organic manures help in mitigating multiple nutrient deficiencies. Application of organic manures to acidic soil can reduce the soluble and exchangeable aluminium temporarily by forming complex and provides better environment for growth and development by improvement in physical, chemical and biological properties of soil” (Jamir *et al.*, 2013). “Enhancing productivity and sustaining the yield of crop primarily depends on nutrient availability and nutrient use efficiency. Though many factors are responsible for the existing yield gap in aggregatum onion, the major factor attributable is the poor availability and uptake

of nutrients by this short duration crop and such a trend in crop production warrants finding of suitable alternate approach and integrated nutrient management practices with the use of bio compost and bioenhancers is a recent approach towards achieving this objective. The use of manures and compost are helpful in maintaining soil health by increasing organic matter content, besides enhancing the crop productivity” (Miglani *et al.*, 2017). Keeping the above facts in view, the present investigation was conducted to study the effect of “Integrated Plant Nutrient Management on Bulb Quality and Nutrient Uptake of Aggregatum Onion (*Allium cepa* var. *aggregatum*)”.

## 2. MATERIALS AND METHODS

The experiment on Aggregatum onion (*Allium cepa* var. *aggregatum*)” was carried out in the Department of Horticulture, Pandit Jawaharlal Nehru College of Agriculture and Research Institute, Karaikal, U.T. of Puducherry, India, during the summer season of 2023. Aggregatum onion type ‘Perambalur Local’ collected from farmer’s field in Padalur village of Perambalur district of Tamil Nadu (India) was used for the study. The treatment materials for the study comprised of vermicompost, poultry manure, panchagavya and jeevamirtham, besides recommended FYM and N, P, K fertilisers. The experiment was a 6 x 3 factorial laid out in a randomized complete block design and replicated two times. The treatments consisted of six fertilizer types and three bioenhancers levels with two replications constituting eighteen treatments combinations (Table 1).

The soil of the experimental area belongs to Kottucherry series, having the textural class of sandy soil with a pH of 7.15, electrical conductivity of 0.35 dSm<sup>-1</sup>, organic carbon of 0.39 per cent, available nitrogen 140 kg ha<sup>-1</sup>, available phosphorus of 9.59 kg ha<sup>-1</sup> and potassium availability of 133.33 kg ha<sup>-1</sup>.

**Table 1. Treatment particulars**

Sl. No.	Treatment	Treatment Particulars
1.	T <sub>1</sub> - F <sub>0</sub> B <sub>0</sub>	Absolute control
2.	T <sub>2</sub> - F <sub>0</sub> B <sub>1</sub>	3 % Panchagavya foliar spray on 15, 30 and 45 <sup>th</sup> DAP
3.	T <sub>3</sub> - F <sub>0</sub> B <sub>2</sub>	Jeevamirtham 500L ha <sup>-1</sup> as soil application thrice with irrigation viz., at planting, 20 <sup>th</sup> and 45 <sup>th</sup> DAP
4.	T <sub>4</sub> - F <sub>1</sub> B <sub>0</sub>	RDF
5.	T <sub>5</sub> - F <sub>1</sub> B <sub>1</sub>	RDF + 3 % Panchagavya foliar spray on 15, 30 and 45 <sup>th</sup> DAP
6.	T <sub>6</sub> - F <sub>1</sub> B <sub>2</sub>	RDF + Jeevamirtham 500L ha <sup>-1</sup> as soil application thrice with irrigation viz., at planting, 20 <sup>th</sup> and 45 <sup>th</sup> DAP
7.	T <sub>7</sub> - F <sub>2</sub> B <sub>0</sub>	RDFYM + 75 % N + RDP + RDK + 25 % N through Vermicompost
8.	T <sub>8</sub> - F <sub>2</sub> B <sub>1</sub>	RDFYM + 75 % N + RDP + RDK + 25 % N through Vermicompost + 3% Panchagavya foliar spray on 15, 30 and 45 <sup>th</sup> DAP
9.	T <sub>9</sub> - F <sub>2</sub> B <sub>2</sub>	RDFYM + 75 % N + RDP + RDK + 25 % N through Vermicompost + Jeevamirtham 500L ha <sup>-1</sup> as soil application thrice with irrigation viz., at planting, 20 <sup>th</sup> and 45 <sup>th</sup> DAP
10.	T <sub>10</sub> - F <sub>3</sub> B <sub>0</sub>	RDFYM + 75 % N + RDP + RDK + 25 % N through Poultry manure
11.	T <sub>11</sub> - F <sub>3</sub> B <sub>1</sub>	RDFYM + 75 % N + RDP + RDK + 25 % N through Poultry manure + 3 % Panchagavya foliar spray on 15, 30 and 45 <sup>th</sup> DAP
12.	T <sub>12</sub> - F <sub>3</sub> B <sub>2</sub>	RDFYM + 75 % N + RDP + RDK + 25 % N through Poultry manure + Jeevamirtham 500L ha <sup>-1</sup> as soil application thrice with irrigation viz., at planting, 20 <sup>th</sup> and 45 <sup>th</sup> DAP
13.	T <sub>13</sub> - F <sub>4</sub> B <sub>0</sub>	RDFYM + 50 % N + RDP + RDK + 50 % N through Vermicompost
14.	T <sub>14</sub> - F <sub>4</sub> B <sub>1</sub>	RDFYM + 50 % N + RDP + RDK + 50 % N through Vermicompost + 3 % Panchagavya foliar spray on 15, 30 and 45 <sup>th</sup> DAP
15.	T <sub>15</sub> - F <sub>4</sub> B <sub>2</sub>	RDFYM + 50 % N + RDP + RDK + 50 % N through Vermicompost + Jeevamirtham 500 L ha <sup>-1</sup> as soil application thrice with irrigation viz., at planting, 20 <sup>th</sup> and 45 <sup>th</sup> DAP
16.	T <sub>16</sub> - F <sub>5</sub> B <sub>0</sub>	RDFYM + 50 % N + RDP + RDK + 50 % N through Poultry manure
17.	T <sub>17</sub> - F <sub>5</sub> B <sub>1</sub>	RDFYM + 50 % N + RDP + RDK + 50 % N through Poultry manure + 3% Panchagavya foliar spray on 15, 30 and 45 <sup>th</sup> DAP
18.	T <sub>18</sub> - F <sub>5</sub> B <sub>2</sub>	RDFYM + 50 % N + RDP + RDK + 50 % N through Poultry manure + Jeevamirtham 500L ha <sup>-1</sup> as soil application thrice with irrigation viz., at planting, 20 <sup>th</sup> and 45 <sup>th</sup> DAP

A fertilizer dose of 60:60:30 kg NPK ha<sup>-1</sup> along with 25 t ha<sup>-1</sup> of FYM (Crop Production Guide of Horticultural Crops - 2020) was taken as the reference for fertiliser application. The reduced level of N fertiliser was compensated with vermicompost in T<sub>7</sub> to T<sub>9</sub> and T<sub>13</sub> to T<sub>15</sub> and by poultry manure in T<sub>10</sub> to T<sub>12</sub> and T<sub>16</sub> to T<sub>18</sub>. The quantity of vermicompost and poultry manure applied in each plot were calculated to compensate the requirement based on the N content in vermicompost (1.50 per cent) and poultry manure (3.03 per cent). The post harvest soil and plant samples collected were analysed for nutrient content as indicated in Table 2.

The quality parameters viz., TSS and ascorbic acid content in bulbs were analysed as proposed by (Tigchelaar, 1986; Sadasivam and Balasubraminan1987).

**Table 2. Details of analytical methods employed for soil and plant analysis**

Sl. No.	Analysis	Methodology	Reference
<b>I. Soil Analysis</b>			
<b>A</b>	<b>Physico - chemical properties</b>		
1	Soil reaction (1:2.5)	Using glass electrode in the ELICO (LI 120) pH meter	Jackson (1973)
2	Electrical conductivity (1:2.5)	Using ELICO (CM 180) conductivity meter	Jackson (1973)
<b>B</b>	<b>Chemical properties</b>		
1	Organic carbon	Chromic acid wet digestion Method	Walkley and Black (1934)
2	Available nitrogen	Alkaline permanganate Method	Subbiah and Asija (1956)
3	Available phosphorous	Using 0.5 M NaHCO <sub>3</sub> of pH 8.5	Olsen <i>et al.</i> (1954)
4	Available potassium	Flame photometric method	Stanford and English (1949)
<b>II. Plant Analysis</b>			
1	Total nitrogen	Kjeldahl's method	Bremner (1965)
2	Total phosphorous	Vanadomolybdate yellow colour method using diacid Extract	Jackson (1973)
3	Total potassium	Flame photometric method using the neutralised diacid extract	Jackson (1973)

### 3. RESULTS AND DISCUSSION

#### 3.1 Effect of organics and bioenhancers on quality parameters in aggregatum onion

##### 3.1.1 Quality parameters

In view of the increased awareness of the consumers on quality aspects of vegetables, the present study with different fertilizer treatments and bioenhancers revealed the existence of significant influence of various fertilizer treatments, bioenhancers used and their interaction effect of the factors studied for total soluble solids and ascorbic acid content. The highest value for TSS (12.64 °Brix, which differed significantly from rest of the treatments) and ascorbic acid content (15.81 mg 100 g<sup>-1</sup> of bulbs) was recorded for F<sub>3</sub> B<sub>2</sub> (Table 3). This treatment involving the application of inorganic fertilizer with 25 per cent reduction in N fertilizer duly compensated with the application of poultry manure has recorded the highest total soluble solids as well as ascorbic acid content in bulbs of onion and this was found to be in conformity with the earlier findings of Kumar *et al.* (2018) and Akhil and Singh (2022) for total soluble solids and Kumar *et al.* (2018) and Vishwaraj *et al.* (2022) for ascorbic acid content. The increase in total soluble solids with the combined application of fertilisers and poultry manure is attributed to the effective translocation of sucrose from the photosynthesizing tissue to the storage tissue through source and sink relationship as reported by Kumar *et al.* (2018), besides the enhanced metabolic activities under integrated nutrient application as recorded by Dilpreet *et al.* (2016). The increase in ascorbic acid content is attributed to the better nutrient availability, improved plant metabolic function and enhanced enzyme activity as reported by Vishwaraj *et al.* (2022). The improved quality parameters *viz.*, total soluble solids and ascorbic acid observed with soil application of the bioenhancer jeevamirtham is likely to be the result of quick build up of soil fertility, resulting in increased availability and uptake of nutrients, improved beneficial biota in the rhizosphere and growth promoting substances arising out of soil application of jeevamirtham, as suggested by Chakraborty and Sarkar (2019) and Reddy and Menon (2021).

**Table 3. Effect of nutrients on total soluble solids (°Brix) and ascorbic acid content (mg 100 g<sup>-1</sup>) in aggregatum onion cv. Perambalur Local**

Treatment	Total soluble solids (°Brix)		Ascorbic acid content (mg 100 g <sup>-1</sup> )	
T <sub>1</sub> - F <sub>0</sub> B <sub>0</sub>	8.85		9.26	
T <sub>2</sub> - F <sub>0</sub> B <sub>1</sub>	9.83		10.13	
T <sub>3</sub> - F <sub>0</sub> B <sub>2</sub>	10.29		10.37	
T <sub>4</sub> - F <sub>1</sub> B <sub>0</sub>	11.22		12.84	
T <sub>5</sub> - F <sub>1</sub> B <sub>1</sub>	11.44		13.58	
T <sub>6</sub> - F <sub>1</sub> B <sub>2</sub>	11.65		13.83	
T <sub>7</sub> - F <sub>2</sub> B <sub>0</sub>	11.37		13.21	
T <sub>8</sub> - F <sub>2</sub> B <sub>1</sub>	11.88		14.69	
T <sub>9</sub> - F <sub>2</sub> B <sub>2</sub>	11.80		15.06	
T <sub>10</sub> - F <sub>3</sub> B <sub>0</sub>	11.20		13.34	
T <sub>11</sub> - F <sub>3</sub> B <sub>1</sub>	11.78		15.56	
T <sub>12</sub> - F <sub>3</sub> B <sub>2</sub>	12.64		15.81	
T <sub>13</sub> - F <sub>4</sub> B <sub>0</sub>	10.38		10.87	
T <sub>14</sub> - F <sub>4</sub> B <sub>1</sub>	10.86		11.98	
T <sub>15</sub> - F <sub>4</sub> B <sub>2</sub>	10.65		12.10	
T <sub>16</sub> - F <sub>5</sub> B <sub>0</sub>	10.65		11.61	
T <sub>17</sub> - F <sub>5</sub> B <sub>1</sub>	11.21		12.47	
T <sub>18</sub> - F <sub>5</sub> B <sub>2</sub>	11.06		12.72	
Factor	SEd	CD (p=0.05)	SEd	CD (p=0.05)
Fertilisers	0.15	0.31	0.06	0.12
Bioenhancers	0.11	0.22	0.04	0.09
Fertilisers x Bioenhancers (Interaction effect)	0.26	0.54	0.10	0.22

### 3.2 Effect of organics and bioenhancers on post harvest soil properties of aggregatum onion

The differences observed for post harvest soil pH was found significant only for bioenhancers studied whereas, post harvest soil EC assessed was found to show significant variation only among fertilizer treatments (Table 4). The marginal reduction in soil pH (F<sub>1</sub>B<sub>0</sub> - 7.28) over the initial soil pH observed with soil application of jeevamirtham might be

the result of cow's urine used as an important ingredient. Acidic nature of freshly prepared jeevamirtham has been recorded earlier by Kaur (2020). However, this is found to be in controversy to the report of Chakraborty and Sarkar (2019). The treatment receiving 50 per cent N through poultry manure was found to record higher ( $F_5B_2$  -  $0.453 \text{ dSm}^{-1}$ ) EC over the initial value and this might be the effect of soluble salts present in poultry manure and such a finding has been reported already by Singh *et al.* (1992). Significant differences among fertilizer treatments and bioenhancers tried could be observed for organic carbon content with maximum values observed in  $F_4B_2$  (0.43 per cent) and it was found to be on par with  $F_4B_1$  (0.43 per cent). This is in accordance with earlier findings of Rani and Jha (2018) and Brar *et al.* (2019), who proposed that the addition of organic matter either as solid manure or liquid manure could be responsible for a slight increase in soil organic carbon content over the initial value. Significant variation with regard to the post harvest soil N, P and K status could be observed among the fertilizer treatments, bioenhancers studied as well as their interaction in (Table 4). The post harvest available N, P and K was found to be the highest in the treatment RDFYM + 50 % N + RDP + RDK + 50 % N through Poultry manure and jeevamirtham  $500 \text{ L ha}^{-1}$  as soil drench thrice with irrigation *viz.*, at planting, 20 and 45 DAP ( $F_5 B_2$  -  $179.60 \text{ kg ha}^{-1}$  of N,  $12.66 \text{ kg ha}^{-1}$  of P and  $171.58 \text{ kg ha}^{-1}$  of K). The available nutrient content was found to be higher over the initial soil value in all the treated plots except control. The highest values with regard to all the three major nutrients were observed in plots with higher replacement of inorganic 'N' through organic 'N'. This is found to be in agreement with the findings of Meena *et al.* (2019) and could be the result of slow and sustained decomposition of organic matter in releasing nutrients to the soil solution. The conjunctive use of organic manure with inorganic fertiliser could have led to the build up of active pool of NPK as suggested by Ramesh *et al.* (2017) and enhanced beneficial microbial population from the liquid manure as proposed by Brar *et al.* (2019).

**Table 4. Effect of organics and bioenhancers on post harvest soil properties in aggregatum onion cv. Perambalur Local**

Treatment	pH		EC (dSm <sup>-1</sup> )		Organic carbon (per cent)		Nitrogen (kg ha <sup>-1</sup> )		Phosphorus (kg ha <sup>-1</sup> )		Potassium (kg ha <sup>-1</sup> )	
T <sub>1</sub> - F <sub>0</sub> B <sub>0</sub>	7.24		0.306		0.37		129.80		8.94		123.77	
T <sub>2</sub> - F <sub>0</sub> B <sub>1</sub>	7.22		0.310		0.41		134.20		9.16		127.78	
T <sub>3</sub> - F <sub>0</sub> B <sub>2</sub>	7.10		0.323		0.39		136.50		9.23		128.49	
T <sub>4</sub> - F <sub>1</sub> B <sub>0</sub>	7.28		0.324		0.37		150.45		10.30		144.00	
T <sub>5</sub> - F <sub>1</sub> B <sub>1</sub>	7.22		0.328		0.40		154.45		10.39		147.25	
T <sub>6</sub> - F <sub>1</sub> B <sub>2</sub>	7.05		0.339		0.40		158.30		11.02		151.73	
T <sub>7</sub> - F <sub>2</sub> B <sub>0</sub>	7.15		0.349		0.39		160.30		11.38		154.86	
T <sub>8</sub> - F <sub>2</sub> B <sub>1</sub>	7.13		0.358		0.42		164.31		11.40		159.65	
T <sub>9</sub> - F <sub>2</sub> B <sub>2</sub>	7.11		0.367		0.42		165.20		11.42		160.09	
T <sub>10</sub> - F <sub>3</sub> B <sub>0</sub>	7.19		0.360		0.37		165.45		11.50		159.67	
T <sub>11</sub> - F <sub>3</sub> B <sub>1</sub>	7.15		0.364		0.41		169.85		11.61		162.29	
T <sub>12</sub> - F <sub>3</sub> B <sub>2</sub>	6.93		0.380		0.41		171.10		11.72		163.71	
T <sub>13</sub> - F <sub>4</sub> B <sub>0</sub>	7.16		0.427		0.40		169.21		11.79		162.27	
T <sub>14</sub> - F <sub>4</sub> B <sub>1</sub>	7.15		0.439		0.43		173.00		12.01		166.05	
T <sub>15</sub> - F <sub>4</sub> B <sub>2</sub>	6.86		0.445		0.43		175.20		12.08		165.06	
T <sub>16</sub> - F <sub>5</sub> B <sub>0</sub>	7.08		0.434		0.37		171.01		11.34		167.20	
T <sub>17</sub> - F <sub>5</sub> B <sub>1</sub>	7.09		0.442		0.42		175.89		12.10		170.20	
T <sub>18</sub> - F <sub>5</sub> B <sub>2</sub>	6.93		0.453		0.41		179.60		12.66		171.58	
Factor	SEd	CD (p=0.05)	SEd	CD (p=0.05)	SEd	CD (p=0.05)	SEd	CD (p=0.05)	SEd	CD (p=0.05)	SEd	CD (p=0.05)
Fertilisers	0.09	NS	0.01	0.02	0.01	0.02	0.09	0.19	0.09	0.19	0.09	0.19
Bioenhancers	0.06	0.13	0.01	NS	0.01	0.01	0.06	0.13	0.06	0.13	0.06	0.13
Fertilisers x Bioenhancers (Interaction effect)	0.15	NS	0.02	NS	0.02	NS	0.15	0.33	0.15	0.33	0.15	0.33

### 3 3.3. Effect of organics and bioenhancers on nutrient uptake by onion bulbs

#### 3.3.1 Nitrogen, phosphorus and potassium uptake by bulbs (kg ha<sup>-1</sup>)

The study of nutrient uptake by onion bulbs revealed significant differences among the fertiliser treatments, bioenhancers used as well as their interaction for 'N' uptake by bulbs, while the interaction effect was found insignificant for 'P' uptake of bulbs. The differences observed with regard to 'K' uptake of onion was significant only for the various fertilizer treatments used whereas, the bioenhancers used as well the interaction effect on bulbs did not show any significant variation in (Table 5). The study of nutrient uptake by bulbs which is dependent on the concentration of nutrients becomes important as it is considered to be a function of nutrient availability in soil solution. While, all the fertilizer plots were superior to control for this trait, the maximum uptake of 15.27 kg ha<sup>-1</sup>, 3.38 kg ha<sup>-1</sup> and 15.27 kg ha<sup>-1</sup> of N, P and K respectively by onion bulbs was also observed with RDFYM + 75 % N + RDP + RDK + 25 % N through Poultry manure along with application of jeevamirtham 500 L ha<sup>-1</sup> as soil drench thrice with irrigation *viz.*, at planting, 20 and 45 DAP (F<sub>3</sub>B<sub>2</sub>). The increased nutrient uptake under poultry manure treated plots has already been reported by Mahala *et al.* (2018). Significant influence of bioenhancers on nutrient uptake of onion bulbs was observed only for 'N' and 'P' in the present study. This influence of enhanced nutrient uptake by onion bulbs with the use of solid and liquid manures might be the result of breakdown of complex nitrogenous compounds to nitrate nitrogen by the action of microorganisms present in it. Increase in P could be due to greater solubilization of native P from the soil due to action of various organic acids liberated on decomposition of organics. Increased K uptake could be ascribed to the improved soil properties due to the action of organics, leading to better penetration of roots, thereby resulting in greater uptake of K as reported earlier by Ramesh *et al.* (2017). Further, the enhanced uptake of P and K by onion bulbs might also be the result of increased availability of these to nutrients as they are also available from the added poultry manure over and above the recommended dose P and K.

**Table 5. Effect of organics and bioenhancers on nutrient uptake by bulbs (kg ha<sup>-1</sup>) in aggregatum onion cv. Perambalur Local**

Treatment	'N' uptake (kg ha <sup>-1</sup> )		'P' uptake (kg ha <sup>-1</sup> )		'K' uptake (kg ha <sup>-1</sup> )	
T <sub>1</sub> - F <sub>0</sub> B <sub>0</sub>	4.54		0.91		4.52	
T <sub>2</sub> - F <sub>0</sub> B <sub>1</sub>	4.58		0.97		4.58	
T <sub>3</sub> - F <sub>0</sub> B <sub>2</sub>	4.64		1.03		4.67	
T <sub>4</sub> - F <sub>1</sub> B <sub>0</sub>	10.96		2.32		11.25	
T <sub>5</sub> - F <sub>1</sub> B <sub>1</sub>	11.08		2.44		11.31	
T <sub>6</sub> - F <sub>1</sub> B <sub>2</sub>	11.19		2.56		11.37	
T <sub>7</sub> - F <sub>2</sub> B <sub>0</sub>	12.01		2.58		12.26	
T <sub>8</sub> - F <sub>2</sub> B <sub>1</sub>	12.14		2.65		12.32	
T <sub>9</sub> - F <sub>2</sub> B <sub>2</sub>	12.29		2.72		12.41	
T <sub>10</sub> - F <sub>3</sub> B <sub>0</sub>	14.97		3.23		14.96	
T <sub>11</sub> - F <sub>3</sub> B <sub>1</sub>	15.14		3.31		15.41	
T <sub>12</sub> - F <sub>3</sub> B <sub>2</sub>	15.27		3.38		15.27	
T <sub>13</sub> - F <sub>4</sub> B <sub>0</sub>	7.69		1.56		7.78	
T <sub>14</sub> - F <sub>4</sub> B <sub>1</sub>	7.76		1.65		7.83	
T <sub>15</sub> - F <sub>4</sub> B <sub>2</sub>	7.77		1.73		7.86	
T <sub>16</sub> - F <sub>5</sub> B <sub>0</sub>	8.69		1.85		8.88	
T <sub>17</sub> - F <sub>5</sub> B <sub>1</sub>	8.60		1.90		8.94	
T <sub>18</sub> - F <sub>5</sub> B <sub>2</sub>	8.92		1.95		9.01	
Factor	SEd	CD (p=0.05)	SEd	CD (p=0.05)	SEd	CD (p=0.05)
Fertilisers	0.02	0.03	0.02	0.03	0.18	0.37
Bioenhancers	0.01	0.02	0.01	0.02	0.13	NS
Fertilisers x Bioenhancers (Interaction effect)	0.03	0.06	0.03	NS	0.31	NS

## 5. CONCLUSION

The study emphasizes the significance of optimizing soil fertility and nutrient management techniques, particularly through the judicious use of organic amendments such as poultry manure and bioenhancers like jeevamirtham, to enhance the quality attributes and nutrient uptake efficiency of aggregatum onion crops. Such integrated approaches hold promise for improving agricultural sustainability and crop productivity in onion cultivation systems.

## 6. RECOMMENDATIONS

The enhancement in yield and quality, uptake of N, P and K observed with the application of poultry manure and jeevamirtham can be attributed to the improved soil microflora, increased nutrient availability and uptake by plants, efficient movement of nutrients to the plant's sink, and better stress tolerance. Therefore, using both solid (poultry manure) and liquid (jeevamirtham) forms of these amendments is recommended as a beneficial practice for cultivating short-duration aggregatum onions.

### Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

### REFERENCES

1. Akhil and A. Singh. 2022. Effect of organic manure and inorganic fertilizer on growth, yield and quality of garlic (*Allium sativum* L.) *The Pharma Innovation J.* **11** (9): 656-658.
2. Brar, P. S., R. Kaushal and G. Bhardwaj. 2019. A Review on Beneficial of PGPR and Noble Liquid Manures in Enhancing Soil Fertility and Sustainability. *Int. J. Curr. Microbiol. & App. Sci.*, **8** (4): 409-415.
3. Bremner, J. M. 1965. Total nitrogen. Methods of soil analysis. Chemical and microbiological properties, *Agronomy*, **9** (2): pp. 1149-1178.
4. Chakraborty, B. and I. Sarkar. 2019. Quality Analysis and Characterization of Panchagavya, Jeevumrutha and Sasyamrutha. *Int. J. Curr. Microbiol. & App. Sci.*, **8** (5): 2018-2026.
5. Crop Production Guide - Horticulture Crops. 2020. Directorate of Horticulture and Plantation Crops, T. N. Agric. Univ., Coimbatore. pp. 149-151.

6. Dilpreet, T., S. Kulbir, S. Varinder and S. Jagdish. 2016. Growth Yield and Quality of Onion as Influenced by Integrated Nutrient Management. *Int. J. Agric. Sci.*, **8** (51): 2295-2298.
7. Jackson, M. L. 1973. Soil chemical analysis. *Prentice-Hall of India Pvt. Ltd.*, New Delhi, **498**: 151-154.
8. Jamir, S., V.B. Singh, S.P. Kanaujia and A.K. Singh. 2013. Effect of integrated nutrient management on growth, yield and quality of onion (*Allium cepa* L) *Progressive Horticulture*. **45** (2): 373-380.
9. Kaur, A. 2020. JEEVAMRUTHAM: An effective activator of soil microorganisms. *Just Agriculture e-Newsletter*. **1**(1):1-5.
10. Kumar, S., D.K. Rana, K.N. Sha and V. Singh. 2018. Responses of onion (*Allium cepa* L.) cv. Pusa Red to Various organic manures under subtropical condition of Garhwal, Himalaya. *J. Pharmacogn & Phytochem.*, **7** (1): 2294-2297.
11. Mahala, P., M.R. Chaudhary and O.P. Garhwal. 2018. Yield and Quality of *rabi* Onion (*Allium cepa* L.) Influenced by Integrated Nutrient Management. *Int. J. Curr. Microbiol. & App. Sci.*, **7** (5): 3313-3321.
12. Meena, R.N., A.K. Meena and K. Singh. 2019. Yield, Quality, Economics and Nutrient uptake of Onion (*Allium cepa* L.) Influenced by Organic Nitrogen Mngement. *Int. J. Curr. Microbiol. & App. Sci.*, **8** (10): 16-23.
13. Miglani, A., N. Gandhi, N. Singh and J. Kaur. 2017. Influence of different organic manures on the growth and yield of okra. *Int. J. Advance Res. Sci. & Engg.*, **6** (1): 886-892.
14. Olsen, S. R., C. V. Cole, P. S. Watanabe and L. A. Dean. 1954. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. *USDA Circular*, p. 939.
15. Ramesh, G., K. Ajithkumar, Y.S. Amaresh and A.S. Savitha. 2017. Influences of Integrated Nutrient Management on Growth Parameters, Yield and Severity of Diseases in Onion (*Allium cepa* L.) *Int. J. Curr. Microbiol. & App. Sci.*, **6** (8): 1020-1028.

16. Rani, M. and A.K. Jha. 2018. Effect of potassium management on yield, nutrient uptake and storability of *kharif* onion (*Allium cepa* L.) and residual fertility of soil under the Alluvial Zone of Bihar. *J. Pharmacogn. & Phytochem.*, **7** (2): 377-382.
17. Reddy, A. V. V. and S. Menon. 2021. A study on role of jeevamirth in natural farming: A replacement for synthetic fertilizers. *J. Emerg. Technol. Innov. Res.*, **8**(5): a89-a93.
18. Sadasivam, S and T. Balasubraminan. 1987. In: Practical manual in Biochemistry. T. N. Agric. Univ., Coimbatore. p.14.
19. Sharma, S.K., O.P. Garhwal, P. Mahala and T.V. Yadav. 2018. Influence of integrated nutrient management on yield attributes and economics of *kharif* onion (*Allium cepa* L.) under loamy soils. *Int. J. Curr. Microbiol. & App. Sci.*, **7** (6): 2806-2811.
20. Singh, Y., B. Singh and C. S. Khind. 1992. Nutrient transformation in soils amended with green manure. *Advances in soil. sci.*, **20**: 237-309.
21. Stanford. S and L. English. 1949. Use of flame photometer in rapid soil tests for K and Ca. *Agron. J.*, **41**(9): 446-447.
22. Subbiah, B.V. and C. L. Asija. 1956. A rapid procedure for method for the estimation of available nitrogen in soils. *Current Sci.*, **25** (8): 259-260.
23. Tigchelaar, E. C. 1986. Tomato breeding. In: Bisset (Ed.) Breeding Vegetable crops. *AVIP Publishing Co.*, pp. 135-170.
24. Vishwaraj, A., M. Singh, V. Kumar, S. K. Lodhi, U. P. Shahi and K. Alam. 2022. Effect of Nutrient Management on Growth and Quality of Garlic (*Allium sativum* L.) cv. Yamuna Safed-3. *Biological Forum- An International journal*. **14** (3): 1058-1062.
25. Walkley, A. and I. A. Black. 1934. An examination of the Degtjareff Method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Sci.*, **37** (1): 29-38.
26. Yadav, R., D.H. Dwivedi, Govind and S. Maji. 2015. Effect of integrated nutrient management on growth and yield of onion (*Allium cepa* L.) cv. Pusa Madhavi. *J. crop & weed*. **11** (1): 49-53.