

## Case report

# Progressive multifocal leukoencephalitis of the immunocompromised subject: a case report from Zinder National Hospital.

### ABSTRACT

Progressive multifocal leukoencephalitis (PML) is a rare opportunistic infection of HIV-infected individuals. It is a viral infection caused by the John Cunningham (JC) virus, which occurs during the AIDS stage. An elderly patient immunosuppressed by HIV for more than 5 years presented to the HNZ SMIT with headache, vomiting, visual disturbances, speech disorders and numbness of the limbs, all of which had been evolving in a febrile context for more than seven months. Neurological examination revealed left hemiparesis, frontal syndrome, temporal syndrome and homolateral hemianopia. Cerebral computed tomography revealed multiple lesions of leukoencephalitis. PCR revealed JC virus. The patient was put on corticosteroids, rehydration and nursing care. The course was marked by a progressive deterioration in the patient's neurological condition, culminating in death. We report the clinical, biological and CT features of this progressive multifocal leukoencephalitis.

*Keywords: PML, JC Virus, HIV*

### 1. INTRODUCTION

Progressive multifocal leukoencephalopathy (PML) is a central nervous system disorder characterized by the destructive infection of oligodendrocytes by a Polyomaviridae, specifically the John Cunningham virus (JC virus), or its reactivation [1]. More rarely, BK virus or SV40 virus can be causative agents[2]. The diagnosis of PML can be confirmed by detecting the presence of the JC virus in the cerebrospinal fluid (CSF) using PCR [3]. PML typically occurs in the context of immunodeficiency due to human immunodeficiency virus (HIV) infection, hematologic malignancies, various immunosuppressive and modulatory drugs, and rheumatologic disorders [3,4]. PML is now recognized as an infection that can occur in immunocompromised individuals due to HIV. Approximately 5-10% of all patients with AIDS may develop PML [5,6]. Thus, PML is considered one of the opportunistic infections with neurological localization in severely immunocompromised HIV patients[7]. The natural progression of the disease involves progressive deterioration, leading to death within a few months if the patient remains immunodeficient. PML causes progressively developing neurological deficits[8]. Initially, there are multiple demyelinating foci, sparsely distributed in the subcortical white matter. As the disease progresses, each focus enlarges as the virus spreads from cell to cell. Typically, PML pathology includes oligodendrocyte hypertrophy and widespread destruction [15]. It is rare for PML to manifest in patients without known predisposing factors in the context of presumed immunocompetence, but it has been described in numerous case reports and series [3,9]. Progressive multifocal leukoencephalitis is one of the rare complications that can occur in immunocompromised patients. This study should enable other clinicians to consider the possibility of its occurrence in the face of presumptive clinical signs.


### 3. CASE PRESENTATION


This study examines a case of PML in an HIV-immunocompromised patient at the infectious and tropical diseases department of the National Hospital of Zinder. The study involved the analysis of the medical records of a patient hospitalized in October 2023. The methodology used to collect the data was based on questioning, followed by a physical examination and paraclinical tests.

Mr. HW, a 55-year-old known to be immunocompromised due to HIV for over 5 years, was non-adherent to his TDF+3TC+DTG treatment. He was admitted to the infectious and tropical diseases department of the National Hospital of Zinder. The patient had no history of autoimmune, renal, or hepatic disease, nor any history of alcohol consumption, liver disease, or malignant tumors. The reasons for hospitalization included fever, headaches, visual disturbances, language impairment, and left hemiparesis, all occurring in a non-traumatic context. These symptoms had been progressing over the past 7 months and had worsened 9 days prior to admission with the onset of dyspnea, severe acute headaches resistant to usual analgesics, persistent intense fever accompanied by chills and sweats, diffuse abdominal pain resembling a torsion, food vomiting, and a dry cough. Upon admission, the physical examination revealed a conscious patient with an oxygen saturation (SpO<sub>2</sub>) of 98%, a weight of 62 kg, a height of 1.80 m, and a BMI of 19.13 kg/m<sup>2</sup>. The clinical neurological examination was significant for spontaneous eye opening, groaning, and the inability to follow or obey commands. The patient exhibited spontaneous eye movements.

#### Clinical Findings:

1. Infectious syndrome: fever, tachycardia
2. Intracranial hypertension syndrome: headaches, vomiting, and visual disturbances
3. Immunodeficiency syndrome: fever, asthenia, weight loss
4. Frontal syndrome: intellectual dysfunction, impulsivity
5. Temporal syndrome: Aphasia, quadrant hemianopia, mental confusion
6. Parietal involvement: Loss of sensation
7. Occipital involvement: Homonymous hemianopia
8. Left spastic hemiparesis: predominantly affecting the face and arm given the clinical presentation and the context of immunodeficiency, a cranial computed tomography (CT) scan was performed.





Figures 1: The cranial CT images were acquired using helical scanning in parenchymal windows with axial slices, without contrast injection (a, c, e). These images revealed multiple hypodense foci located in the precentral frontal, periventricular, and right temporo-occipital regions, affecting the white matter while sparing the cortex and midline and showing no enhancement with contrast injection (b, d, f). Additionally, cortical and subcortical atrophy were noted.

These CT aspects were therefore indicative of multifocal leukoencephalopathy. A lumbar puncture was indicated and performed, revealing macroscopically clear cerebrospinal fluid with hypocellularity on cytology (03 cells/mm<sup>3</sup>). Biochemical analysis of the cerebrospinal fluid was not performed. PCR analysis of the cerebrospinal fluid identified the JC virus genome. The diagnosis of PML was definitively established based on anamnesis, clinical, biological, and CT findings. The patient subsequently received corticosteroid therapy, intravenous hydration, and nursing care. The patient's neurological condition progressively deteriorated, and he passed away two months after hospitalization in a state of coma.

### 3. DISCUSSION

This case suggests that non-adherence to antiretroviral therapy (ART) was a major aggravating factor in this patient's immunodepression, thereby facilitating the onset of PML. However, PML can also occur in other immunodeficiency contexts, such as lymphoid malignancies, post-transplantation, and immunosuppressive therapies [10,11]. It is important to note that cases of PML in immunocompetent individuals have been described by Varun Jain et al. in 2023 [3].

The diagnosis of PML was based on the characteristic appearance of lesions on cranial CT scans, in conjunction with the patient's clinical context. PML is characterized by white matter alterations, predominantly in the parieto-occipital regions, as seen in our patient, and rarely affects the gray matter[12]. In this patient, a succession of different clinical manifestations occurred, starting with muscle weakness and then difficulty in speaking. Visual problems followed, and the patient remained bedridden. Headaches, episodes of vomiting, fever and hemi-body deficit gradually set in. The clinical course of this patient's symptoms contrasts with the findings of other authors [13,14]. In HIV-infected patients, PML often presents with atypical clinical features [11]. The main differential diagnosis was HIV-associated leukoencephalopathy or HIV encephalitis [15].

The CT scan findings strongly suggested PML. This underscores the importance of cranial CT in suspected PML cases, although MRI, which was not performed, could have highlighted T1 hypointense and T2 hyperintense lesions [16]. Biologically, the diagnosis of PML in this case was based on PCR analysis of the cerebrospinal fluid, which identified the JC virus. The most crucial diagnostic criterion is the PCR amplification of the JC virus from CNS tissue and/or cerebrospinal fluid [5]. JC virus infection can also be diagnosed via biopsy, revealing inclusion bodies in the nuclei of affected oligodendrocytes. PCR testing of CSF for the JC virus has revolutionized the diagnosis in recent years [17]. However, this test can yield false negatives. Therefore, it should be repeated if clinical and imaging contexts strongly suggest the diagnosis, given its high sensitivity (up to 93%) and specificity (up to 100%). The diagnosis of PML in this patient was based on anamnesis, clinical, biological, and CT criteria [15]. Besides PML, other JC virus-related neurological conditions include JC virus granule cell neuronopathy, JC virus encephalopathy, and JC virus meningitis [18]. To date, there is no defined treatment for PML, and the prognosis is fatal in most cases [19,20].

## 4. CONCLUSION

Progressive multifocal leukoencephalopathy is a rare demyelinating brain infection affecting the central nervous system's white matter, predominantly occurring in HIV-immunocompromised patients. No specific treatment has proven effective to date, and death typically occurs within 1 to 9 months after onset.

## Disclaimer (Artificial intelligence)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

## CONSENT

All authors declare that written informed consent was obtained from the patient for the publication of this case report and accompanying images.

## ETHICAL APPROVAL

This study was conducted by ethical principles and received approval from the institutional review board (IRB) OF THE Faculty of Medical Sciences University of Zinder (FSS-UAS), NIGER. The IRB reviewed the research plan, ensuring that it adhered to ethical standards and guidelines for conducting research involving human subjects. The approval from the IRB signifies compliance with the principles of confidentiality, informed consent, and the protection of participants' rights.

## ACKNOWLEDGEMENTS

The authors thank all study participants as well as the clinic staff at the National Hospital, Zinder.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. La leucoencéphalopathie multifocale progressive (LEMP) | Société canadienne de la SP [Internet]. [cité 26 déc 2023]. Disponible sur: <https://spscanada.ca/progressive-multifocal-leukoencephalopathy-pml>
2. Cortese I, Beck ES, Al-Louzi O, Ohayon J, Andrada F, Osuorah I, et al. BK virus-specific T cells for immunotherapy of progressive multifocal leukoencephalopathy: an open-label, single-cohort pilot study. *Lancet Neurol.* 1 août 2021;20(8):639-52.
3. Jain V, Branstetter H, Savaram S, Vasquez M, Swords G, Aghili-Mehrizi S, et al. Progressive multifocal leukoencephalopathy without overt immunosuppression. *Medicine (Baltimore).* 29 sept 2023;102(39):e35265.
4. Progressive neurologic dysfunction in a psoriasis patient treated with dimethyl fumarate - Bartsch - 2015 - *Annals of Neurology* - Wiley Online Library [Internet]. [cité 24 déc 2023]. Disponible sur: <https://onlinelibrary.wiley.com/doi/abs/10.1002/ana.24471>
5. Koralnik IJ. Progressive multifocal leukoencephalopathy revisited: Has the disease outgrown its name? *Ann Neurol.* août 2006;60(2):162-73.
6. The Prognosis of Progressive Multifocal Leukoencephalopathy in HIV/AIDS Patients Undergoing Highly Active Antiretroviral Treatment: A Systematic Review - PMC [Internet]. [cité 26 juill 2024]. Disponible sur: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10575567/>

7. Berger JR. Progressive multifocal leukoencephalopathy. *Handb Clin Neurol*. 2007;85:169-83.
8. Leucoencéphalopathie multifocale progressive - StatPearls - Bibliothèque NCBI [Internet]. [cité 26 juill 2024]. Disponible sur: <https://www.ncbi.nlm.nih.gov/books/NBK560584/>
9. Berciano J. Progressive multifocal leukoencephalopathy in an immunocompetent patient. *Neurologia*. 2020;35(1):58.
10. Weissert R. Progressive multifocal leukoencephalopathy. *J Neuroimmunol*. 2011;231(1-2):73-7.
11. Progressive multifocal leukoencephalopathy presenting as a solitary gray matter lesion - PubMed [Internet]. [cité 26 déc 2023]. Disponible sur: <https://pubmed.ncbi.nlm.nih.gov/9747922/>
12. Soni N, Ora M, Mangla R, Singh R, Ellika S, Agarwal A, et al. Radiological abnormalities in progressive multifocal leukoencephalopathy: Identifying typical and atypical imaging patterns for early diagnosis and differential considerations. *Mult Scler Relat Disord*. 1 sept 2023;77:104830.
13. Fiche maladie : Leucoencéphalopathie multifocale progressive [Internet]. Radeos.org. [cité 26 juill 2024]. Disponible sur: [https://www.radeos.org/maladie/fiche-leucoencephalopathie-multifocale-progressive\\_411.html](https://www.radeos.org/maladie/fiche-leucoencephalopathie-multifocale-progressive_411.html)
14. Leucoencéphalopathie multifocale progressive (LEMP) - Troubles du cerveau, de la moelle épinière et des nerfs [Internet]. Manuels MSD pour le grand public. [cité 26 juill 2024]. Disponible sur: <https://www.msmanuals.com/fr/accueil/troubles-du-cerveau,-de-la-moelle-epiniere-et-des-nerfs/infections-cerebrales/leucoencéphalopathie-multifocale-progressive-lemp>
15. Leucoencéphalite multifocale progressive :: Infections :: ePOPI [Internet]. [cité 26 déc 2023]. Disponible sur: [https://epopi.fr/?page=fiche&id=374&cat\\_id=299&tk=66730d82444338dfb7cc51de52338866](https://epopi.fr/?page=fiche&id=374&cat_id=299&tk=66730d82444338dfb7cc51de52338866)
16. Leucoencéphalopathie multifocale progressive - Troubles neurologiques [Internet]. Édition professionnelle du Manuel MSD. [cité 26 déc 2023]. Disponible sur: <https://www.msmanuals.com/fr/professional/troubles-neurologiques/infections-cerebrales/leucoencéphalopathie-multifocale-progressive>
17. Progressive Multifocal Leukoencephalopathy in AIDS: Are There Any MR Findings Useful to Patient Management and Predictive of Patient Survival? - PMC [Internet]. [cité 26 déc 2023]. Disponible sur: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7657792/>
18. Complications neurologiques de l'infection par le virus JC : revue générale - EM consulte [Internet]. [cité 26 déc 2023]. Disponible sur: <https://www.em-consulte.com/article/1430318/complications-neurologiques-de-l-infection-par-le->
19. Ali F, Lemp F, Patel D, Schiers K. Fatal Purulent Pericarditis Due to Viridans Group Streptococcus. In: B51 **Critical care case reports: cardiovascular diseases in the ICU II** [Internet]. American Thoracic Society; 2019 [cité 26 déc 2023]. p. A3529-A3529. Disponible sur: [https://www.atsjournals.org/doi/10.1164/ajrccm-conference.2019.199.1\\_MeetingAbstracts.A3529](https://www.atsjournals.org/doi/10.1164/ajrccm-conference.2019.199.1_MeetingAbstracts.A3529)
20. Dekeyser M, De Herve MDG, Hendel-Chavez H, Labeyrie C, Adams D, Adebbs-Nasser G, et al. Leucoencéphalopathie multifocale progressive et anergie lymphocytaire réfractaire au traitement après thérapie d'inhibition du deuxième signal d'activation lymphocytaire (bêlatacept). *Néphrologie Thérapeutique*. 2017;13(5):308-9.