

Original Research Article

Estimates of the fugacity of CO₂ in the Vridichannel(Côte d'Ivoire)

Abstract

From July to November 2023 and in April 2024, surface waters were collected at three separate sampling sites within the Vridi channel, to measure physical and chemical parameters such as salinity, temperature, pH, Dissolved Inorganic Carbon (DIC), pH. The fCO₂ and Total Alkalinity were calculated from pH and DIC. These surface waters were sampling during 4 seasons to study the variability of the fugacity of CO₂ (f CO₂).

Low salinity and low temperature were observed in July -August because this period corresponded to high precipitation inducing salinity dilution and the appearance of coastal upwelling. High salinity and temperature were observed in a great dry season. In the whole, salinity and temperature have same trends of variability. Whatever the season, surface waters were oversaturated in fCO₂ with respect to atmospheric equilibrium in the Vridi channel except in dry season. On average the fCO₂ decreased from the great cold season to the hot season following marine season. Indeed, the high coastal upwelling appeared in July and disappeared in October. The variability of the channel fCO₂ followed the upwelling events along the coastal areas. Moreover, parameters such Total Alkalinity (TA), Dissolved Inorganic Carbon and pH undergone the influence the coastal upwelling as fCO₂ in the Vridi channel.

Keywords: Atlantic Ocean, fugacity of CO₂, Total Alkalinity, Vridi channel, Ébrié system, lagoon

1. Introduction

Inner estuaries and other near-shore ecosystems act as sources of CO₂ to the atmosphere due to their heterotrophic ecosystem metabolic status[1–4]. Lagoons are among the most common near-shore coastal environments occupying 13% of the World's coastline[5,6]. At the interface between terrestrial and marine environments, lagoons are subject to both continental and marine influences. Rivers, rain and ground water are the mainly continental inputs into the lagoons. These waters did not only carry large amounts of particulate material in the form of clay particles and organic detritus but also dissolved material in the form of dissolved organic matter and nutrients arising from human activity in the vicinity of the lagoons. Coastal lagoon ecosystems are directly related to many processes *i.e.* coastal lagoons are dynamic, open systems where functions are dominated and controlled by the physical and chemical environment[7].

In coastal lagoons, dissolved inorganic carbon (DIC), and particularly partial pressures of CO₂ (pCO₂) are driven by several thermodynamic and biotic factors, induced by river water inputs, tidal exchanges, mixing of water masses, photosynthesis and aerobic/anaerobic degradation of organic matter, precipitation/dissolution of calcium carbonate, benthic/pelagic couplings, and air–water exchanges [8–10]. Seasonal measurements of the carbonate system parameters have been performed for instance across salinity gradient transects or strategically located stations along lagoon channels or estuaries [11–14].

In Côte d'Ivoire,[14] found that three lagoons (Potou, Ebrié, and Grand-Lahou) were oversaturated in CO₂ during all seasons. They behaved as CO₂ sources to the atmosphere due to net ecosystem heterotrophy and inputs of riverine CO₂ rich waters. While two lagoons (Aby and Tendo) were CO₂ sinks because they were undersaturated in CO₂ with respect to the atmosphere. Indeed, there was higher biological activities in these lagoons due too much phytoplankton concentration. Yet, very few systematic scientific works have been carried out in coastal lagoons in comparison to estuaries.

In this work, we describe the variability of CO₂ system parameters such as Dissolved Inorganic Carbon (DIC), pH, Total Alkalinity (TA) and fugacity of CO₂ (fCO₂) during four seasons in Vridi

channel. This study characterizes the extent of seasonal variability in $f\text{CO}_2$ over an annual cycle, then to compare it with $f\text{CO}_2$ impacted by environmental factors in tidal waters.

2. Materials and Methods

2.1 The study area and its hydrography



Fig 1. Three sampling sites location in Vridi channel during the study period (adapted from [15])

Lagoons are the most prominent coastal ecosystems of Côte d'Ivoire covering an area of 1,200 km^2 , corresponding to one of large fractions of the surface area of lagoons in West Africa [16]

The Vridi channel is situated in southern Côte d'Ivoire (Fig. 1). It is precisely located at longitude $4^{\circ}0'50''$ W, latitude $5^{\circ}15'23''$ N. This artificial estuary is achieved by the breakthrough of the spit near Jacquerville dam when performing the autonomous harbor of Abidjan in 1933. It runs from southwest (west jetty; channel entrance) to northeast (east jetty; channel exit). The Vridi channel is 2,700 m long, with an average width of 370 m and an average depth of around 18 m. Its eastern jetty, oriented towards the lagoon environment, has an average width of 370 m, an arm's length slope of 7.5 m and an average depth of -23.6 m. Its ocean-facing jetty has an average width of 350 m, a 6.5 m shoreline and an average depth of 29.5 m. Its draught is around 16 m, with an air draught of around 15 m.

The water seasons of the Vridi channel have five seasons as follow: a hot season (HS) from February to April; a rainy season (RS) from May to July; a great cold season (GSC) from August to September; a flood season (FS) from October to November and; a short cold season (SCS) from December to January [17,18].

The rivers in Côte d'Ivoire have two different hydrological regimes [19]. The Tanoé, the Bia, the La Mé and the Agnéby rivers have an equatorial transition regime with two flooding periods in June–July and October–November. The Comoé and Bandama rivers have a mixed regime with only one flooding period in September–October [14,20].

The watershed of this estuary is invaluable because of the presence of the Atlantic Ocean. On the continent, its watershed is related to that of Ebrié system [15]. Indeed, the aquatic ecosystem is an important autonomous Harbor of Abidjan, which is the one of the largest ports in West Africa. This port is home to the great industrial area of Côte d'Ivoire [21,22].

2.2 Data

This study was carried out over the period from July to November 2023 and in April 2024. During these periods, surface waters were collected at three separate sampling sites within the Vridi channel, to ensure optimum measurement accuracy of physical and chemical parameters. These samples were collected for a period of nine hours, with the aim of estimating the variability of carbonate system. Sampling was carried out as follows: the first sample was collected on July 2023 and the last on April 2024. Each measurement is assigned a number and a date with a 30-minute interval. On average, there were 18 measurements per day. 72 water samples were taken from a maximum depth of 2 m below the surface of the channel in polyethylene bottles. Once collected, the pH, salinity and temperature of these samples were measured in situ using a WTW 3630 IDS multiparameter with a precision of ± 0.01 , ± 0.01 and ± 0.1 °C respectively. Next, the water samples were poisoned with a saturated HgCl₂ solution, to reduce the pH of the samples to below 2 in order to inhibit the potential actions of the microorganisms. The samples were then hermetically sealed and stored in a cooler containing a significant quantity of ice. Sample processing and in situ storage were carried out in compliance with ISO 5667-3. Once in the laboratory, the Dissolved Inorganic Carbon (DIC) was determined using an automatic potentiometric titrator (titrand Metrohm) with 0.1 M HCl as titrant. TA and fCO₂ were computed from pH and DIC with based on the "CO2SYS" program configured by [23]. The estimated precision of fCO₂ and TA were estimated accuracy of ± 4 μ atm and ± 4 μ mol kg⁻¹ respectively [24,25].

3. Results and discussions

3.1 Physical features

In the Vridi channel, the measured salinity and temperature ranged from 4.62 to 31.65, and from 24.9 °C to 32.7 °C respectively. On average, measured salinity and temperature in 2023 were still lower than those recorded in 2024 (table 1). It is noticed that the maximum salinity as maximum temperature was observed in April 2024 and their weak values were reported in July-August 2023 (Fig.2). Indeed, March is the end of great dry season along the coast [14,18,21,26,27]. During this period, there is no water from the rivers to dilute salinity in the Ebrié lagoon because of low precipitations on the coast and the temperature is high (> 30 °C). As salinity and temperature were collected early in April so they were impacted by a long dry season. Also, low salinity and low temperature were observed in July -August because in this period (Long Rainy Season) the precipitation is still high, salinity is diluted. Vridi channel is connected to the Atlantic Ocean then the coastal upwelling water with low temperature seems to be recorded in our sampling sites. In the whole, salinity and temperature have same trends which are different when these two parameters are measured in the seawater. In coastal or open ocean and marginal sea, salinity and temperature exhibited contrasting variations [28–30].

Our measured salinity values are significantly higher than those obtained by [31] and [32]. Indeed, these authors carried out salinity measurements in the Adjin and Potou [31] and Aby [32] lagoons. Their measurements are below 1, often close to zero despite a season. However, [14] carried out salinity measurements in four lagoons (Aby, Ebrié, Tendo, Potou and Tagba) whose values (15-20) seem close to our own. Also, they highlighted that salinity in Ebrié lagoon presented strong longitudinal gradients with the highest salinities at the vridi channel but decreasing eastward and westward due to the freshwater inputs from some rivers.

Taking in account the season, if we compare our temperature values (Table 1) with those obtained by certain authors [14,31–33], they are slightly lower. Indeed, [33] found that the seasonal variation in temperature for Potou lagoon was 25.19-33.50 °C while for a recent period it averaged slightly high (29.21 \pm 0.43 °C) according to [31]. In situ temperatures for the Aby lagoon [32] have a range of values (27.72-30.73 °C) slightly exceeding those observed in our campaigns. Similar results were provided by [14] for study carried out on 4 lagoons, as in the case of salinity. Nevertheless, [31] found an average temperature for Adjin lagoon (26.72 \pm 0.09 °C) quite close to our measurements (Table 1) following the season in April particularly.

Table 1: Mean of physical and chemical parameters in Vridi channel

date	Salinity	pH	Temperature (°C)	DIC (μ mol/kg)	TA (μ mol/kg)	fCO ₂ (μ atm)
16/07/2023	7.12 \pm 1.86	7.36 \pm 0.13	25.49 \pm 1.35	345.24 \pm 18.27	672.98 \pm 121.25	972.38 \pm 297.73
26/08/2023	6.38 \pm 1.57	7.09 \pm 0.21	25.79 \pm 0.43	349.30 \pm 5.78	482.02 \pm 147.50	1517.39 \pm 937.48

26/11/2023	15.01±1.62	7.41±0.16	27.55±0.53	494.18±244.32	496.80±242.50	562.40±329.85
06/04/2024	28.16±2.62	7.65±0.12	31.46±0.76	142.11±64.03	189.75±64.58	84.03±55.19

Mean ± 1σ: standard deviation (± 1σ)




Fig.2.Distribution of salinity (red: top panel) and temperature (blue: bottom panel) during the sampling period in Vridi channel.

3.2 Distribution of the CO₂ system parameters (pH, DIC, TA and fCO₂)

3.2.1 Distribution of pH

On the sampling sites, the measurements of pH ranged between 6.75 and 7.85 (Fig.3). The measured pH shows that in two different periods, the water in the Vridi canal is on average neutral. But in high rainy season and little dry season, pH values are slightly lower than those in November and April. This result is corroborated by [15] showing that from October 2018 to September 2019, highest pH was found in February to April period and its lowest value between October and November. Indeed, during the dry season the channel waters were fed essentially by the warm waters from the Atlantic Ocean. During the low rainy season, the warm Sudanese waters of the Comoé River settle completely in this channel.

According to the authors or the periods, the pH values obtained are slightly acidic, then neutral or slightly basic. For Potou lagoon, [33] found a range of values fairly close to acidity and basicity (5.20-7.90). The range of variation in pH remains fairly wide. Authors such as [31] and [14] provided pH values of 6.26-7.02 and 6.52-7.22 respectively in the neutral range, whose amplitudes are low.

While [32] showed that Aby lagoon was almost basic (7.12-7.87). Only the latter result differs slightly from our measurements. Again, [15] mentioned that the superficial sediments were exposed to slightly neutral to basic water due the range of measured pH during all season.




Fig. 3. Measurements of pH in Vridi channel during our study periods

3.2.2 Total Alkalinity (TA) and Dissolved Inorganic Carbon (DIC) dynamics

TA as DIC values ranged from 102 to 1014 $\mu\text{mol}/\text{kg}$, and from 69 to 1016 $\mu\text{mol}/\text{kg}$ respectively. On average TA and DIC were high in rainy season in comparison to the dry season. During the great cold period, the coastal upwelling water from the Atlantic Ocean moved to the Vridi channel. The upwelled water rich in CO_2 impacted the concentrations of TA and DIC. [14] found good relationships between TA and salinity according to the season and the sides of the Ebrié lagoon. Indeed, TA variability is due to the influence of many rivers such as Mé, Comoé and Agneby in eastern and western sides respectively. This paradox between the presence of these waters and the upwelling waters is explained by the exceptionally abundant rainfall experienced by Côte d'Ivoire during the long rainy season with on average 954 mm [34].

In addition, this channel is the transit point for many industries pollutant carried by the coastal rivers inputs from the continent particularly, and by the Atlantic Ocean [18,21,27]. This pollution is due to human activities such as the intensification of agriculture with the intensive use of chemical fertilizers in watersheds.

Using the whole dataset, we determined relationships between TA and salinity, DIC and salinity. The equations of the fit are given as follow:

$$\text{TA} = -16.66(\pm 2.35) * \text{Sal} + 696.48(\pm 39.5) \quad r^2 = 0.42 \quad (1)$$

$$\text{DIC} = -8.31(\pm 2.12) * \text{Sal} + 450.5(\pm 39.5) \quad r^2 = 0.18. \quad (2)$$

The standard error on the predicted TA and on the predicted DIC is $\pm 179 \mu\text{mol}/\text{kg}^{-1}$, $\pm 161 \mu\text{mol}/\text{kg}^{-1}$ respectively. The standard errors were higher than those (TA: $\pm 7.2 \mu\text{mol}/\text{kg}^{-1}$; DIC: $\pm 16.6 \mu\text{mol}/\text{kg}^{-1}$) found during the EGEE cruises [28] showing clear discrepancy between measured and calculated TA and DIC respectively.

In the context of our study, an examination of the all of the channel data revealed a linear regression between TA and salinity with an r^2 value of 0.42 ($p = 9.1949\text{E}-10$). This means that only 42% of the variation in TA can be explained by the changes in salinity. It is important to notice that the correlation is relatively weak despite this correlation is significant. The weak correlation underlines that certain factors (not well captured in this study) other than salinity could significantly influence TA. Indeed, the

carbon cycle and marine ecosystems are influenced by terrestrial inputs, such as river outflow and upwelling dynamics in coastal areas [35,36]. As TA, relationship between DIC and salinity was established with an r^2 value (0.18) very weak ($p=0.0002$) but it is also significant. Consequently, the classic empirical relationships could not reproduce the concentrations of DIC in coastal waters, as it does not reflect the entire coastal lagoon dynamics that impact DIC spatio-temporal variability[37].

3.2.3 Distribution of CO₂fugacity (fCO₂) in Vridi channel

In the channel, the highest fCO₂ (>3000 μatm) appeared in August 2023 and the lowest value (35 μatm) in April 2024 (Fig. 4). The peak-to-peak value is too much (~3000 μatm). This variability is related to the seasons as defined previously. Indeed, in July-August the channel undergoes the intrusion of the Atlantic Ocean. During this period, there are upwellings (equatorial and coastal) in the Atlantic Ocean which are responsible for the large CO₂ outgassing from the deep to the surface water[28,38].

As presented in table 1, on average the fCO₂ decreased from the great cold season to the hot season. In fact, the high coastal upwelling appeared in July and disappeared in October, and the minor coastal upwelling was observed between February to March [39]. During dry season, water circulation is very weak in the lagoon except in its central part under tidal forcing [40]. So, the variability of the channel fCO₂ followed the upwelling events along the coastal areas.

In 2009, [14] showed up the range of fCO₂ variability in three rivers close to the Vridi channel. The mean of fCO₂ for these three rivers (Comoé, Tanoé and Bia rivers) was higher than the atmospheric equilibrium. values of fCO₂ ranged 1925 μatm to 9595 μatm . These highest fCO₂ were due to the lowest biological activities or higher freshwater discharge. In addition, during rainy season, the Comoé freshwater discharge induces a strong circulation in the eastern part of the lagoon [40]. Thus, the precipitation recorded during the rainy periods have considerable impact on the coastal water characteristics at this location. By superimposing pH and fCO₂, it is noticed that water became slightly acidic when there was an increase in fCO₂ at the surface water. The peak-to-peak value fCO₂ is about twice of our fCO₂ gap value.

During ROAM cruises from October 2018 to December 2020 in Jacqueline (Côte d'Ivoire) [29] close to the coast, calculated fCO₂ values often exceeded 1000 μatm in the July-August period because in this period, coastal upwelling was intense. Outside this period, mean fCO₂ values ranged from 500 μatm to 600 μatm . These show that coastal areas remain a source of CO₂ for the atmosphere, like the Vridi Channel, even if fCO₂ values did not reach those of the Vridi Channel.

In the temperate coastal lagoon (France), some authors [41] found that fCO₂ values showed weak seasonal variations increasing from April 2008 (474±14 μatm) to September 2008 (515±36 μatm) before decreasing towards November 2008 (463 μatm). Indeed, they provided the impact of tide and biological activities on fCO₂ variability on a daytime or a year. Despite surface waters of the Arcachon lagoon were oversaturated in CO₂ with respect to the atmosphere fCO₂ (390 μatm), they are still lower in comparison of our results.




Fig. 4. Calculated fugacity of CO₂ (fCO₂) in Vridi channel

Also, in a tidal creek in the Duplin River salt marsh-estuary coastal ecosystem (Georgia, USA), strong seasonal and tidal/diurnal variations in fCO₂ were measured by [42] with values ranging from 500 µatm at high tide to 4,000 µatm at low tide and from 1,600 µatm at high tide to 12,000 µatm at low tide during the coldest and warmest months, respectively. Horizontal advection with the intrusion of river water into the creek at high tide can explain the lower measured values. These surface waters seemed oversaturated in CO₂ because this lagoon is still a source CO₂ all the season.

4. Conclusion

From July to November 2023 and in April 2024, surface waters were collected at three separate sampling sites within the Vridi channel, to ensure optimum measurement accuracy of physical and chemical parameters such as salinity, temperature, pH, Dissolved Inorganic Carbon (DIC), pH. The fCO₂ and Total Alkalinity were calculated from pH and DIC.

Low salinity and low temperature were observed in July -August because in this period because this period corresponded to high precipitation (salinity dilution) and the appearance of coastal upwelling (low temperature). High salinity and temperature were observed in a long dry season. In the whole, salinity and temperature have same trends of variability. Whatever the season, surface waters were oversaturated in fCO₂ with respect to atmospheric equilibrium in the Vridi channel except in dry season. On average the fCO₂ decreased from the great cold season to the hot season. Indeed, the high coastal upwelling appeared in July and disappeared in October. The variability of the channel fCO₂ followed the upwelling events along the coastal areas. Moreover, parameters such Total Alkalinity (TA), Dissolved Inorganic Carbon and pH undergo the influence of the coastal upwelling as fCO₂ in the Vridi channel. In the temperate coastal lagoon (France), fCO₂ values showed weak seasonal variations increasing from April 2008 to September 2008 before decreasing towards November 2008. Despite the fCO₂ values were oversaturated in CO₂ with respect to the atmosphere fCO₂, they are still weaker in comparison of our results. Nevertheless, some lagoon areas have high fCO₂ values, as shown by [14], with values reaching 9500 µatm often. Also, others study [42] have shown the impact of tides inducing fCO₂ values of over 10000 µatm.

During ROAM cruises in Jacqueville (Côte d'Ivoire) close to the coast, $f\text{CO}_2$ values often exceeded 1000 μatm in the July-August period because in this period, coastal upwelling was intense. Despite these high $f\text{CO}_2$ values, the study area released on average less CO_2 to the atmosphere in comparison to the Vridi channel.

This study is limited on a seasonal variability chemical and physical parameters because the measurements did not take account the impact of the tides on these parameters. In addition, it will be useful to have a time-series to better estimate the $f\text{CO}_2$ with uncertainty in the Vridi channel.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

REFERENCES

1. Kemp WM, Smith EM, Marvin-DiPasquale M, Boynton WR. Organic carbon balance and net ecosystem metabolism in Chesapeake Bay. *Mar Ecol Prog Ser.* 1997;150:229–48.
2. Gattuso J-P, Frankignoulle M, Wollast R. CARBON AND CARBONATE METABOLISM IN COASTAL AQUATIC ECOSYSTEMS. *Annu Rev Ecol Syst.* 1998;29:405–34.
3. Gazeau F, Smith SV, Gentili B, Frankignoulle M, Gattuso J-P. The European coastal zone: characterization and first assessment of ecosystem metabolism. *Estuar Coast Shelf Sci.* 2004;60:673–94.
4. Borges AV, Frankignoulle M. Distribution of surface carbon dioxide and air-sea exchange in the English Channel and adjacent areas. *J Geophys Res Oceans.* 2003;108:2000JC000571.
5. Kjerfve B. Comparative oceanography of coastal lagoons. *Estuar Var [Internet]. Elsevier; 1986.* p. 63–81.
6. Barnes RSK. Coastal lagoons [Internet]. Cambridge University Press; 1980
7. Castel J, Caumette P, Herbert R. Eutrophication gradients in coastal lagoons as exemplified by the Bassin d'Arcachon and the étang du Prévost. *Hydrobiologia.* 1996;329:ix–xxviii.
8. Cai W-J. Estuarine and Coastal Ocean Carbon Paradox: CO_2 Sinks or Sites of Terrestrial Carbon Incineration? *Annu Rev Mar Sci.* 2011;3:123–45.
9. Cotovicz Júnior LC, Knoppers BA, Brandini N, Costa Sousa SJ, Abril G. A large CO_2 sink enhanced by eutrophication in a tropical coastal embayment (Guanabara Bay, Rio de Janeiro, Brazil). 2015
10. Ribas-Ribas M, Gómez-Parra A, Forja JM. Air–sea CO_2 fluxes in the north-eastern shelf of the Gulf of Cádiz (southwest Iberian Peninsula). *Mar Chem.* 2011;123:56–66.
11. Cai W, Pomeroy LR, Moran MA, Wang Y. Oxygen and carbon dioxide mass balance for the estuarine-intertidal marsh complex of five rivers in the southeastern U.S. *Limnol Oceanogr.* 1999;44:639–49.
12. Burgos M, Ortega T, Forja J. Carbon Dioxide and Methane Dynamics in Three Coastal Systems of Cadiz Bay (SW Spain). *Estuaries Coasts.* 2018;41:1069–88.

13. Frankignoulle M, Bourge I, Canon C, Dauby P. Distribution of surface seawater partial CO₂ pressure in the English Channel and in the Southern Bight of the North Sea. *Cont Shelf Res.* 1996;16:381–95.
14. Koné YJM, Abril G, Kouadio KN, Delille B, Borges AV. Seasonal Variability of Carbon Dioxide in the Rivers and Lagoons of Ivory Coast (West Africa). *Estuaries Coasts.* 2009;32:246–60.
15. N'da Samuel YMK, Albert T. Simultaneous effects of the climate change and the recent hydromorphological modification of Vridi channel (Côte d'Ivoire) on some its hydrochemical characteristics. *ecosystems.* 2020;9:10.
16. Binet D, Le Reste L, Diouf PS. THE INFLUENCE OF RUNOFF AND FLUVIAL OUTFLOW ON THE ECOSYSTEMS AND LIVING RESOURCES OF WEST AFRICAN COASTAL WATERS. *Can J Fish Aquat Sci.* 40:1692–701.
17. Yao MK, Trokourey A. Influence de l'hydroclimat sur la dynamique saisonnière de certains éléments traces métalliques dans un estuaire marin: Cas d'étude. *J Société Ouest-Afr Chim.* 2018;45:31–41.
18. Konan YM, Albert T. Fractionation distribution and ecological risk assessment of some trace metals in artificial estuary: Vridi channel (Côte d'Ivoire). *Adv Nat Appl Sci.* 2018;12:1–6.
19. Jallow BP, Toure S, Barrow MM, Mathieu AA. Coastal zone of the Gambia and the Abidjan region in Côte d'Ivoire: Sea level rise vulnerability, response strategies, and adaptation options. *Clim Res.* 1999;12:129–36.
20. Koné YJM, Abril G, Delille B, Borges AV. Seasonal variability of methane in the rivers and lagoons of Ivory Coast (West Africa). *Biogeochemistry.* 2010;100:21–37.
21. Yao MK, Brou YS, Trokourey A, Soro MB. Metal pollution and ecological risk assessment in sediment of artificial estuary: Case of Vridi Channel, Côte d'Ivoire. *J Appl Sci Environ Manag.* 2017;21:785–92.
22. Gnagne YA, Yapo BO, Meite L, Kouame VK, Gadji AA, Mambo V, et al. Caractérisation physico-chimique et bactériologique des eaux usées brutes du réseau d'égout de la ville d'Abidjan. *Int J Biol Chem Sci.* 2015;9:1082–93.
23. Pierrot D, Lewis E, Wallace D. MS Excel Program Developed for CO₂ System Calculations Program developed for CO₂ system calculations. ORNL/CDIAC-105a.
24. Millero FJ. Thermodynamics of the carbon dioxide system in the oceans. *Geochim Cosmochim Acta.* 1995;59:661–77.
25. Frankignoulle M, Borges AV. Direct and Indirect pCO₂ Measurements in a Wide Range of pCO₂ and Salinity Values (The Scheldt Estuary). *Aquat Geochem.* 2001;7:267–73.
26. Yoroba F, Kouassi BK, Diawara A, Yapo LAM, Kouadio K, Tiemoko DT, et al. Evaluation of Rainfall and Temperature Conditions for a Perennial Crop in Tropical Wetland: A Case Study of Cocoa in Côte d'Ivoire. *Adv Meteorol.* 2019;2019:1–10.
27. Yao MK, Trokourey A. Influence of the hydroclimate on the seasonal dynamics of certain metallic trace elements in a marine estuary: Case study. *J Soc Ouest-Afr Chim Fr* 2018b. 2018;45:31–41.
28. Koffi U, Lefèvre N, Kouadio G, Boutin J. Surface CO₂ parameters and air–sea CO₂ flux distribution in the eastern equatorial Atlantic Ocean. *J Mar Syst.* 2010;82:135–44.
29. Koffi KU, Konan ES, Hassoun AER, Kouadio Y. Relationship between the carbonate system and phytoplankton community in the Gulf of Guinea-Africa. *Front Mar Sci.* 2024;11:1286338.
30. Yao KM, Marcou O, Goyet C, Guglielmi V, Touratier F, Savy J-P. Time variability of the north-western Mediterranean Sea pH over 1995–2011. *Mar Environ Res.* 2016;116:51–60.

31. Yéo KM, Komoé K, Konan ES, Goné DL. Seasonal and Spatial Variations of Phytoplankton in Relation to Physico-Chemical Parameters in Adjin Lagoon, Abidjan, Côte d'Ivoire, West Africa. *J Water Resour Prot.* 2023;15:509–25.
32. Assemian-Niango S, Kpan KGK, Aka N. Seasonal variation of some major ions in a tropical lagoon waters, Aby lagoon (Côte d'Ivoire): Relationship with temperature, salinity and pH. *Int J Biol Chem Sci.* 2020;14:3391–9.
33. Marthe YK, Lanciné GD, Bamory K, Aristide DG, Ardjouma D. Seasonal and Spatial Variations in Water Physicochemical Quality of Coastal Potou Lagoon (Côte d'Ivoire, Western Africa). *J Water Resour Prot.* 2015;7:741–8.
34. Yao C, Kacou M, Koffi ES, Dao A, Dutremble C, Guilliod M, et al. Rainfall risk over the city of Abidjan (Côte d'Ivoire): first contribution of the joint analysis of daily rainfall from a historical record and a recent network of rain gauges. *Proc IAHS.* 2024;385:259–65.
35. Cramer W, Marini K, Guiot J. Current situation and risks for the future First Mediterranean Assessment Report [Internet]. Union for the Mediterranean, Plan Bleu, UNEP/MAP, Marseille, France; 2020 p. 632.
36. Abboud-Abi Saab M, Hassoun AER. Effects of organic pollution on environmental conditions and the phytoplankton community in the central Lebanese coastal waters with special attention to toxic algae. *Reg Stud Mar Sci.* 2017;10:38–51.
37. Vance JM, Currie K, Zeldis J, Dillingham PW, Law CS. An empirical MLR for estimating surface layer DIC and a comparative assessment to other gap-filling techniques for ocean carbon time series. *Biogeosciences.* 2022;19:241–69.
38. Lefèvre N. Low CO₂ concentrations in the Gulf of Guinea during the upwelling season in 2006. *Mar Chem.* 2009;113:93–101.
39. Djakouré S, Penven P, Bourlès B, Koné V, Veitch J. Respective Roles of the Guinea Current and Local Winds on the Coastal Upwelling in the Northern Gulf of Guinea. *J Phys Oceanogr.* 2017;47:1367–87.
40. Brenon I, Mondé S, Pouvreau N, Maurin JC. Modeling hydrodynamics in the Ebrié Lagoon (Côte d'Ivoire). *J Afr Earth Sci.* 2004;39:535–40.
41. Polsenaere P, Delille B, Poirier D, Charbonnier C, Deborde J, Mouret A, et al. Seasonal, Diurnal, and Tidal Variations of Dissolved Inorganic Carbon and pCO₂ in Surface Waters of a Temperate Coastal Lagoon (Arcachon, SW France). *Estuaries Coasts.* 2023;46:128–48.
42. Wang SR, Di Iorio D, Cai W, Hopkinson CS. Inorganic carbon and oxygen dynamics in a marsh-dominated estuary. *Limnol Oceanogr.* 2018;63:47–71.