

Genetic analysis for popping traits in tropical inbred lines of popcorn (*Zea mays* var. *everta*)

**ABSTRACT**

**Aim:** Combining ability studies in popcorn to identify superior parents for commercial hybrid breeding.

**Study design:** line x tester mating fashion

**Place and Duration of study:** MRC, PJTSAU, Rajendranagar Hyderabad, between 2023-2024.

**Methodology:** Popcorn (*Zea mays* var. *everta*) is a popular snack item both in India and other countries, the genetic base of popcorn germplasm is very narrow, and they resulted in few popcorn hybrids. In this study, a set of 44 popcorn hybrids obtained by crossing 22 inbred lines with two testers in line x tester fashion were evaluated along with parents and two standard checks *i.e.* Amber popcorn and BPCH-6.

**Results:** Combining ability analysis implied significant differences among parents, crosses, parent's v/s crosses for all the traits indicating variation in the selected lines of popcorn. For grain yield, lines PC 1, PC 116, PC 6-1, PC 12-2, PC 25, PC 201-1 and tester CML 451 were identified as good general combiners, whereas crosses, PC 1 x CML 451, PC 25-1 x CML 451, PC 71-1 x CML 451 and PC 5 x CML 286 showed high desirable *sca* effects for grain yield. The popping expansion trait in crosses *viz.*, PC 22 x CML 451, PC 5 x CML 286, PC 72-2 x CML 451 displayed desirable *sca* effects. In addition to these hybrids PC 5 x CML 286 exhibited positive significant *sca* values along with high *per se* performance for both the traits.

**Conclusion:** The lines which show desirable performance for yield and popping traits will be utilized in further development of popcorn hybrids.

**Keywords:** Popcorn, *Zea mays* var. *everta*, line x tester, Combining ability, Good general combiners, Specific combining ability.

**INTRODUCTION**

popcorn is one of the most popular snack foods in larger parts of the world. Popcorn [*Zea mays* L. *sspeverta*] is a form of flint corn and differs from dent and other soft corns in two ways. The first is that it contains almost entirely hard starch. The second is that it has a very hard pericarp and outer layers of endosperm, which permit the internal pressure and temperature to rise high enough to pop (Cretors, 2001). Indian popcorn market is estimated to be worth of \$209 million by 2025 (<https://www.researchandmarkets.com>). India is the topmost popcorn importer in the world, and it

imports majorly from Argentina and USA. During 2020-21, total value of popcorn imported in India was \$5.09 Million.

The surge in popcorn consumption over the last decade, is mainly due to the proliferation of multiplexes, malls, and food courts in both major cities and Tier-II locations. Despite this growth, the market still offers a very limited variety of popcorn. Very few popcorn varieties i.e., Pearl Popcorn variety of PAU, Amber Popcorn and BPCH 6 hybrid from PJTSAU, Hyderabad, Jawahar Popcorn from JNKVV, Bajaura Popcorn from CSK, Palampur, and Pant Popcorn 1 from GBPUA & T, Pantnagar are predominantly under cultivation.

However, these existing popcorn varieties and hybrids have low popping volume and yield. Hence, there is need to develop high-yielding popcorn hybrids with increased popping volume. Popping traits are majorly governed by polygenes (Clary 1954; Ziegler 2001) and are influenced by environmental factors (Li *et al.* 2003) as well as popping techniques (Dofinger *et al.* 1990). Hence, both genetic and environmental factors should be considered for enhancing popping traits in popcorn varieties and hybrids. Genetical factors include balancing high yield with desirable popping traits, with efforts focused on developing hybrids that excel both in productivity and popping quality (Amaral *et al.*, 2016; Srdic and Pajic, 2011). Introducing traits like popping expansion represents a strategic approach in breeding high yielding pop corn hybrids with good popping expansion.

In popcorn, to exploit maximum heterotic potential for yield and quality, where in parental selection plays a key role for a sound hybrid breeding programme. In this context, line × Tester analysis by Kempthorne (1957) has widely been used for identification of promising inbred lines by crossing lines them with testers. The value of any inbred line in hybrid breeding ultimately depends on its ability to combine very well with other lines to produce superior hybrid combinations. The present study involving a Line × Tester analysis was aimed at to determine both the general combining ability (GCA) of inbreds and specific combining ability (SCA) of crosses for grain yield, its components and quality in popcorn and to identify promising inbred lines and high heterotic combinations for grain yield coupling with popping volume.

## **Material and Methods**

Experimental material consisted of 22 popcorn inbred lines collected from diverse populations. In *kharif*, 2023a total of 44 F<sub>1</sub> crosses were generated by mating 22 popcorn lines with two designated maize testers i.e., CML 286 and CML 451 following a Line × Tester mating design. These hybrids along with parents two standard checks viz., Amber popcorn and BPCH 6 were evaluated during the *rabi*, 2023-2024 in RBD in two replications at MRC, ARI, PJTSAU, Rajendranagar, Hyderabad under the specific environmental conditions of Hyderabad, characterized by an altitude of 542.6 m and coordinates approximately 79°23'E longitude and 17°19'N latitude. For each entry, row length of 4 m was maintained and row-row and plant-plant distance of 60cm and 20 cm were followed, respectively. All the recommended package of practices were followed to raise a good crop. Data were recorded on eight yield and yielding attributing traits viz., days to 50 percent tasseling, days to 50% silking, plant height (cm), number of kernel rows per ear, number of

kernels per row, starch content, popping expansion (%), and grain yield (kg/ha). Popping expansion ratio was recorded as per the guidelines of ICAR Indian Institute of Maize Research, Ludhiana(<https://iimr.icar.gov.in>). **The data analysis was conducted using R software Agricole package**, following standard procedures for analysis of variance by (Panse and Sukhatme, 1985) and Line x Tester analysis (Kempthorne, 1957).

## Results and Discussion

### Analysis of Variance

The analysis of variance revealed that there were no significant differences among the replications for all the studied eight traits. This suggests that the experiment was conducted under consistent conditions so that the performance of entry was similar in each of the two replications. Whereas, all the genotypes had shown significant differences for all the traits (Table 1), indicating sufficient amount of variability among the genotypes. The analysis of variance for parents and crosses were also significant indicating that parents were diverse and transmitted their valuable potential to their crosses for all the traits. Significant differences were also found between parents vs. crosses for all traits suggesting expression of heterosis for all the traits. These results agreed with those obtained by El-Shamarka *et al.* (2020) and Ismail *et al.* (2020).

The analysis of variance of crosses was further divided into three components *i.e.*, lines, testers, and line x tester interactions. Significant differences for lines were observed for four traits *viz.*, days to 50% tasseling, days to 50% silking, starch content, popping expansion and for testers significant differences were found for plant height, grain yield and popping expansion. Whereas for line x tester, significant differences were found for all the traits except days to 50% tasseling, days to 50% silking and starch content indicating the importance of SCA of parents in getting good hybrid combinations and the influence of dominant or non-additive gene action in the inheritance of these traits.

**Table 1. ANOVA for L x T analysis for yield, its components and quality traits in pop corn**

Source of Variation	d.f.	Days to 50% tasseling	Days to 50% silking	Plant height	Number of kernel rows per ear	Number of kernels per row	Grain yield	Starch content	Popping expansion
Replications	1	2.7	1.9	4.06	0.6	0.5	302395	0.898	0.2
Genotypes	67	6.6	6.3	555.8	9.3	51.	6450971	18.6	22.2
Parents	23	7.5	9.5	146.4	5.7	45.2	1476618	18.2	33.2
P. vs. C.	1	1.8	111.3	26258.7	16.4	1646.2	243290219	105.9	13.9
Crosses	43	1.9	2.1	177.1	11.0	17.9	3603783	16.7	16.5
Lines	21	2.7	3.2	180.4	14.1	21.4	2368046	32.4	29.7
Testers	1	1.7	0.1	1722.2	4.7	11.1	41327514	0.3	14.7
Line x tester	21	1.214	1.245	100.22	8.3	14.7	3043151	1.9	3.4
Error	67	0.810	0.871	12.550	1.300	3.509	186229	1.7	0.5

Note: \*, \*\*: significant at 0.05 and 0.01 levels of probability, respectively

### Mean performance

Mean performance of 44 crosses and two checks BPCH 6 and Amber popcorn were mentioned in the Table 2. For the trait days to 50 % tasseling and days to 50% silking, none of the crosses were found significantly earlier over the checks but, two crosses PC1 x CML 286 (DT-54 days; DS-56 days) and PC10 x CML 451 (DT-54.5 days;DS-56.5 days) were found on par to the checks BPCH 6 (DT-54.5 days; DS- 56.5 days) and Amber popcorn (DT-54.0 days; DS- 56.5 days).For plant height,crosses PC 24-1x CML 286 (131 cm), PC 116 x CML 286 (139 cm) PC 12-2x CML 286 (140.5 cm),PC 22x CML 286(140.5 cm) had significant dwarf stature against checks BPCH 6 (147.0 cm) and Amber popcorn (148.0 cm).

**Table 2. Mean performance of 44 crosses and two checks for eight traits**

	Days to 50% tasseling	Days to 50% silking	Plant height	Number of kernel rows per ear	Number of kernels per row	Grain yield	Starch content	Popping expansion
PC1xCML286	54.0	56.0	162.0	16.4	40.9	5546.5	64.50	13.7
PC1xCML451	57.5	58.5	159.5	18.2	38.1	8413.5	65.25	12.2
PC5xCML286	56.5	58.5	154.0	12.3	40.7	6686.5	62.40	14.8
PC5xCML451	57.0	59.0	165.0	14.4	36.0	4460.0	62.65	9.6
PC6-1xCML286	57.0	59.0	150.5	18.3	36.4	5840.0	69.95	12.8
PC6-1xCML451	57.2	59.0	173.5	16.4	37.3	8013.5	65.30	10.2
PC10xCML286	57.1	59.0	158.0	12.4	39.1	5253.5	66.15	12.0
PC 10x CML 451	54.5	56.5	163.5	14.0	39.6	7293.5	64.35	8.85
PC12-2xCML286	56.5	57.5	140.5	16.3	37.0	6360.0	68.80	18.15
PC12-2xCML451	57.0	59.0	163.5	12.3	41.3	8113.5	69.70	15.3
PC22xCML286	57.0	59.0	140.5	12.2	33.8	5080.0	66.10	18.55
PC22xCML451	57.0	59.0	170.5	18.0	31.9	5610.0	66.30	19.9
PC24xCML286	57.0	59.0	143.5	16.4	34.3	5253.5	65.10	10.70
PC24xCML451	57.5	59.5	154.0	14.2	37.2	5200.0	64.30	10.1
PC24-1xCML286	57.0	59.0	131.0	18.1	38.7	3353.5	69.35	12.7
PC24-1xCML451	58.0	60.0	152.5	18.4	37.2	6360.0	69.75	13.2
PC25xCML286	58.0	60.5	151.5	12.4	38.0	5747.0	61.80	19.7
PC25xCML451	57.5	59.5	154.5	12.0	40.0	7243.5	63.60	19.7
PC25-1xCML286	57.5	59.0	143.5	18.0	37.3	4646.5	69.85	14.8
PC25-1xCML451	56.5	58.5	162.0	14.0	40.3	7266.5	70.00	13.25
PC26xCML286	58.0	59.5	164.5	18.0	42.0	4966.5	68.75	16.3
PC26xCML451	57.5	59.5	172.5	16.0	37.3	6593.5	69.55	14.5
PC28xCML286	58.0	60.5	158.0	16.5	41.3	4760.0	64.60	10.0
PC28xCML451	57.5	60.0	159.5	16.0	38.0	4747.0	63.25	12.35
PC38xCML286	56.5	58.5	149.5	16.4	38.0	6627.0	61.95	13.1
PC38xCML451	56.0	58.0	151.0	12.4	35.0	2773.0	61.00	12.1
PC 71-1xCML 286	58.5	59.5	152.5	14.0	39.1	4740.0	67.85	17.3
PC 71-1xCML 451	58.0	61.0	154.0	18.2	41.8	7793.5	66.80	15.75
PC72-2xCML286	58.0	61.0	151.5	12.0	34.0	4800.0	65.15	12.85
PC72-2xCML451	57.5	59.5	152.5	14.1	37.6	6893.5	65.45	14.6
PC116xCML286	57.5	59.5	139.0	14.0	42.2	6073.5	68.65	15.7
PC116xCML451	57.0	59.0	159.0	18.4	38.7	7560.0	68.40	13.2
PC117AxCML286	56.5	58.5	161.5	16.4	33.3	5213.5	63.55	11.9
PC117AxCML451	55.5	57.5	161.0	18.4	38.3	6853.0	62.20	12.7
PC200xCML286	58.0	59.5	162.0	12.2	38.7	4300.0	70.05	12.25
PC200xCML451	58.5	60.5	174.0	16.6	38.3	6840.0	70.65	11.45

PC201-1xCML286	58.5	60.5	158.0	12.4	37.3	5880.0	64.65	12.6
PC201-1xCML451	58.0	60.0	162.5	12.5	39.6	6800.0	65.80	13.8
PC209xCML286	57.5	59.5	157.0	16.3	35.4	5380.0	68.70	12.1
PC209xCML451	58.0	60.0	153.5	14.5	36.9	6446.5	69.65	12.4
PC213xCML286	57.5	59.5	157.0	12.1	32.1	4526.5	68.95	15.3
PC213xCML451	58.5	60.5	151.5	14.5	33.2	6073.5	69.95	15.6
PC506xCML286	56.5	58.5	142.0	12.0	28.3	2720.0	69.75	9.5
PC506xCML451	56.5	58.5	149.5	12.0	40.0	6560.0	70.15	8.5
Checks								
BPCH 6	54.5	56.5	147	14.2	36.8	5019.5	65.5	23.2
Amber popcorn	54.0	56.5	148.0	14.0	34.9	4894.0	66.5	24.6
C.D. 5%	1.8863	1.9046	7.1362	2.2629	3.7628	937.89	2.6431	1.6086

For the trait number of kernel rows per ear, eleven crosses out of forty-four were significantly superior over both the checks and highest was observed in three crosses viz., PC 24-1 x CML 451, PC 116 x CML 451 and PC 117A x CML-451 with a value of 18.4. Seventeen crosses were found significantly superior over both the checks for number of kernels per row and the cross PC 116 x CML 286 (42.2) followed by PC 71-1 x CML 451 (41) showed high number of kernels per row. For grain yield, among the 44 hybrids, twenty-three hybrids had significantly superior grain yield over both the checks and three hybrids viz., PC1 x CML451, PC12-2 x CML451 and PC6-1 x CML451 gave high grain yields of 8414, 8114 and 8014 kg/ha, respectively. Similar results were reported by (Aslam *et al.* 2017, El-Hosary *et al.* 2018, Abdel-Moneam *et al.* 2020, Zandrato *et al.* 2024, Mishra *et al.* 2024) Two hybrids i.e., PC 200 x CML 451 and PC 506 x CML 451 reported high starch content of 70% and the findings align with Zulkadiret *et al.* (2020). Mean value for the trait Popping expansion (%) was 14 % within the range of 8.5 % to 19.9 % these results are consistent with Zulkadiret *et al.* (2020). None of the hybrids had significantly superior popping expansion over both the checks.

**Table 3. GCA effects of lines and testers for yield, its components and quality traits in popcorn.**

	Days to 50% tasseling	Days to 50% silking	Plant height	Number of kernel rows per ear	Number of kernels per row	Grain yield	Starch content	Popping expansion
PC 1	-1.909**	-1.920**	5.094**	2.313**	1.944*	1124.057**	-1.736*	-0.538
PC 5	-0.409	-0.420	3.844*	-1.711**	0.837	-282.693	-4.099**	-1.362**
PC 6-1	-0.159	-0.170	6.469**	2.351**	-0.656	1070.807**	1.008	-2.051**
PC 10	-1.409**	-1.420**	5.219**	-1.779**	1.794	417.557	-1.344*	-3.113**
PC 12-2	-0.409	-0.920	-3.531*	-0.720	1.594	1380.807**	2.641**	3.187**
PC 22	-0.159	-0.170	-0.031	0.089	-4.681**	-510.943	-0.424	5.662**
PC 24	0.091	0.080	-6.781**	0.325	-1.781	-629.193**	-1.914*	-3.163**
PC 24-1	0.341	0.330	-13.906**	3.258**	0.452	-999.193**	2.962**	-0.545
PC 25	0.591	0.830	-2.531	-2.797**	1.469	639.307*	-3.910**	6.137**
PC 25-1	-0.159	-0.420	-2.906	0.978	1.269	100.557	3.314*	0.483
PC 26	0.591	0.330	12.844**	1.960**	2.144	-75.943	2.546**	1.649**
PC 28	0.591	1.080	3.094	1.212	2.094	-1102.443**	-2.685*	-2.371**
PC 38	-0.909	-0.920	-5.281	-0.597	-1.031	-1155.943**	-5.135*	-0.945
PC 71-1	0.841	1.080	-2.406	1.101	2.944**	410.807	0.721	2.974**
PC 72-2	0.591	1.080	-3.656	-1.978**	-1.756	-9.193	-1.322	0.179
PC 116	0.091	0.080	-6.731**	1.153	2.919*	960.807**	1.945**	0.949

PC 117A	-1.159	-1.170	5.719	2.418	-1.756	177.307	-3.707	-1.238
PC 200	1.091	0.830	12.344	-0.625	0.969	-285.943	3.740	-1.701
PC 201-1	1.091	1.080	4.594	-2.570	0.894	484.057	-1.387	-0.326
PC 209	0.591	0.580	-0.281	0.377	-1.356	57.307	2.591	-1.275
PC 213	0.841	0.830	-1.406	-1.741	-4.906	-555.943	2.847	1.962
PC 506	-0.659	-0.670	-9.781	-3.015	-3.406	-1215.943	3.349	-4.551
S.E.± (lines)	0.450	0.466	1.771	0.936	0.936	215.77	0.656	0.367
CML286	0.000	-0.034	-4.424	-0.233	-0.356	-685.307	0.06	0.409
CML451	0.000	0.034	4.424	0.233	0.356	685.307	-0.06	-0.409
S.E.±(testers)	0.135	0.140	0.534	0.282	0.282	65.05	0.198	0.110

, \*\* significant at 0.05 and 0.01 levels of probability, respectively

#### General combining ability effects:

The efficacy of parental line selection in crop improvement can be significantly improved by estimating the combining ability and SCA effects. The GCA effects represents the additive nature of gene action. Estimates of GCA effects of 22 inbred lines and two testers of eight traits are presented in Table 3. Inbred PC 1 was found to be good general combiner for days to 50% tasseling, days to 50% silking, number of kernel rows per ear, number of kernels per row, grain yield and PC 116 for dwarf plant height, number of kernel rows per ear, number of kernels per row, grain yield, starch content and popping expansion. Three inbreds viz., PC 1, PC 10 and PC 117A had significant and negative *gca* effects for flowering traits indicating potential of these inbreds in developing early maturing hybrids. Seven lines and one tester CML 286 were found to be good general combiners for plant height in desirable direction. For grain yield, in addition to PC 1 and PC 116, lines PC 6-1, PC 12-2, PC 25, PC 201-1 and tester CML 451 were good general combiners with significant *gca* effects and these inbreds can be utilized in the development of high yielding hybrids. These findings were in concurrent with work done by Lima *et al.* (2019), Parsons *et al.* (2020), Kamphorst *et al.* (2021), Omy *et al.* (2023). For starch content and popping expansion, PC 12-2, PC 26, PC 116, PC 213 were the good general combiners and could be used in breeding of better popping quality hybrids.

#### Specific combining ability effects:

The SCA effect is important criteria to evaluate the potential of hybrids. SCA effects of 44 test crosses for eight studied traits are presented in Table 4. The results revealed that none of the crosses had significant *sca* effects in desirable direction for all the traits. For grain yield, nine crosses had positive and significant *sca* effects and these crosses involved either high x high or high x low or low x low general combiners and the highest effect was found in PC 38 x CML 286 (2612.30) followed by PC 5 x CML 286 (1798.55) and PC 506 x CML 451 (1234.693). Few crosses viz., PC 1 x CML 451, PC 25-1 x CML 451 and PC 71-1 x CML 451 showing high desirable specific combining ability effects also shown high *per se* performance for grain yield and these crosses had good general combiner as one of the parents. Earlier studies of Parsons *et al.* (2020), Kamphorst *et al.* (2021), Omy *et al.* (2023) had the similar results. For days 50% tasseling, PC 1 x CML 286 had negative and significant effects and for days to silking, none of the crosses were

found significant in desirable direction. Eight, seven and three crosses had significant sca effects in desirable direction for plant height, number of kernel rows per ear and number of kernels per row, respectively. For starch content, PC 6-1 x CML 286 and for popping expansion, PC 5 x CML 286, PC 10 x CML 286, PC 22 x CML 451, PC 28 x CML 451 and PC 72-2 x CML 451 were good specific combiners. It was observed that for majority of the traits, involvement of one good general combiner is desirable to get the better specific combination. In High x low general combiners, gene interaction between positive alleles of high combiners and negative alleles of low combiners could produce good segregants. Gene action is non fixable for such traits and hence could be exploited in heterosis breeding. Similar finding of non-additive gene action in governing in governing these traits were reported by Aminu *et al.* (2014), Abdulazeez, S.D *et al.* (2021). In the present study, majority of hybrids had low to medium popping expansion. So, the best strategy to improve the popping expansion is through intra population approach until it reaches commercial expansion values and then exploit heterosis for grain yield by developing inbred lines and hybrids.

**Table 4. SCA effects of superior crosses for all the traits studied**

	Days to 50% tasseling	Days to 50% silking	Plant height	Number of kernel rows per ear	Number of kernels per row	Grain yield	Starch content	Popping expansion
PC1xCML286	-1.750**	-1.216	5.549	-0.686	1.756	-748.193	-0.438	0.341
PC 1xCML 451	1.750**	1.216	-5.549	0.686	-1.756	748.193	0.438	-0.341
PC 5xCML 286	-0.250	-0.216	-1.201	-0.812	2.713	1798.55**	-0.174	2.189**
PC6-1x CML 286	0.00	0.034	-7.076	1.213	-0.094	-401.443	2.248	0.903
PC 10x CML 286	1.250	1.284	1.674	-0.571	0.106	-334.693	0.860	1.166
PC 12-2x CML 286	-0.250	-0.716	-7.076*	2.243*	-1.794	-191.443	-0.503	1.016
PC 12-2x CML 451	0.250	0.716	7.076**	-2.243**	1.794	191.443	0.503	-1.016
PC 22x CML 286	0.00	0.034	-10.576**	-2.689**	1.281	420.307	-0.150	-1.084*
PC 22x CML 451	0.00	-0.034	10.576**	2.689*	-1.281	-420.307	0.150	1.084*
PC 24x CML 286	-0.250	-0.216	-0.826	1.333	-1.119	712.057	0.352	-0.109
PC 24-1x CML 286	-0.500	-0.466	-6.451	0.111	1.098	-817.943	-0.235	-0.653
PC 24-1x CML 451	0.500	0.466	6.451	-0.111	-1.098	817.943	0.235	0.653
PC 25-1x CML 286	0.500	0.284	-4.951	2.250**	-1.119	-624.693	-0.132	0.344
PC 25-1x CML 451	-0.500	-0.284	4.951	-2.250**	1.119	624.693	0.132	-0.344
PC 26x CML 286	0.250	0.034	0.299	1.233	2.706	-128.193	-0.464	0.753
PC 28x CML 286	0.250	0.284	3.549	0.464	2.006	691.807	0.608	-1.602**
PC 28x CML 451	-0.250	-0.284	-3.549	-0.464	-2.006	-691.807	-0.608	1.602**
PC 38x CML 286	0.250	0.284	3.674	2.225**	1.831	2612.30**	0.421	0.097
PC 71-1x CML 451	0.00	0.716	-3.549	1.881	0.994	841.443**	-0.472	-0.378

PC 72-2x CML 451	-0.250	-0.784	-3.799	0.829	1.444	361.443	0.184	1.302
PC 116x CML 286	0.250	0.284	-5.776	-1.960	2.131	-57.943	0.063	0.828
PC 116x CML 451	-0.250	-0.284	5.776	1.960	-2.131	57.943	-0.063	-0.828
PC 200x CML 286	-0.250	-0.466	-1.701	-1.952	0.581	-584.693	-0.366	-0.003
PC 200x CML 451	0.250	0.466	1.701	1.952	-0.581	584.693	0.366	0.003
PC 209x CML 451	0.250	0.216	-6.174	-1.166	0.394	-152.057	0.537	0.548
PC 213x CML 451	0.500	0.466	-7.049	0.968	0.194	88.193	0.552	0.559
PC 506x CML 451	0.00	-0.034	-0.674	-0.258	5.494	1234.693	0.274	-0.078
S.E.± (Crosses)	0.6367	0.660	2.505	1.324	1.324	305.14	0.782	0.520

*\*, \*\* significant at 0.05 and 0.01 levels of probability, respectively*

#### **Conclusion:**

**Significant differences were observed among parents and crosses across all traits, indicating ample variation among the parental lines for enhancing both yield and quality traits. Several lines (PC 1, PC 116, PC 6-1, PC 12-2, PC 25, PC 201-1) and tester CML 451 demonstrated strong potential as general combiners for grain yield, displaying positive and statistically significant general combining ability (GCA) effects. Additionally, specific crosses such as PC 1 x CML 451, PC 25-1 x CML 451, PC 71-1 x CML 451, and PC 5 x CML 286 exhibited desirable specific combining ability (SCA) effects, particularly for grain yield. For popping expansion, crosses like PC 22 x CML 451, PC 5 x CML 286, and PC 72-2 x CML 451 showed favourable SCA effects. Notably, the hybrid PC 5 x CML 286 demonstrated positive and significant SCA values coupled with strong per se performance for both traits, suggesting its potential for advancing popcorn hybrid breeding programs aimed at improving grain yield and popping expansion simultaneously.**

#### **Disclaimer (Artificial intelligence)**

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.

3.

### References:

Cretors C. Popcorn products. *Snack foods processing*. 2001; 385-420.

<https://www.researchandmarkets.com>

Clary G. A study of the inheritance of expansion in popcorn (Doctoral dissertation, Purdue University). 1954.

Ziegler K. E. Popcorn. In: Hallauer AR (ed.), *Specialty Corns*. CRC Press, Boca Raton. 2001; 199-224.

Li Y.L, Wu, S.W, Niu, S.Z. and Li, Z.Q. Analysis of combining ability of popcorn with normal corn inbreds belonging to different heterotic groups. *J Henan Agric Univ*. 2003; 37:1-5.

Dofing S.M, D'Croz-Mason N, Thomas-Compton M.A. Inheritance of expansion volume and yield in two popcorn  $\times$  dent corn crosses. *Crop Science*. 1991; 31(3):715-718.

Amaral Junior A.T.D, Freitas I.L.D.J, Guimarães, A.G, Maldonado C, Arriagada O, Mora F. Bayesian analysis of quantitative traits in popcorn (*Zea mays L.*) through four cycles of recurrent selection. *Plant Production Science*. 2016; 19(4):574-578.

Srdić J, Pajić Z. Variability of yield and kernel quality parameters of popcorn hybrids (*Zea mays L.* everta). *Journal on Processing and Energy in Agriculture*. 2011; 15(2):84-86.

Kempthorne O. *An Introduction to Genetic Statistics*. John Wiley and Sons, Inc. New York. 1957; 458-471.

<https://iimr.icar.gov.in>

**Lokonon, B.E, Guedezoomè, P.M.B, Essetcheou, S.M.M and Kakaï, R.G.. On the use and interpretation of Generalized Linear Models (GLMs) in agricultural sciences: critical analysis and application guidelines. *African Journal of Applied Statistics*. 2023; 10(1):1435-1450.**

Panse V.G, Sukhatme P.V. *Statistical methods for agricultural research*. ICAR, New Delhi. 1985; 8:308-318.

El-Shamarka S.H, Darwish I.H, Marwa M, Afifi A.A, AM N.E.D. Combining Abilities for Yield and Its Components through Line by Tester Analysis in Maize under Two Plant Densities. 2020.

Ismail M.R, El-Hosary A.A, El-Badawy M.E.M. Genetic analysis of some topcrosses of yellow maize. *Electronic Journal of Plant Breeding*. 2020; 11(4):1099-1104.

Aslam M, Sohail Q, Maqbool M.A, Ahmad S, Shahzad R. Combining ability analysis for yield traits in diallel crosses of maize. *JAPS: Journal of Animal & Plant Sciences*. 2017; 27(1).

El-Hosary A.A.A, Motawea M.H, Elgammaal A.A. Combining ability for yield and some of its attributes in maize across two locations. *Egypt. J. Plant Breed.* 2018; 22(3):625-640.

Abdel-Moneam M.A, Sultan M.S, Abido W.A, Hadrázy Á, Sadek S.A, Shalof, M.S. Investigation of combining ability and superiority percentages for yield and some related traits in yellow maize using linex tester analysis. *Acta Agraria Debreceniensis.* 2020 ;(1):5-14.

**Zendrato, Y.M, Suwarno, W.B. and Marwiyah, S. June. Combining ability and heterosis of tropical maize under acidic soil conditions. In *IOP Conference Series: Earth and Environmental Science*.2024. (Vol. 1359, No. 1, p. 012041). IOP Publishing.**

**Mishra D.K, Singh B, Chakraborty, M, Verma N, Surin S.S and Mishra,S.K. 2024. Combining Ability and Heterosis in Maize (*Zea mays* L.) Inbred Lines. *Journal of Scientific Research and Reports: 30(6) ;943-954.***

Zulkadir G, Idikut L. Determination of popping traits and grain quality of landraces popcorn populations. *Journal of Food Science and Technology.* 2021; 58:1302-1312.

Lima V.J.D, Amaral Júnior A.T.D, Kamphorst S.H, Bispo R.B, Leite J.T, Santos T.D.O, Schmitt K.F.M, Chaves M.M, Oliveira U.A.D, Santos P.H.A.D, Gonçalves G.M.B. Combined dominance and additive gene effects in trait inheritance of drought-stressed and full irrigated popcorn. *Agronomy.* 2019; 9(12):782.

Kamphorst S.H, do Amaral Junior A.T, De Lima V.J, Carena M.J, Azeredo V.C, Mafra G.S, Santos P.H.A.D, Leite J.T, Schmitt K.F.M, dos Santos Junior D.R, Bispo R.B. Driving sustainable popcorn breeding for drought tolerance in Brazil. *Frontiers in Plant Science.* 2021; 12:732285.

Omy S, Hossain M, Karim A, Amiruzzaman M, Akhi A. Combining ability assessment of popcorn considering yield and kernel popping quality.

Parsons L, Ren Y, Yobi A, Hurst P, Angelovici R, Rodriguez O, Holding D.R. Production and selection of quality protein popcorn hybrids using a novel ranking system and combining ability estimates. *Frontiers in Plant Science.* 2020; 11:539857.

Aminu D, Dawud M.A, Modu A. Combining ability and heterosis for different agronomic traits in maize (*Zea mays* L.) under drought stress in the Sudan Savanna of Borno State, Nigeria. *Plant Breeding Crop Sci.* 2014; 6(10):128-134.

Abdulazez S. D, Kakarash S. A, Ismael N.B. May. Estimation of heterosis and combining ability for yield, yield component using linex tester methods in maize (*Zea Mays* L.). In *IOP Conference Series: Earth and Environmental Science.* 2021; (Vol. 761, No. 1: 012083). IOP Publishing.

UNDER PEER REVIEW