

## **Assessment of heavy metals and physico-chemical parameters of sewage water, soil in different blocks jaipur district of rajasthanindia**

### **ABSTRACT**

Jaipur is the capital state of Rajasthan. During the year 2021, District Jaipur of 3 blocks (Chomu, Sanganer, and Shahpura) in 9 villages [Keshav Nagar (V<sub>1</sub>), Morija(V<sub>2</sub>), Nindola(V<sub>3</sub>), Goner (V<sub>4</sub>), Shrikishanpura (V<sub>5</sub>), Durgapura (V<sub>6</sub>), Shivpuri (V<sub>7</sub>), Manoharpur (V<sub>8</sub>), Nwalpura (V<sub>9</sub>)] were selected for estimation of physico-chemical properties and Heavy metal contamination in sewerage channels. A total of 9 samples were collected from 9 villages sewerage channels. This study aimed to analyze the physico-chemical properties of agricultural soil and irrigation water in three blocks of Jaipur, Rajasthan. The findings revealed significant variations in the physico-chemical properties of the agricultural soil and water in Chomu, Sanganer, and Shahpura. Standard methods were employed to assess the physico-chemical characteristics of soil samples contaminated with sewerage effluents in these regions, which exhibited considerable diversity. The pH levels of the soil samples were predominantly slightly acidic and alkaline ranging from 6.22 to 7.31. Electrical conductivity values varied between 0.41 to 0.62mmhos/cm. The percentages of organic matter and organic carbon in the soil samples ranged from 0.17% to 0.25% and 0.29% to 0.43%, respectively. Additionally, the analysis included heavy metals such as zinc (Zn), manganese (Mn), iron (Fe), and copper (Cu). The soil samples were found to contain elevated levels of heavy metals, surpassing the permissible limits. The discharge of sewerage effluent into the environment, primarily originating from the textile industry, house, etc. constitutes a major source of pollution that can adversely impact the local flora and fauna. Therefore, it is imperative to implement effluent treatment measures before their release into the environment.

**Keywords:** Jaipur district, sampling sites, physico-chemical properties, heavy metals, etc.

### **INTRODUCTION**

Environmental contamination is a sewage ecological crisis that poses significant challenges to our planet today. Developing countries in South Asia, such as Pakistan, Nepal, Bangladesh, and India, particularly experience the detrimental effects of pollution resulting from urbanization, increased household and vehicular fuel usage, and industrialization (Karn and Harada, 2001). The city of Jaipur in Rajasthan, situated in the central region, has witnessed rapid urban growth and industrial development in the past two decades. Numerous industrial areas have emerged throughout Jaipur and its neighboring regions during this period. This industrialization has led to a prevalent issue of heavy metal pollution, referring to the presence of toxic metallic elements with high density and low concentration tolerance, including copper, manganese, lead, cadmium, iron, mercury, zinc, and nickel (Raikwar *et al.*, 2008). While heavy metals occur naturally in the Earth's crust, the majority of environmental contamination and exposure stem from human activities, such as mining, smelting operations, and industrial processes encompassing metal refinement, coal burning, petroleum combustion, nuclear power generation, and manufacturing in various sectors like plastics,

textiles, microelectronics, wood preservation, and paper processing. Birds are particularly sensitive to environmental contaminants resulting from human activities and serve as vital indicators of ecological conditions in specific areas. Therefore, they are widely used in monitoring environmental changes (Medona *et al.*, 2015). The analysis of heavy metal excretion through bird fecal matter has gained attention due to the Wildlife Protection Act of 1972, which prohibits the capturing and killing of birds in India. Consequently, researchers in this field focus on analytical studies that utilize bird tissues and organs less frequently. Moreover, limited studies on heavy metal pollution in bird feces exist within the Indian context. Additionally, heavy metals possess the ability to bioaccumulate and biomagnify within food chains (Zhuang *et al.*, 2009). As a result, fecal matter serves as a valuable indicator of metal contamination, effectively reflecting the level of metal pollution in the environment. Therefore, the examination of bird feces provides an estimation of the adverse impact of trace metals on these organisms and their surrounding environment. This study aims to assess the concentration of heavy metals in bird fecal matter within industrial environments and explore the potential use of fecal matter as a non-invasive method for determining metal concentrations in the environment. Metals occur naturally in our environment, especially in the Earth's crusts where they contribute to the balance of the planet. However, as a result of human activities they are distributed, concentrated and chemically modified, which may increase their toxicity (Mihaly *et al.*, 2005).

## **MATERIALS AND METHODS**

Jaipur, the capital city and the largest city in the state of Rajasthan, is located at a geographical position of approximately 26.9124° N latitude and 75.7873° E longitude. The district of Jaipur covers a total geographical area of 11,06,148 hectares or 11,061.48 square kilometers (Government of Rajasthan, n.d.). In terms of agricultural land use, the Gross Cropped Area in Jaipur district is estimated to be around 8,48,313 hectares. Out of this, the Net Sown Area is approximately 6,63,167 hectares. However, it's important to note that only 3,02,428 hectares of the Net Sown Area are under irrigation. These figures provide an overview of the agricultural landscape in Jaipur district, highlighting the extent of cultivated land and the proportion that benefits from irrigation. These statistics are essential for understanding the agricultural potential and the distribution of agricultural activities in the region. (District Fact Book., 2019)

Nine samples were obtained from three different blocks in the Jaipur district prior to the monsoon season. These samples included soil and sewage water effluent. The purpose was to assess the physico-chemical characteristics of the collected samples. The collection took place in the morning of June 2021, with temperature and pH measurements recorded in the field. To maintain sample integrity, acid-washed plastic bottles and sterilized plastic bags were used for collection, and the samples were stored at 4°C. The soil samples underwent analysis to determine various parameters such as pH, electrical conductivity (EC), percentage of organic carbon (OC), percentage organic matter (OM), and Heavy metals zinc (Zn), iron (Fe), copper (Cu), manganese (Mn). On the other hand, the effluent samples were analyzed

for pH, electrical conductivity (EC), as well as cations and anions present. In summary, a total of nine samples consisting of soil and sewage water effluent were collected from different blocks in Jaipur district before the onset of the monsoon season. These samples were analyzed to assess their physico-chemical properties. The collection process involved maintaining cleanliness and sample preservation, and subsequent laboratory analysis focused on various parameters specific to each sample types. These heavy metals and organic compounds affect quality of soil and ground water of the area (Bhattacharjee et al., 2003). Heavy metals enter in the human body by different pathways and causes harmful effects (Gitimoniet al., 2009). Environmental pollution and continuous exposure of human beings to toxic heavy metals such as Hg, Cd, and Pb is serious growing problem throughout the world (Yusuf and Sonibare 2004). The properties of dry soil along with its type have a great importance in agriculture (Ahire et al., 2013). Soil is not only important for agriculture but also have more useful for living organism. The differences in soil characteristics associated with landscape position are usually attributed to differences in the runoff, erosion and deposition processes that affect soil genesis (Dengđz, 2010).

### Sample collection Areas

table-1 shows, were took on different locations seweragechannel water and soil samples

S.No.	Blocks	Villages	Latitude( <sup>0</sup> N)	Longitude( <sup>0</sup> E)
1.	Chomu(B <sub>1</sub> )	KeshavNagar (V <sub>1</sub> )	26.9039 <sup>0</sup>	75.7844 <sup>0</sup>
		Moriya(V <sub>2</sub> )	27.2068 <sup>0</sup>	75.7582 <sup>0</sup>
		Nindola(V <sub>3</sub> )	27.3185 <sup>0</sup>	75.7081 <sup>0</sup>
2.	Sanganer(B <sub>2</sub> )	Goner(V <sub>4</sub> )	26.8865 <sup>0</sup>	75.8341 <sup>0</sup>
		Shrikishanpura(V <sub>5</sub> )	26.7998 <sup>0</sup>	75.8582 <sup>0</sup>
		Durgapura(V <sub>6</sub> )	26.8518 <sup>0</sup>	75.7862 <sup>0</sup>
3.	Shahpura(B <sub>3</sub> )	Shivpuri (V <sub>7</sub> )	26.9426 <sup>0</sup>	75.7526 <sup>0</sup>
		Manoharpur(V <sub>8</sub> )	26.2994 <sup>0</sup>	75.9571 <sup>0</sup>
		Nwalpura(V <sub>9</sub> )	26.8103 <sup>0</sup>	75.8365 <sup>0</sup>

## RESULTS AND DISCUSSION

The soil samples adjoining the sewage water effluent, agriculture land regions of Jaipur. They also show different variation in the physico-chemical properties. The soil samples were analyzed for pH, Electrical Conductivity (E.C.), % organic carbon, % organic matter. The

colour of soil samples were brown and dark gray. The pH of soil samples were ranged 6.22 to 7.31 slightly acidic to alkaline nature. High pH can be casues acidic soils. Similar results have been also reported by kumaret *al.*, 2014. The electrical conductivity (E.C.) value from 0.41 to 0.62 mmhos/cm. In the variations of electrical conductivity due to present of high concentration of ions. Same type results were reported by Rahi *et al.*, 2018. The percentages of organic matter and organic carbon in the soil samples were ranged from 0.17% to 0.25% and 0.29% to 0.43%. Due to high organic waste discharged with sewage water so organic carbon and organic matter % contents increases.Similar results reported by Singh et al., 2010. The amount of heavy metals zinc, iron, copper and manganese ranges were reported 1.10 to 1.81 ppm, 3.10 to 4.71 ppm, 0.25 to 0.55 ppm and 2.90 to 4.70 ppm. The amount of heavy metals in sewage water crossed their standard limit. Toxicity symptoms are reported in the sewage water. Similar results have been also reported Singh *et al.*, 2006.

Parameters	Critical limit
pH	7-8.5
Electrical Conductivity mmhos/cm	0-1.5
Organic Carbon %	0.5-0.75
Organic Matter %	0.8-1.29
Zinc ppm	0.6
Iron ppm	4.5
Copper ppm	0.2
Manganese ppm	2.0

**Table no.-2: Standard Critical limits of different parameters**

sites	pH	E.C.mmhos/cm	O.C.%	O.M.%	Zn Ppm	Fe ppm	Cu ppm	Mn Ppm
V <sub>1</sub>	7.17	0.51	0.20	0.34	1.15	4.35	0.30	3.10
V <sub>2</sub>	7.24	0.60	0.18	0.31	1.10	3.65	0.31	3.50
V <sub>3</sub>	7.10	0.53	0.17	0.29	1.45	3.10	0.52	2.90
V <sub>4</sub>	6.76	0.41	0.19	0.32	1.57	4.28	0.46	3.78
V <sub>5</sub>	6.35	0.57	0.24	0.41	1.42	3.50	0.44	4.55
V <sub>6</sub>	7.31	0.61	0.18	0.31	1.68	4.69	0.37	2.92

V <sub>7</sub>	6.37	0.53	0.21	0.36	1.32	4.71	0.55	4.10
V <sub>8</sub>	7.25	0.62	0.19	0.32	1.81	4.00	0.38	4.70
V <sub>9</sub>	6.22	0.51	0.25	0.43	1.62	3.85	0.40	2.98

**Tableno.-3:DfferentVillages of Sewage water and soil Results**

### Conclusion

This research has demonstrated that soil samples in Chomu, Sanganer, and Shahpura blocks have been affected by sewage water effluent, resulting in a noticeable increase in coloration, unpleasant odor, and a slightly acidic to alkaline pH. Furthermore, these soil samples contain trace amounts of metal ions at concentrations that do not meet the required standards. The study has also revealed that the effluent released from the sewage channel is significantly polluted. It is crucial to promptly implement appropriate methods for treating the effluent before it is discharged into surface water sources in order to mitigate the potential environmental risks it poses.

### References

- District Factbook. Rajasthan District Factbook Jaipur district. Key Socio-economic Data of Jaipur district, Rajasthan. District Profile – Krishi Vigyan Kendra, Jaipur 2019.
- Karn, S.K. and Harada, H. (2001): Surface water pollution in three urban territories of Nepal, India, and Bangladesh. *Environ. Manage*, 28(4): 483–496.
- Raikwar M.K., Kumar P., Singh M. and Singh A. (2008): Toxic effect of heavy metals in livestock health, *Vet World.*, 1(1): 28-30.
- Medona Mary, R., Nirmala, T. and Delphine Rose, M.R. (2015): Feeding guild and diversity of avifauna at sothuparai reservoir, Periyakulam, Theni district, Tamilnadu, India. *Int J RecentSci Res.*, 6(12): 7997-8001.
- Zhuang, P., Zou, H. and Shu, W. (2009): Biotransfer of heavy metals along a soil-plant insect-chicken food chain: *Field study. J. Environ. Sci.*, 21(6): 849-853.
- Mihaly-Cozmuta A, Mihaly-Cozmuta L, Viman V, Vatca G and Varga (2005). Spectrometric methods used to determine heavy metals and total cyanides in accidental polluted soils. *American Journal of Applied Sciences* 12(1) 358-362.
- Gitimoni Deka and Bhattacharjee KG (2009). Assesment of water quality in an area receiving effluent discharge from a textile mill. *Indian Journal of Environmental Protection* 29(6) 539-543.

- Yusuf RO and Sonibare JA (2004). Characterization of textile industries effluents in Kaduna, Nigeria and Olayinka, Studies on industrial pollution in Nigeria. The effects of textile effluents on the quality of ground water in some parts of Lagos. *Nigerian Journal of Health and Biomedical Sciences* 3 44-50.
- Ahire DV, Chaudhari PR, Ahire VD and Patil AA (2013). Correlations of Electrical Conductivity and Dielectric Constant with Physico-Chemical Properties of Black Soils. *International Journal of Scientific and Research Publications* 3(2) 1-16.
- Dengđz O (2010). Morphology, Physico-Chemical Properties and Classification of Soils on Terraces of the Tigris River in the South-east Anatolia Region of Turkey. *Journal of Agricultural Science* 16(206) 205-212.
- Kumar, Pradeep, Yadav, B. L., Rajput, S. G., Yadav, Brijesh, & Singh, K. (2014). Status of major nutrient in relation to soil properties of Jaipur district of Rajasthan under groundnut cultivation. *J. Soil Water Conservation*, 13(1), 31-35.
- Rahi, R. K., Prasad, R. N., & Gupta, V. (2018). Analysis of physico-chemical properties of textile effluents collected from Sanganer, Jaipur. *Int. J. Adv. Sci. Res. Manag*, 3(7), 119-123.
- Singh, M., Lodha, P., & Singh, G. P. (2010). Seasonal diatom variations with reference to physico-chemical properties of water of Mansagar Lake of Jaipur, Rajasthan. *Res J Agric Sci*, 1(4), 451-457.
- Singh, V., & Singh Chandel, C. P. (2006). Analytical study of heavy metals of industrial effluents at Jaipur, Rajasthan (India). *Journal of environmental science & engineering*, 48(2), 103–108.